



CryOcean-QCV

IOP Daily Data Quality report for 14/05/2014

| DOCUMENT INFORMATION | | | |
|----------------------|----------|-------------------|---------------|
| Document history | | | |
| Version | Issued | Reason for change | Author |
| 1.0r1 | 08/12/14 | Creation | F. M. Calafat |

| This Version | | |
|--|---------------|--|
| Version 1.0r1 – issued by National Oceanography Centre on 08/12/14 | | |
| Written by: | F. M. Calafat | |
| Checked by: | P. Cipollini | |
| Approved by: | | |
| Distribution List | | |
| Company | Delivered via | Names |
| ESA | ftp | Jerome Bouffard, EOP-GMQ Pierre Féménias, EOP-GMQ |
| NOC | ftp | P.Cipollini H.Snaith |



IOP DAILY QUALITY ASSESSMENT FOR 14/05/2014

Report issued on **08/12/14**

Date and time of the first record: **14 05 2014 00:00:00.505**

Date and time of the last record: **14 05 2014 23:54:23.658**

Range of complete orbits in present day: **21717 to 21730**

WARNINGS (see Appendix A for details)

orbit_dropout: 76.9% of records over ocean for orbit 21724

orbit_dropout: 51.2% of records over ocean for orbit 21727

orbit_dropout: 42.4% of records over ocean for orbit 21728

orbit_dropout: 54.1% of records over ocean for orbit 21729



1 Data latency

Median latency [range std]: 1.4 days [1.1 - 1.7]




Figure 1. Box-and-whiskers plot for the latency showing for each day in last 30-day window the first and third quartiles (bottom and top of the box), the median (thick black), the 5% and 95% percentiles (lower and upper whiskers), the mean (blue) and the standard deviation (blue dashed line). The percentage of records delivered within 3 days is also shown (red, right y-axis). The yellow shading highlights the day analyzed in this report whereas the horizontal black line denotes the 3 days threshold.




Figure 2. Histogram of the data latency for 14/05/2014.

2 Data coverage and completeness

| | Present in day | Theoretical max. | Percentage (%) |
|-------------------------|----------------|------------------|----------------|
| Total | 70235 | 91584 | 76.7 |
| Oceans and lakes | 49985 | 61035 | 81.9 |

Table 1. Number of total (land and ocean/lake) and only ocean/lake records together with their percentage relative to the theoretically expected number of measurements from the orbits ground tracks for 14/05/2014. Theoretical values are also shown.




Figure 3. Percentage of 1-Hz records over land and ocean/lake (red) and only over ocean/lake (blue) relative to the theoretically expected number from the orbits ground tracks for each day in last 30-day window. The black circle at the end of the time series highlights the day analyzed in this report. The values in parentheses denote linear trends and their standard errors.




Figure 4. Percentage of 1-Hz records over land and ocean/lake (red) and only over ocean/lake (blue) relative to the theoretically expected number from the orbits ground tracks for each orbit in last 30-day window. The yellow shading highlights the day analyzed in this report. The values in parentheses denote linear trends and their standard errors.

3 SSH anomaly coverage and validity

Note 3.1: unless otherwise stated, measurements taken over polar regions have been excluded from the computation of all statistics shown in this section.

Note 3.2: the extent of the regions defining the geographical mode mask of the reference mission is shown in Figure A1 (Appendix A).

Note 3.3: most statistics shown in this section have been computed separately for the low resolution mode (LRM) and the pseudo low resolution mode (PLRM).

3.1. Validity based on flags

Note 3.4: here “flag-valid” refers to those records that have not been flagged as bad by either the average status flag or the measurement confidence flag.

Number of flag-valid 1-Hz SSH records over oceans and lakes (including polar regions):
42786

Percentage of flag-valid 1-Hz SSH records over oceans and lakes (including polar regions) relative to theoretical maximum: **70.1%**




Figure 5. Geographical distribution of flag-valid SSH anomaly data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the SSH anomaly in cm and are calculated separately for LRM and PLRM regions. Measurements taken over polar regions have been excluded from the computation of the statistical values. The black lines mark the outer limit of the Arctic and Antarctic polar regions.




Figure 6. Percentage of flag-valid 1-Hz SSH records over ocean and lakes (including polar regions) relative to theory for each day in last 30-day window. The linear trend and its standard deviation is also shown.




Figure 7. Histogram of flag-valid SSH anomaly over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014.

3.2. Analysis of measurement noise

20-Hz measurement noise (average std of 20-Hz sample blocks) for the flag-valid SSH anomaly over oceans and lakes for LRM and PLRM, respectively: **9.1 cm** and **14.2 cm**
Equivalent 1-Hz measurement noise for the flag-valid SSH anomaly for LRM and PLRM, respectively: **2.0 cm** and **3.2 cm**




Figure 8. Geographical distribution of flag-valid 20-Hz SSH anomaly measurement noise over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the SSH anomaly noise and are calculated separately for LRM and PLRM regions. Measurements taken over polar regions have been excluded from the computation of the

statistical values. The black lines mark the outer limit of the Arctic and Antarctic polar regions.




Figure 9. 2D histogram showing flag-valid SSH anomaly noise as a function of SWH for LRM (top) and PLRM (bottom) for 14/05/2014. The black line denotes the median SSH anomaly as a function of SWH.



3.3. Validity based on scientific editing criteria

Note 3.5: here “science-valid” refers to the flag-valid SSH records over oceans and lakes (excluding polar regions) that meet the editing criteria described in Table 2.

| Parameter | Min threshold | Max threshold |
|---|---------------|---------------|
| SSH anomaly | -10 m | 10 m |
| Standard deviation of SSH anomaly (1-Hz block) | 0 m | 0.15 m |
| Inverse barometer correction | -2 m | 2 m |
| Biased orbit | -0.9 m | 0.9 m |

Table 2. Editing criteria. The last criteria in the table refers to the orbits highlighted in the 'Warnings' table at the beginning of this report as being suspicious of suffering from a significant orbit bias. For such orbits, any record having a value outside the range determined by the minimum and maximum thresholds is rejected.

Number of science-valid 1-Hz SSH records over oceans and lakes: **36897**

Percentage of science-valid 1-Hz SSH records over oceans and lakes relative to theoretical maximum: **60.5%**

20-Hz measurement noise (average std of 20-Hz sample blocks) for the science-valid SSH anomaly for LRM and PLRM, respectively: **7.7 cm** and **10.5 cm**

Equivalent 1-Hz measurement noise for the science-valid SSH anomaly for LRM and PLRM, respectively: **1.7 cm** and **2.3 cm**




Figure 10. Geographical distribution of science-valid SSH anomaly data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the SSH anomaly in cm and are calculated separately for LRM and PLRM regions. The black lines mark the outer limit of the Arctic and Antarctic polar regions. Gray circles denote records that have been rejected based on the editing criteria described in Table 2.




Figure 11. Histogram of science-valid SSH anomaly over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014.




Figure 12. Mean science-valid SSH anomaly for LRM (blue dot) and PLRM (red dot). The corresponding standard deviation (blue error bar and thin red lines, respectively) for each day in last 30-day window is also shown. The yellow shading highlights the day analyzed in this report.




Figure 13. Average science-valid SSH anomaly noise for LRM (blue dot) and PLRM (red dot). The corresponding standard deviation (blue error bar and thin red lines, respectively) for each day in last 30-day window is also shown. The yellow shading highlights the day analyzed in this report.

3.4. Crossover analysis




Figure 14. Mean crossover differences (absolute values) for the science-valid SSH anomaly (blue dot) and the corresponding standard deviation (error bar) for each day in last 30-day window. Mean crossover differences are calculated daily by averaging all crossover differences in a day (there are about 12-13 crossovers per day). The difference at each crossover is computed as the difference between median values over 2-s windows centered about the crossover. The yellow shading highlights the day analyzed in this report.

4 SWH coverage and validity

Note 4.1: unless otherwise stated, measurements taken over polar regions have been excluded from the computation of all statistics shown in this section.

Note 4.2: the extent of the regions defining the geographical mode mask of the reference mission is shown in Figure A1 (Appendix A).

Note 4.3: most statistics shown in this section have been computed separately for the low resolution mode (LRM) and the pseudo low resolution mode (PLRM).

4.1. Validity based on flags

Note 4.4: here “flag-valid” refers to those records that have not been flagged as bad by either the average status flag or the measurement confidence flag.

Number of flag-valid 1-Hz SWH records over oceans and lakes (including polar regions):
42809

Percentage of flag-valid 1-Hz SWH records over oceans and lakes (including polar regions) relative to theoretical maximum: **70.1%**




Figure 15. Geographical distribution of flag-valid SWH data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the SWH in m and are calculated separately for LRM and PLRM regions. Measurements taken over polar regions have been excluded from the computation of the statistical values. The black lines mark the outer limit of the Arctic and Antarctic polar regions.




Figure 16. Percentage of flag-valid 1-Hz SWH records over ocean and lakes (including polar regions) relative to theory for each day in last 30-day window. The linear trend and its standard deviation is also shown.




Figure 17. Histogram of flag-valid SWH over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014.

4.2. Analysis of measurement noise

20-Hz measurement noise (average std of 20-Hz sample blocks) for the flag-valid SWH over oceans and lakes for LRM and PLRM, respectively: **53.3 cm** and **82.6 cm**

Equivalent 1-Hz measurement noise for the flag-valid SWH for LRM and PLRM, respectively: **11.9 cm** and **18.5 cm**




Figure 18. Geographical distribution of flag-valid 20-Hz SWH measurement noise over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the



SWH noise and are calculated separately for LRM and PLRM regions. Measurements taken over polar regions have been excluded from the computation of the statistical values. The black lines mark the outer limit of the Arctic and Antarctic polar regions.

4.3. Validity based on scientific editing criteria

Note 4.5: here “science-valid” refers to the flag-valid SWH records over oceans and lakes that meet the editing criteria described in Table 3.

| Parameter | Min threshold | Max threshold |
|--|---------------|---------------|
| SWH | 0 m | 15 m |
| Standard deviation of SWH (1-Hz block) | 0 m | 1 m |

Table 3. Editing criteria.

Number of science-valid 1-Hz SWH records over oceans and lakes: **37435**

Percentage of science-valid 1-Hz SWH records over oceans and lakes relative to theoretical maximum: **61.3%**

20-Hz measurement noise (average std of 20-Hz sample blocks) for the science-valid SWH for LRM and PLRM, respectively: **49.6 cm** and **70.3 cm**

Equivalent 1-Hz measurement noise for the science-valid SWH for LRM and PLRM, respectively: **11.1 cm** and **15.7 cm**




Figure 19. Geographical distribution of science-valid SWH data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the SWH in m and are calculated separately for LRM and PLRM regions. The black lines mark the outer limit of the Arctic and Antarctic polar regions. Gray circles denote records that have been rejected based on the editing criteria described in Table 3.




Figure 20. Histogram of science-valid SWH over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014.




Figure 21. Box-and-whiskers plot for the science-valid SWH showing for each day in last 30-day window the first and third quartiles (bottom and top of the box), the median (thick black), the 5% and 95% percentiles (lower and upper whiskers), and the mean (blue). The yellow shading highlights the day analyzed in this report.




Figure 22. Average science-valid SWH noise for LRM (blue dot) and PLRM (red dot). The corresponding standard deviation (blue error bar and thin red lines, respectively) for each day in last 30-day window is also shown. The yellow shading highlights the day analyzed in this report.

5 Sigma0 coverage and validity

Note 5.1: unless otherwise stated, measurements taken over polar regions have been excluded from the computation of all statistics shown in this section.

Note 5.2: the extent of the regions defining the geographical mode mask of the reference mission is shown in Figure A1 (Appendix A).

Note 5.3: most statistics shown in this section have been computed separately for the low resolution mode (LRM) and the pseudo low resolution mode (PLRM).

5.1. Validity based on flags

Note 5.4: here “flag-valid” refers to those records that have not been flagged as bad by either the average status flag or the measurement confidence.

Number of flag-valid 1-Hz sigma0 records over oceans and lakes (including polar regions):
42839

Percentage of flag-valid 1-Hz sigma0 records over oceans and lakes (including polar regions) relative to theoretical maximum: **70.2%**




Figure 23. Geographical distribution of flag-valid sigma0 data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the sigma0 in dB and are calculated separately for LRM and PLRM regions. Measurements taken over polar regions have been excluded from the computation of the statistical values. The black lines mark the outer limit of the Arctic and Antarctic polar regions.




Figure 24. Percentage of flag-valid 1-Hz sigma0 records over ocean and lakes (including polar regions) relative to theory for each day in last 30-day window. The linear trend and its standard deviation is also shown.




Figure 25. Histogram of flag-valid sigma0 over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014.

5.2. Analysis of measurement noise

220-Hz measurement noise (average std of 20-Hz sample blocks) for the flag-valid SWH over oceans and lakes for LRM and PLRM, respectively: **12.1 10^{-2} dB** and **14.5 10^{-2} dB**

Equivalent 1-Hz measurement noise for the flag-valid SWH for LRM and PLRM, respectively: **2.7 10^{-2} dB** and **3.2 10^{-2} dB**




Figure 26. Geographical distribution of flag-valid 20-Hz sigma0 measurement noise over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the sigma0 noise and are calculated separately for LRM and PLRM regions. Measurements taken over polar regions have been excluded from the computation of the statistical values. The black lines mark the outer limit of the Arctic and Antarctic polar regions.

5.3. Validity based on scientific editing criteria

Note 5.5: here “science-valid” refers to the flag-valid sigma0 records over oceans and lakes that meet the editing criteria described in Table 4.

| Parameter | Min threshold | Max threshold |
|--|---------------|---------------|
| Sigma0 | 7 dB | 30 dB |
| Standard deviation of Sigma0 (1-Hz block) | 0 dB | 0.23 dB |

Table 4. Editing criteria.



Number of science-valid 1-Hz sigma0 records over oceans and lakes: **36302**

Percentage of science-valid 1-Hz sigma0 records over oceans and lakes relative to theoretical maximum: **59.5%**

20-Hz measurement noise (average std of 20-Hz sample blocks) for the science-valid sigma0 for LRM and PLRM, respectively: **$10.5 \cdot 10^{-2}$ dB** and **$10.7 \cdot 10^{-2}$ dB**

Equivalent 1-Hz measurement noise for the science-valid sigma0 for LRM and PLRM, respectively: **$2.4 \cdot 10^{-2}$ dB** and **$2.4 \cdot 10^{-2}$ dB**




Figure 27. Geographical distribution of science-valid sigma0 data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the sigma0 in dB and are calculated separately for LRM and PLRM regions. The black lines mark the outer limit of the Arctic and Antarctic polar regions. Gray circles denote records that have been rejected based on the editing criteria described in Table 4.




Figure 28. Histogram of science-valid sigma0 over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014.




Figure 29. Box-and-whiskers plot for science-valid sigma0 showing for each day in last 30-day window the first and third quartiles (bottom and top of the box), the median (thick black), the 5% and 95% percentiles (lower and upper whiskers), and the mean (blue). The yellow shading highlights the day analyzed in this report.




Figure 30. Average science-valid sigma0 noise for LRM (blue dot) and PLRM (red dot). The corresponding standard deviation (blue error bar and thin red lines, respectively) for each day in last 30-day window is also shown. The yellow shading highlights the day analyzed in this report.

6 Altimeter wind speed coverage and validity

Note 6.1: unless otherwise stated, measurements taken over polar regions have been excluded from the computation of all statistics shown in this section.

Note 6.2: the extent of the regions defining the geographical mode mask of the reference mission is shown in Figure A1 (Appendix A).

Note 6.3: most statistics shown in this section have been computed separately for the low resolution mode (LRM) and the pseudo low resolution mode (PLRM).

6.1. Validity based on flags

Note 6.4: here “flag-valid” refers to those records that have not been flagged as bad by either the average status flag or the measurement confidence flag.

Number of flag-valid 1-Hz wind records over oceans and lakes (including polar regions):
45655

Percentage of flag-valid 1-Hz wind records over oceans and lakes (including polar regions) relative to theoretical maximum: **74.8%**




Figure 31. Geographical distribution of flag-valid wind data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the wind in m/s and are calculated separately for LRM and PLRM regions. Measurements taken over polar regions have been excluded from the computation of the statistical values. The black lines mark the outer limit of the Arctic and Antarctic polar regions.




Figure 32. Percentage of flag-valid 1-Hz wind records over ocean and lakes (including polar regions) relative to theory for each day in last 30-day window. The linear trend and its standard deviation is also shown.




Figure 33. Histogram of flag-valid wind over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014.

6.2. Validity based on scientific editing criteria

Note 6.5: here “science-valid” refers to the flag-valid wind records over oceans and lakes that meet the editing criteria described in Table 5.

| Parameter | Min threshold | Max threshold |
|----------------------|---------------|---------------|
| Altimeter wind speed | 0 m/s | 30 m/s |

Table 5. Editing criteria.

Number of science-valid 1-Hz wind records over oceans and lakes: **38424**

Percentage of science-valid 1-Hz wind records over oceans and lakes relative to theoretical maximum: **63.0%**




Figure 34. Geographical distribution of science-valid wind data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the wind in m/s and are calculated separately for LRM and PLRM regions. The black lines mark the outer limit of the Arctic and Antarctic polar regions. Gray circles denote records that have been rejected based on the editing criteria described in Table 5.




Figure 35. Histogram of science-valid wind over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014.




Figure 36. Box-and-whiskers plot for the science-valid wind showing for each day in last 30-day window the first and third quartiles (bottom and top of the box), the median (thick black), the 5% and 95% percentiles (lower and upper whiskers), and the mean (blue). The yellow shading highlights the day analyzed in this report.

7 Mispointing coverage and validity

Note 7.1: unless otherwise stated, measurements taken over polar regions have been excluded from the computation of all statistics shown in this section.

Note 7.2: the extent of the regions defining the geographical mode mask of the reference mission is shown in Figure A1 (Appendix A).

Note 7.3: most statistics shown in this section have been computed separately for the low resolution mode (LRM) and the pseudo low resolution mode (PLRM).

Note 7.4: here “flag-valid” refers to those records that have not been flagged as bad by either the average status flag or the measurement confidence flag.

Number of flag-valid 1-Hz mispointing records over oceans and lakes (including polar regions): **45655**

Percentage of flag-valid 1-Hz mispointing records over oceans and lakes (including polar regions) relative to theoretical maximum: **74.8%**

Percentage of positive flag-valid 1-Hz mispointing records over oceans and lakes (including polar regions) relative to theoretical maximum: **63.6%**




Figure 37. Geographical distribution of flag-valid mispointing data over oceans and lakes for 14/05/2014. The statistical values shown in the table refer to the mispointing in 10^{-2} deg^2 and are calculated separately for LRM and PLRM regions. Measurements taken over polar regions have been excluded from the computation of the statistical values. The black lines mark the outer limit of the Arctic and Antarctic polar regions.




Figure 38. Percentage of flag-valid 1-Hz mispointing records over ocean and lakes (including polar regions) relative to theory for all values (blue) and only positive values (red) for each day in last 30-day window. The linear trend and its standard deviation is also shown for each time series.




Figure 39. Histogram of flag-valid mispointing over oceans and lakes for LRM (blue) and PLRM (red) for 14/05/2014. Only positive values have been considered.




Figure 40. Box-and-whiskers plot for flag-valid mispointing showing for each day in last 30-day window the first and third quartiles (bottom and top of the box), the median (thick black), the 5% and 95% percentiles (lower and upper whiskers), and the mean (blue). The yellow shading highlights the day analyzed in this report. Only positive values have been considered.

APPENDIX A

In this appendix we briefly describe the meaning of the warning codes summarized in page 2 of the document.

- **latency_fail:** this warning is issued whenever one or more data files for the day analyzed in this report are made available with a delay of more than 3 days with respect to the mean time of the records stored in the file.
- **latency_mean_high:** this warning is issued whenever the mean of the latencies over all data files for the day analyzed in this report is larger than 2 days.
- **ocean_dropout:** this warning is issued whenever the percentage of 1-Hz records over ocean and lakes relative to the theoretically expected number per day is smaller than 80% for the day analyzed in this report.
- **orbit_dropout:** this warning refers to individual orbits and is issued whenever the percentage of 1-Hz records over oceans and lakes relative to the theoretically expected number per orbit is smaller than 80% for one or more orbits for the day analyzed in this report.
- **large_orbit_bias:** this warning refers to individual orbits and is issued whenever there are more than 100 flag-valid records with a SSH anomaly larger than 90 cm.




Figure A1. Extent of the regions defining the geographical mode mask for 14/05/2014. The regions where SAR, and SARIn are operated are denoted by the orange and blue polygons, respectively. LRM is operated over areas of the continental ice sheets (red polygons) and over oceans and land not covered by other modes.

(end of Document)