


---

## **ENVISAT GOMOS report: August 2009**



---

Prepared by:	L. Saavedra de Miguel - SERCO
Approved by:	Angelika Dehn - SERCO
Inputs from:	GOMOS Quality Working Group, ECMWF
Issue:	1.0
Reference:	ENVI-SPPA-EOPG-TN-09-0032
Date of issue:	14 <sup>th</sup> September 2009
Status:	Reviewed
Document type:	Technical Note

# **TABLE OF CONTENTS**

<b>1 INTRODUCTION.....</b>	<b>3</b>
1.1 Scope .....	3
1.2 References.....	3
1.3 Acronyms and Abbreviations.....	4
<b>2 SUMMARY .....</b>	<b>6</b>
<b>3 INSTRUMENT AND DATA AVAILABILITY.....</b>	<b>8</b>
3.1 GOMOS Unavailability Periods.....	8
3.2 Stars Lost in Centering.....	8
3.3 Stars lost due to VCCS anomaly.....	10
3.4 Data Generation Gaps.....	10
3.5 Data availability to users.....	11
<b>4 INSTRUMENT CONFIGURATION AND PERFORMANCE.....</b>	<b>12</b>
4.1 Instrument Operation and Configuration.....	12
4.1.1 Operations since beginning of mission .....	12
4.1.2 Current operations and configuration.....	15
4.2 Limb, Illumination conditions and instrument gain setting.....	16
4.3 Thermal Performance.....	17
4.4 Optomechanical Performance.....	21
4.5 Electronic Performance.....	22
4.5.1 Dark Charge Evolution and Trend.....	22
4.5.2 Signal Modulation .....	25
4.5.3 Electronic Chain Gain and Offset.....	26
4.6 Acquisition, Detection and Pointing Performance.....	27
4.6.1 SATU Noise Equivalent Angle .....	27
4.6.2 Tracking Loss Information.....	29
4.6.3 Most Illuminated Pixel (MIP).....	32
<b>5 LEVEL 1 PRODUCT QUALITY MONITORING .....</b>	<b>33</b>
5.1 Processor Configuration.....	33
5.1.1 Version.....	33
5.1.2 Auxiliary Data files (ADF).....	36
5.2 Quality Flags Monitoring.....	39
5.2.1 Quality Flags Monitoring (extracted from Level 2 products).....	41
5.3 Spectral Performance.....	45
5.4 Radiometric Performance.....	45
5.4.1 Radiometric Sensitivity.....	45
5.4.2 Pixel Response Non Uniformity.....	48
5.5 Other Calibration Results.....	48
<b>6 LEVEL 2 PRODUCT QUALITY MONITORING .....</b>	<b>48</b>
6.1 Processor Configuration.....	48
6.1.1 Version.....	48
6.1.2 Auxiliary Data Files (ADF).....	50
6.1.3 Re-Processing Status.....	51
6.2 Quality Flags Monitoring.....	52
6.3 Other Level 2 Performance Issues .....	53
6.3.1 Monthly Ozone average.....	53
6.3.2 Ozone dispersion monitoring .....	54

**7 VALIDATION ACTIVITIES AND RESULTS ..... 56**  
7.1 GOMOS-ECMWF Comparisons (Rossana Dragani, ECMWF input)..... 56

**APPENDIX A ..... 58**

# 1 INTRODUCTION

The GOMOS monthly report documents the current status and recent changes to the GOMOS instrument, its data processing chain, and its data products.

The Monthly Report (hereafter MR) is composed of analysis results obtained by the Data Processing and Quality Control, combined with inputs received from the different entities working on GOMOS operation, calibration, product validation and data quality. These teams participate in the GOMOS Quality Working Group:

- European Space Agency (ESRIN, ESOC, ESTEC-PLSO)
- IDEAS
- ACRI
- Service d'Aeronomie
- Finnish Meteorological Institute
- IASB-Belgian Institute for Space Aeronomy
- Astrium Space
- ECMWF

In addition, the group interfaces with the Atmospheric Chemistry Validation Team.

## 1.1 *Scope*

The main objective of the Monthly Report is to give, on a regular basis, the status of GOMOS instrument performance, data acquisition, results of anomaly investigations, calibration activities and validation campaigns. The following six sections compose the MR:

- Summary
- Unavailability
- Instrument Configuration and Performance
- Level 1 Product Quality Monitoring
- Level 2 Product Quality Monitoring
- Validation Activities and Results

## 1.2 *References*

- [1] ENVISAT Weekly Mission Operations Report #368, #369, #370, #371 and #372
- [2] ECMWF GOMOS Monthly Reports
- [3] Routine update of the wavelength assignment, Gilbert Barrot (ACRI-ST), Issue 1 Revision 1, September 19, 2007

### 1.3 *Acronyms and Abbreviations*

ACVT	Atmospheric Chemistry Validation Team
ADC	Analogue-to-Digital Converter
ADF	Auxiliary Data File
ADS	Auxiliary Data Server
ANX	Ascending Node Crossing
AOCS	Attitude and Orbit Control System
ARB	Anomaly Review Board
ARF	Archiving Facility (PDS)
CCU	Central Communication Unit
CFS	CCU Flight Software
CNES	Centre National d'Études Spatiales
CTI	Configuration Table Interface / Configurable Transfer Item
CR	Cyclic Report
DC	Dark Charge
DMOP	Detailed Mission Operation Plan
DPM	Detailed Processing Model
DS	Data Server
DSA	Dark Sky Area
DSD	Data Set Descriptor
ECMWF	European Centre for Medium Weather Forecast\
EO	Earth Observation
EQSOL	Equipment Switch Off Line
ESA	European Space Agency
ESL	Expert Support Laboratory
ESRIN	European Space Research Institute
ESTEC	European Space Research & Technology Centre
ESOC	European Space Operations Centre
FCM	Fine Control Mode
FinCoPAC	Finnish Products Archiving Center
FMI	Finnish Meteorological Institute
FOCC	Flight Operations Control Centre (ENVISAT)
FP1	Fast Photometer 1
FP2	Fast Photometer 2
GADS	Global Annotations Data Set
GOMOS	Global Ozone Monitoring by Occultation of Stars
GOPR	Gomos Prototype
GS	Ground Segment
HK	Housekeeping
IASB	Institut d'Aeronomie Spatiale de Belgique
IAT	Interactive Analysis Tool
ICU	Instrument Control Unit
IDEAS	Instrument Data quality Evaluation and Analysis
IDL	Interactive Data Language
IECF	Instrument Engineering and Calibration Facilities
IMK	Institute of Meteorology Karlsruhe (Meteorologisch Institut Karlsruhe)
INV	Inventory Facilities (PDS)

IPF	Instrument Processing Facilities (PDS)
JPL	Jet Propulsion Laboratory
LAN	Local Area Network
LMA	Levenberg-Marquardt Algorithm
LPCE	Laboratoire de Physique et Chimie de l'Environnement
LRAC	Low Rate Archiving Center
LUT	Look Up Table
MCMD	Macro Command
MDE	Mechanism Drive Electronics
MIP	Most Illuminated Pixel
MPH	Main Product Header
MPS	Mission Planning System
MR	Monthly Report
NRT	Near Real Time
OBDAH	On-Board Data Handling
OBT	On Board Time
OCM	Orbit Control Manoeuvre
OOP	Out-of-plane
OP	Operational Phase of ENVISAT
PAC	Processing and Archiving Centre (PDS)
PCF	Product Control Facility
PDCC	Payload Data Control Centre (PDS)
PDHS	Payload Data Handling Station (PDS)
PDHS-E	Payload Data Handling Station – ESRIN
PDHS-K	Payload Data Handling Station – Kiruna
PDS	Payload Data Segment
PEB	Payload Equipment Bay
PLSOL	Payload Switch off Line
PMC	Payload Module Computer
PRNU	Pixel Response Non Uniformity
PSO	On-Orbit Position
QC	Quality Control
QUARC	Quality Analysis and Reporting Computer
QWG	Quality Working Group
RDV	RenDez-Vous
RGT	ROP Generation Tool
RIVM	Rijksinstituut voor Volksgezondheid en Milieu
ROP	Reference Operations Plan
RRM	Rate Reduction Mode
RTS	Random Telegraphic Signal
SA	Service d'Aeronomie
SAA	South Atlantic Anomaly
SATU	Star Acquisition and Tracking Unit
SFA	Steering Front Assembly
SFCM	Stellar Fine Control Mode
SFM	Steering Front Mechanism
SM	Service Module
SMNA	Servicio Meteorológico Nacional de Argentina

SMP	Set Measurement Parameter
SODAP	Switch On and Data Acquisition Phase
SPA1	Spectrometer A CCD 1
SPA2	Spectrometer A CCD 2
SPB1	Spectrometer B CCD 1
SPB2	Spectrometer B CCD 2
SPH	Specific Product Header
SQADS	Summary Quality Annotation Data Set
SSP	Sun Shade Position
STP	Set Thermal Parameter
SYSTEM	Stellar Yaw Steering Mode
SZA	Solar Zenith Angle
VCCS	Voice Coil Command Saturation

## 2 SUMMARY

**Instrument availability** (section 3.1): There were no instrument unavailability periods during the reporting month.

**Instrument operations** (section 4.1.2): Since 17<sup>th</sup> July 2009 the instrument is operated in an azimuth window of [25, 50] degrees due to the less occurrences of VCCS anomalies. No monitoring sequence was planned for the reporting month.

**Voice-Coil Command Saturation (VCCS) Anomaly** (section 3.3): No Voice Coil Command Saturation anomalies occurred during the reporting month.

**Data availability** when instrument was in operation (section 3.4): The availability of Level 0 and Level 1b was higher than 99% for the whole month.

**Data availability for users** (section 3.5): Routine dissemination of Level 1b and Level 2 products produced by the PDS to the users is enabled. Level 1b data are available on request to the EO Helpdesk ([EOHelp@esa.int](mailto:EOHelp@esa.int)), while level 2 data are available for the whole mission on different ftp sites. All data (reprocessed, NRT and consolidated) are processed with the same version of GOMOS processor.

**Wavelength monitoring** (section 5.3): the wavelength shifts show a variation which was not expected after the implementation of the routine calibration on 14<sup>th</sup> December 2007. The QWG has been informed and is investigating this issue.

**Pointing performance** (section 4.6.1):

- **Sudden increase on September 2005:** as can be seen in fig. 4-6.1, the SATU NEA had a sudden increase on 8<sup>th</sup> September 2005 mainly in ‘Y’ axis. These values remained high, fluctuating between 1 and 1.8 microrad until December 2005 when they came back to the values they used to be before the increase of September. The reason why there was higher noise in the data causing the jump in daily SATU average is not known.
- **Gradual increase on mid April 2006:** a different problem was present since mid April 2006 until October 2007. A gradual increase of the daily SATU ‘Y’ mean was observed. This increase was due to fluctuations of the SATU ‘Y’ data observed at the beginning of nominal occultations

(starting at 130 km that corresponds to an elevation angle of around 65°). The decrease of the start elevation angle of the occultation has no impact on the amplitude of the SATU 'Y' fluctuations. Investigations carried out by the ESL, ESA and industry pointed to a problem on the SFM (mechanical or electrical) and not to a problem on the SATU itself. Since October 2007 the fluctuations have disappeared and as a consequence the daily SATU 'Y' average has come back below the threshold set to 3 micro radians.

- **Current anomaly: sudden increase on December 2008:** similarly to the anomaly happened on April 2006, the SATU NEA had an increase on 29<sup>th</sup> December 2008 due to fluctuations of the SATU 'Y' data. The difference with respect to the previous anomaly is that this time, the increase was quite sudden and the fluctuations are present during the whole occultation, not only at the beginning of the occultation. The most critical effect of this anomaly is the loss of the star measurement high in the atmosphere, which means that many times the corresponding ozone profiles do not include the ozone peak present at around 25-30 km. After the increase of the elevation filter from 100 to 150 on 29<sup>th</sup> June 2009, the abortion of the star measurements is now deeper in the atmosphere but still premature. A new increase on 6<sup>th</sup> July (from 150 to 200) did not produce an immediate improvement but it seems that after that date a gradual improvement took place.

**Temperatures** (section 4.3): The CCD temperatures show the expected global increase due to the radiator ageing. Another expected variation of the temperatures, the seasonal one, with amplitude of around 1.5 degree can also be observed.

**Modulation signal** (section 4.5.2): The values of the modulation are daily extracted and plotted; they should not be very different from the ones coded into the processor: 1.40 ADU for SPA1 and 0.76 ADU for SPA2. The modulation signal shows high values during summer time for the ESRIN data, it now being confirmed that the South Atlantic Anomaly is the cause of these unexpected peaks. The quality of ESRIN data, in particular over the SAA zone, is impacted but the measure of this impact is under investigation. However, in the second half of the months of October of all years (2004-2008) the peaks are smaller because the DSA zone where the data are taken for this analysis is moving towards the Northern Hemisphere. At the end of October the DSA zone is definitely chosen by the planning system in the Northern Hemisphere (to fill the criteria 'DSA in full dark limb conditions') and the high peaks disappear.

**Star detection performance** (section 4.6.3): the stars should be detected not far from the SATU center, that is, pixel number 145 in elevation and number 205 in azimuth. The elevation MIP (Most Illuminated Pixel, which is the pixel at the moment of the detection) had a significant variation until 12<sup>th</sup> December 2003 when a new PSO algorithm was activated in order to reduce the deviations of the ENVISAT platform attitude with respect to the nominal one. Afterwards, the MIP position was quite stable around its nominal pixel values until the occurrence of the VCCS anomaly on January 2005. The reason for the change in trend observed after the anomaly is, at the moment, not understood. This behavior, currently stable at pixel 127 in elevation and 193 in azimuth, does not impact the data quality but may invalidate attitude monitoring by GOMOS and could represent a hidden anomaly.

**Radiometric sensitivity monitoring** (section 5.4.1): for stars 25 and 9, the UV ratio is greater than the threshold 10%. It is clear that there is a global decrease of UV ratios for all the stars. This confirms the expected degradation suffered by the UV optics that is, anyway, very small considering also the small variation for the rest of the stars. For the photometers radiometric sensitivity ratios it is observed that every star has a variation that seems to be seasonally related. The variation is significant for stars 25 and



18. After some investigations performed by the QWG that exclude an inaccurate reflectivity correction LUT, it seems that the PH1/2 radiometric sensitivity variations could come from the fact that the spectrometers and the photometers are not illuminated the same way when the straylight appears.

**Auxiliary Data File** (sections 5.1.2 and 5.3): Four GOM\_CAL\_AX files with updated DC maps and new wavelength assignment have been disseminated during the reporting period.

### 3 INSTRUMENT AND DATA AVAILABILITY

#### 3.1 GOMOS Unavailability Periods

No unavailability periods were recorded for the reporting month.

**Table 3.1-1: List of unavailability periods issued during the reporting month**

Reference of unavailability report	Start time Star orbit	Stop time Stop orbit	Description
-	-	-	-

#### 3.2 Stars Lost in Centering

The acquisition of a star initiates with a rallying phase where the telescope mechanism is directed towards the expected position of the star. Subsequently the acquisition procedure enters into detection mode, where the SATU star tracker output signal is pre-processed for spot presence survey and for the location of the most illuminated couple of adjacent pixels for two added lines, over the detection field. The Most Illuminated Pixel (MIP) defines the position of the first SATU centering window. The following step in the acquisition sequence is then initiated and consists of a centering phase where the SATU output signal is pre-processed for spot presence survey over the maximum of 10x10 pixel field. This allows the third phase to begin: the tracking phase.

The centering phase has occasionally resulted in loss of the star from the field of view. Fig. 3.2-1 reports the percentage of the stars lost in centering for the period 3<sup>rd</sup> February 2003 to 30<sup>th</sup> August 2009. It can be seen that only two stars, mainly weak stars (higher star id means higher magnitude) are lost during the centering phase between 4% and 9 % of their planned observations. The majority of those are geo-localized over the SAA.

As the monitoring shows neither a trend nor excessively high percentages of loss, there is no need for the moment to reject any star from the catalogue, and there is no indication of instrument-related problems. Now with the instrument in a new operation scenario, the stars could be also lost due to the anomaly “elevation voice coil command saturation” even if the instrument is not going anymore to Stand by / Refuse mode (section 3.3).

### Statistics on stars lost in centering: 03-FEB-2003 until 30-AUG-2009




Figure 3.2-1: Statistics on stars that have been lost during the centering phase. The number above the columns corresponds to the Star ID

### 3.3 Stars lost due to VCCS anomaly

No VCCS anomalies occurred during the reporting month. The information provided in table 3.3-1 (when available) is:

- **UTC anomaly:** the UTC of the anomaly occurrence
- **Star from:** star id of the last successful occultation before the anomaly occurrence
- **Star to:** star id of the star to be occulted when the anomaly occurred
- **Az. star from:** the start azimuth of the “Star from”
- **Az. star to:** the start azimuth of the “Star to”
- **Diff:** the azimuth angle difference between the last azimuth of “Star from” and the start azimuth of the “Star to”
- **Nb stars:** number of consecutive stars lost due to the anomaly

Table 3.3-1: VCCS Anomaly occurred during the reporting period

UTC Anomaly	Star from	Star to	Az. Star from	Az. Star to	Diff	Nb stars
-	-	-	-	-	-	-

### 3.4 Data Generation Gaps

The trend in percentage of available NRT data within the archives PDHS-K and PDHS-E is depicted in fig. 3.4-1 (when instrument was in operation). It is a good indicator on how the PDS chain is working in terms of generation and dissemination of data to the archives. The percentage is calculated once per week.

The availability of Level 0 and Level 1b was higher than 99% for the whole month.




Figure 3.4-1: Percentage of level 0 and level 1b data availability on the archives PDHS-E and PDHS-K

Occultations planned to be acquired but for which no GOM\_NL\_\_0P data product has become available are presented in fig. 3.4-2 for the reporting period.




Figure 3.4-2: The red points are the occultation geo-location (starting) corresponding to planned data acquisitions for which no GOMOS level 0 product has become available

### 3.5 Data availability to users

Routine dissemination of higher-level products produced by the PDS to the users is enabled. Level 1b data are available on request to the EO Helpdesk ([EOHelp@esa.int](mailto:EOHelp@esa.int)), while level 2 data are available for the whole mission. For information on the passwords, please, contact the EO Helpdesk ([EOHelp@esa.int](mailto:EOHelp@esa.int)):

- Reprocessed products GOM\_NL\_\_2P are available at the **D-PAC ftp server** (name: **ftp-ops-dp.eo.esa.int**, IP-Address: **195.37.183.37**):  
<ftp://gomo2usr@ftp-ops-dp.eo.esa.int> from **August 2002 to 4<sup>th</sup> July 2006**.
- Near Real Time products GOM\_NL\_\_2P (generated three hours after sensing time) are available on the following servers:  
<ftp://gomosusr@oa-es.eo.esa.int> (ESRIN data). **A seven-day rolling archive has been set-up on this server.**  
<ftp://gomosusr@oa-ks.eo.esa.int> (KIRUNA data). **A seven-day rolling archive has been set-up on this server.**
- Consolidated products GOM\_NL\_\_2P (generated three weeks after sensing time) are available at D-PAC ftp server  
<ftp://gomo2usr@ftp-ops-dp.eo.esa.int> since **23 July 2006**

All data (reprocessed, NRT and consolidated) are processed with the same version of GOMOS processor.

## 4 INSTRUMENT CONFIGURATION AND PERFORMANCE

### 4.1 Instrument Operation and Configuration

#### 4.1.1 OPERATIONS SINCE BEGINNING OF MISSION

GOMOS has had different operational scenarios during the mission:

- **End of March 2003 to July 2003:** during this period the azimuth range had to be decreased in steps (table 4.1-1) to avoid an instrument problem (“Voice\_coil\_command\_saturation” anomaly) that caused GOMOS to go into STAND BY/REFUSE mode
- **July 2003:** the driver assembly was switched to the redundant B-side and since that date the full azimuth range (-10.8, +90.8) was again available
- **25<sup>th</sup> January 2005:** A second major anomaly occurred. Between this date and until the instrument was declared operational again (29<sup>th</sup> August 2005), GOMOS has been operated for testing and anomaly investigation purposes in different operation scenarios.
- **29<sup>th</sup> August 2005:** GOMOS operational again with reduced azimuth window of 20 degrees
- **9<sup>th</sup> October 2005:** azimuth window moved from 20 to 25 degrees
- **12<sup>th</sup> March 2006:** the reduced azimuth window of 25 degrees becomes a sliding window
- **2<sup>nd</sup> February 2008:** azimuth window moved from 25 to 30 degrees
- **21<sup>st</sup> August 2008:** minimum allowed azimuth angle set to +2 degrees

The changes in azimuth configuration during the whole mission until end of reporting period are summarized in table 4.1-1.

**Table 4.1-1: Historical changes in Azimuth configuration when GOMOS is in operations**

Date	Orbit	Minimum Azimuth (°)	Maximum Azimuth (°)	Comment
01-MAR-2002		-10.8	+90.8	Nominal
29-MAR-2003 17:40	5635	0.0	+90.8	Reduced
31-MAY-2003 06:22	6530	+4.0	+90.8	Reduced
16-JUN-2003 16:17	6765	+12.0	+90.8	Reduced
15-JUL-2003 01:39	7200	-10.8	+90.8	Nominal
25-JAN-2005 23:33	15200	tests	tests	Different configurations for testing purposes
29-AUG-2005 02:52	18280	-10	+10	Reduced
26-SEP-2005 01:32	18680	-5	+20	Reduced
03-OCT-2005 01:12	18780	-5	+15	Reduced
09-OCT-2005 21:30	18878	-5	+20	Reduced
12-MAR-2006 17:29	21080	+10	+35	Reduced
09-APR-2006 12:47	21480	+5	+30	Reduced
16-APR-2006 15:48	21580	0	+25	Reduced
30-APR-2006 15:08	21780	-5	+20	Reduced
07-MAY-2006 14:48	21880	0	+25	Reduced
14-MAY-2006 14:28	21980	+15	+40	Reduced
28-MAY-2006 13:47	22180	+20	+45	Reduced
04-JUN-2006 13:27	22280	+15	+40	Reduced
18-JUN-2006 12:47	22480	+20	+45	Reduced
25-JUN-2006 12:27	22580	0	+25	Reduced
02-JUL-2006 12:07	22680	-5	+20	Reduced
16-JUL-2006 11:27	22880	0	+25	Reduced
23-JUL-2006 11:07	22980	+10	+35	Reduced

06-AUG-2006 10:26	23180	0	+25	Reduced
27-AUG-2006 09:26	23480	+5	+30	Reduced
03-SEP-2006 09:06	23580	0	+25	Reduced
10-SEP-2006 08:46	23680	-5	+20	Reduced
01-OCT-2006 07:45	23980	+5	+30	Reduced
15-OCT-2006 07:05	24180	-5	+20	Reduced
22-OCT-2006 06:45	24280	0	+25	Reduced
29-OCT-2006 06:25	24380	-5	+20	Reduced
05-NOV-2006 06.05	24480	10	+35	Reduced
12-NOV-2006 05.45	24580	5	+30	Reduced
03-DEC-2006 04.44	24880	20	+45	Reduced
10-DEC-2006 04.24	24980	10	+35	Reduced
17-DEC-2006 20.50	25090	0	+25	Reduced
24-DEC-2006 03.44	25180	5	+30	Reduced
07-JAN-2007 03.04	25380	0	+25	Reduced
14-JAN-2007 02.44	25480	-5	+20	Reduced
21-JAN-2007 02.23	25580	0	+25	Reduced
28-JAN-2007 02.03	25680	-5	+20	Reduced
04-FEB-2007 01.43	25780	-10	+15	Reduced
11-FEB-2007 01.23	25880	-5	+20	Reduced
18-FEB-2007 01.03	25980	0	+25	Reduced
25-FEB-2007 00.43	26080	+5	+30	Reduced
04-MAR-2007 00.23	26180	+15	+40	Reduced
11-MAR-2007 00.03	26280	+20	+45	Reduced
24-MAR-2007 23.22	26480	0	+45	Reduced
31-MAR-2007 23.02	26580	+5	+30	Reduced
07-APR-2007 22.42	26680	+10	+35	Reduced
14-APR-2007 22.22	26780	+5	+30	Reduced
21-APR-2007 22.02	26880	0	+25	Reduced
28-APR-2007 21.42	26980	-5	+20	Reduced
12-MAY-2007 21.02	27180	20	+45	Reduced
19-MAY 2007 20.41	27280	+10	+35	Reduced
09-JUN-2007 19.41	27580	+15	+40	Reduced
16-JUN-2007 19.21	27680	-5	+20	Reduced
23-JUN-2007 19.01	27780	0	+25	Reduced
07-JUL-2007 18.21	27980	-5	+20	Reduced
04-AUG-2007 17:00	28380	0	+25	Reduced
11-AUG-2007 16.40	28480	5	+30	Reduced
18-AUG-2007 16.20	28580	0	+25	Reduced
26-AUG-2007 16.00	28680	10	+35	Reduced
04-SEP-2007 04.01	28816	+65	+90	Reduced: SATU-Y test
05-SEP-2007 06.51	28832	+10	+35	Reduced
08-SEP-2007 15.19	28880	+15	+40	Reduced
15-SEP-2007 14.59	28980	+20	+45	Reduced
22-SEP- 2007 14.39	29080	-5	+15	Reduced
29-SEP-2007 14.19	29180	+5	+30	Reduced
13-OCT-2007 13.39	29378	10	+35	Reduced
20-OCT-2007 13.19	29480	0	+30	Reduced
24-OCT-2007 01.09	29530	0	+25	Reduced
27-OCT- 2007 12.59	29580	10	+35	Reduced
10-NOV-2007 12.18	29780	-5	+20	Reduced
17-NOV-2007 11.58	29880	0	+25	Reduced
24-NOV-2007 11.38	29980	+5	+30	Reduced
01-DEC-2007 11.18	30080	+15	+40	Reduced
08-DEC- 2007 10.58	30180	+10	+35	Reduced
11-DEC- 2007 22.48	30230	+5	+35	Reduced

15-DEC- 2007 10.38	30280	+5	+30	Reduced
22-DEC- 2007 10.18	30380	0	+25	Reduced
05-JAN-2008 09.37	30580	-1	+24	Reduced
12-JAN-2008 09.17	30680	-2	+23	Reduced
19-JAN-2008 08.57	30780	-7	+18	Reduced
26-JAN-2008 08.37	30880	-2	+23	Reduced
02-FEB-2008 08.17	30980	-6	+24	Reduced
16-FEB-2008 07.37	31180	-8	+22	Reduced
23-FEB-2008 07.17	31280	-2	+28	Reduced
01-MAR-2008 06.56	31380	+5	+35	Reduced
08-MAR-2008 06.36	31480	+13	+43	Reduced
15-MAR-2008 06.16	31580	+10	+40	Reduced
22-MAR-2008 16.00	31686	+14	+44	Reduced
29-MAR-2008 05.36	31780	-1	+29	Reduced
05-APR-2008 05:16	31880	-8	+22	Reduced
12-APR-2008 04:56	31980	-4	+26	Reduced
19-APR-2008 04:36	32080	-10	+20	Reduced
03-MAY-2008 03:55	32280	-5	+25	Reduced
10-MAY-2008 03:35	32380	-6	+24	Reduced
17-MAY-2008 03:15	32480	+9	+39	Reduced
24-MAY-2008 02:55	32580	+14	+44	Reduced
31-MAY-2008 12:39	32686	+16	+46	Reduced
07-JUN-2008 02.15	32780	+18	+48	Reduced
14-JUN-2008 01.55	32880	+5	+35	Reduced
21-JUN-2008 01.35	32980	+6	+36	Reduced
28-JUN-2008 01.14	33080	-2	+28	Reduced
05-JUL-2008 00.54	33180	-10	+20	Reduced
19-JUL-2008 00.14	33380	0	+30	Reduced
25-JUL-2008 23.54	33480	+5	+35	Reduced
01-AUG-2008 23.34	33580	-1	+29	Reduced
08-AUG-2008 23.14	33680	-3	+27	Reduced
15-AUG-2008 22.54	33780	+12	+42	Reduced
23-AUG-2008 08.37	33886	+5	+35	Reduced
29-AUG-2008 22.13	33980	+4	+34	Reduced
05 -SEP- 2008 21.53	34080	+6	+36	Reduced
12 -SEP- 2008 21.33	34180	+15	+45	Reduced
27 -SEP- 2008 06.56	34386	+4	+34	Reduced
03-OCT-2008 20.33	34480	+7	+37	Reduced
10-OCT-2008 20.13	34580	+4	+34	Reduced
17-OCT-2008 19.53	34680	+2	+32	Reduced
01-NOV-2008 05.16	34886	+3	+33	Reduced
07-NOV-2008 18.52	34980	+5	+35	Reduced
14-NOV-2008 18.32	35080	+40	+70	Reduced
28-NOV-2008 17.52	35280	+25	+55	Reduced
06-DEC-2008 03.35	35686	+17	+47	Reduced
12-DEC-2008 17.12	35480	+14	+44	Reduced
19-DEC-2008 16.51	35580	+10	+40	Reduced
26-DEC-2008 16.31	35680	+6	+36	Reduced
02-JAN-2009 16.11	35780	+3	+33	Reduced
10-JAN-2009 01.55	35886	+4	+34	Reduced
16-JAN-2009 15.31	35980	+2	+32	Reduced
12-FEB-2009 04.39	36360	+3	+23	Testing
12-FEB-2009 08.00	36362	+20	+40	Testing
12-FEB-2009 11.21	36364	+35	+55	Testing
12-FEB-2009 14.42	36366	+50	+70	Testing
12-FEB-2009 18.03	36368	+65	+85	Testing

02-MAR-2009 15.17	36624	+10	+20	Testing
02-MAR-2009 21.59	36628	+20	+30	Testing
03-MAR-2009 04.41	36632	+30	+40	Testing
03-MAR-2009 11.24	36636	+40	+50	Testing
03-MAR-2009 18.06	36640	+30	+50	Reduced
19 -JUN- 2009 08.08	38180	+25	+50	Testing
21 -JUN- 2009 10.26	38210	+15	+40	Testing
23 -JUN- 2009 12.44	38240	+5	+30	Testing
25 -JUN- 2009 15.02	38270	+20	+45	Testing
26 -JUN- 2009 07.48	38280	+30	+50	Reduced
17-JUL-2009 06.48	38580	+25	+50	Reduced

### 4.1.2 CURRENT OPERATIONS AND CONFIGURATION

The planned GOMOS operations for the reporting period are identified in table 4.1-2. The operation scenario of GOMOS since 29<sup>th</sup> August 2005 until end of reporting month consists of:

- Planning 2 orbits per sequence (nominal were 5): this is done because in case of a VCCS failure with subsequent loss of star observation, the maximum loss of consecutive observations cannot exceed two orbits.
- Reduced azimuth field of view (nominal was [-10°, +90°]): as the anomaly occurs during the rallying of the telescope in the preparation for the star observation, it has been decided to reduce the field of view in order to minimize the failure occurrence probability. Different ranges have been used until 2<sup>nd</sup> March 2009 (table 4.1-1) in order to optimize the number of occultations per orbit. Since 17<sup>th</sup> July 2009 the instrument is operated in an azimuth window of [25, 50] degrees due to the less occurrences of VCCS anomalies.

**Table 4.1-2: GOMOS planned operations. The planning is built on a 2-orbit sequence basis (2 orbits with the same stars)**

UTC Start	Start Orbit	Stop Orbit	Mode (Asynchronous or Synchronous)	Calibration (CAL) Dark Sky Area (DSA) or Nominal (Nom)
31-JUL-2009 06.08.07	38780	38877	S	Nom
07-AUG-2009 02.26.48	38878	38878	A	Nom
07-AUG-2009 05.47.59	38880	38977	S	Nom
14-AUG-2009 02.06.40	38978	38978	A	Nom
14-AUG-2009 05.27.52	38980	39077	S	Nom
21-AUG-2009 01.46.33	39078	39078	A	Nom
21-AUG-2009 05.07.45	39080	39177	S	Nom
28-AUG-2009 01.26.26	39178	39178	A	Nom
28-AUG-2009 04.47.38	39180	39277	S	Nom

There was no new Configurable Table Interface (CTI) uploaded to the instrument. The files used since the beginning of the mission are in table 4.1-3. The yellow ones are the current ones in use.



**Table 4.1-3: Historic CTI Tables**

CTI filename		Dissemination to FOCC
SMP	CTI_SMP_GMVIEC20030716_123904_00000000_00000004_20030715_000000_20781231_235959.N1	16-JUL-2003
	CTI_SMP_GMVIEC20021104_075734_00000000_00000003_20021002_000000_20781231_235959.N1	06-NOV-2002
	CTI_SMP_GMVIEC20021002_082339_00000000_00000002_20021002_000000_20781231_235959.N1	07-OCT-2002
	CTI_SMP_GMVIEC20020207_154455_00000000_00000000_20020301_032709_20781231_235959.N1	21-FEB-2002
STP	CTI_STP_GMTIEC20021104_080137_00000000_00000000_20021002_000000_20781231_235959.N1	04-NOV-2002
	CTI_STP_GMVIEC20021002_083222_00000000_00000000_20021002_000000_20781231_235959.N1	02-OCT-2002

## 4.2 Limb, Illumination conditions and instrument gain setting

The **limb** and the **illumination condition** are two parameters that can confuse the user community. In table 4.2-1 there are specified the product parameter (level 1b and level 2 of processor GOMOS/4.02 operational until 8<sup>th</sup> August 2006) where the flag is located, the meaning and the source. The difference between the limb (SPH/bright\_limb) and the illumination condition (SUMMARY\_QUALITY/limb\_flag) is that the first one is coming from the mission scenario and the second is coming from the processing (defined from the computation of the sun zenith and azimuth angles at both instrument and tangent point locations). The SPH/bright\_limb is for some occultations set to “dark” in the mission scenario while they are in fact in bright limb illumination conditions. To select the highest quality data for scientific applications, data with SUMMARY\_QUALITY/limb\_flag equal to ‘0’ should be used (see also the disclaimer: <http://envisat.esa.int/dataproducts/availability/disclaimers>). The instrument gain settings are also specified in table 4.2-1 (they depend on the mission scenario flags) just for completeness of information. The same is valid for the prototype version Gopr\_6.0a\_6.0a and following ones (including the one that was used for the second reprocessing of 2002-2005 years), where the **limb** is in fields SPH/bright\_limb and SUMMARY\_QUALITY/dark\_bright\_limb and the **illumination condition** is in field SUMMARY\_QUALITY/obs\_ill\_cond. For these prototypes **and the processor GOMOS/5.00 in operations since 8<sup>th</sup> August 2006**, the illumination condition can have five values (see table 4.2-2).

**Table 4.2-1: Relationship between limb, illumination condition flags and instrument gain settings (IPF version GOMOS/4.02 and previous)**

Products parameter	SPH/bright_limb	0 = Dark	1 = Bright	Coming from mission scenario
	SUMMARY_QUALITY/limb_flag	0 = Full Dark 1 = Bright 2 = Twilight	1 = Bright 2 = Twilight	In the geolocation process the sun zenith angle is computed and the occultation then is flagged accordingly
Instrument Gain	SPA Gain	3 (2)	0	Gain setting for spectrometer A. In parenthesis, values valid only for Sirius occultations (starID=1)
	SPB Gain	0	0	Gain setting for spectrometer B

**Table 4.2-2: Relationship between limb, illumination condition flags and instrument gain settings (IPF version GOMOS/5.00 and following ones; prototype version GOPR 6.0a\_6.0a and following ones)**

<b>Products parameter</b>	SPH/bright_limb SUMMARY_QUALITY/dark_bright_limb	0 = Dark	1 = Bright	Coming from mission scenario
	SUMMARY_QUALITY/obs_ill_cond	0 = Full Dark 1 = Bright 2 = Twilight 3 = Straylight 4 = Twi.+Stray		In the geolocation process the sun zenith angle is computed and the occultation is then flagged accordingly
<b>Instrument Gain</b>	SPA Gain	3 (2)	0	Gain setting for spectrometer A. In parenthesis, values valid only for Sirius occultations (starID=1)
	SPB Gain	0	0	Gain setting for spectrometer B

### 4.3 Thermal Performance

Since the beginning of the mission, the hot pixel and RTS phenomena have been producing a continuous increase of the dark charge signal within the CCD detectors (see section 4.5.1). In order to minimize this effect, three successive CCD cool downs were performed in orbits 800 (25<sup>th</sup> April 2002), 1050 (13<sup>th</sup> May 2002) and 2780 (11<sup>th</sup> September 2002) with a total decrease in temperature of 14 degrees.

Fig. 4.3-1 and 4.3-2 display, respectively, the overall temperature variation and the temperature variation around the Ascending Node Crossing (ANX) time with a resolution of 0.4 degrees (coding accuracy for level 0 data).

The CCD temperatures show the expected global increase due to the radiator ageing. Another expected variation of the temperatures, the seasonal one, can be also observed: at the beginning of mission the amplitude was around 0.8 but now it is around 1.5 degrees. The peaks that occur mainly in spectrometer B1 and B2 are also to be noted. They happen a little before the ANX for some consecutive orbits and every 8-10 days. Their origin is not known, as we did not find any correlation between these peaks and other activities carried out by other ENVISAT instruments.

The CCD temperature at almost the same latitude location (fig. 4.3-2) is monitored in order to detect any inter-orbital temperature variation. The abnormal decreases observed sometimes in all detectors are after GOMOS switch off periods, when the instrument did not have enough time to reach the nominal temperature before starting the measurements.

During the reporting period, the orbital temperature variation of the detector SPB2 for ascending and descending passes (fig. 4.3-3 and 4.3-4) is nominal (around 2.5 degrees). The stability of the temperature during the orbit is important because it affects the position of the interference patterns. The phenomenon of the interference is present mainly in SPB and this Pixel Response Non-Uniformity (PRNU) is corrected during the processing.




Figure 4.3-1: Level 0 temperature evolution of all GOMOS CCD detectors since October 2002 until the end of the reporting period




Figure 4.3-2: Level 0 temperature evolution of all GOMOS CCD detectors around ANX since November 2002 until the end of the reporting period




Figure 4.3-3: Ascending orbital variation of SPB2 temperature during reporting period




Figure 4.3-4: Descending orbital variation of SPB2 temperature during reporting period

### 4.4 Optomechanical Performance

- Version GOMOS/4.00 and previous ones: in the GOMOS processor versions GOMOS/4.00 and previous, the spectra are expected to be aligned along CCD lines, and therefore use only a single average line index per CCD. In table 4.4-1, the mean values of the location of the star signal for all the calibration analysis done is reported. The ‘left’ and ‘right’ values are calculated (the whole interval is not used) because the spectra present a slight slope, more pronounced in spectrometer B. In table 4.4-2, mean values of the location of the star signal are calculated for some specific wavelength intervals. These intervals have been changed between the calibration performed in September 2002 and the ones performed afterwards (until November 2003). Table 4.4-3 reports the average location of the star spot on the photometer 1 and 2 CCD.
- Version GOMOS/4.02: in this processor version operational since 23<sup>rd</sup> March 2004 until 8<sup>th</sup> August 2006, a Look Up Table (LUT) gives the line index of the spectra location as a function of the wavelength. The values obtained during calibration exercises are shown in table 4.4-4. These values should be similar to the ones of the LUT; otherwise the LUT should be updated. However this characterization curve is not exactly the location of the star spectrum on the CCD but rather a combination of this position and some artefact created by the shape of the instrument optical point spread function (PSF). The exact shape is actually a straight line (especially for SPB) that has been characterised in 2005.
- Current version GOMOS/5.00 (since 8<sup>th</sup> August 2006): the exact shape of the CCD spectra location curve (which is a straight line) that has been characterised in 2005 was implemented in the current set of GOMOS ADFs. The position of the spectra convoluted with the PSF is calculated during the processing.

**Table 4.4-1: Mean value of the location of the star signal during the occultation at the edges of every band (mean over 50 values, filtering the outliers)**

	UV (SPA1) left/right	VIS (SPA2) left/right (Inverted spectra)	IR1 (SPB1) left/right	IR2 (SPB2) left/right
11/09/2002	80.7/80.7	79.8/79.5	82.8/81.9	83.1/82.1
01/01/2003	80.7/80.6	79.8/79.5	82.8/82.0	83.2/82.2
17/07/2003 & 02/08/2003	80.7/80.7	79.8/79.5	82.8/81.9	83.1/82.1
08/11/2003	80.7/80.6	79.8/79.5	82.8/81.9	83.1/82.1

**Table 4.4-2: Mean value of the location of the star signal during the occultation (as table 4.4-1) but now within some wavelength intervals**

	UV (SPA1)	VIS (SPA2)	IR1 (SPB1)	IR2 (SPB2)
11/09/2002 wl range (nm)	80.8 [300-330]	79.8 [500-530]	82.6 [760-765]	82.9 [937-942]
01/01/2003 wl range (nm)	80.6 [350-360]	78.6 [650-670]	81.6 [760-765]	80.3 [935-945]
02/08/2003	80.6	79.7	82.5	82.8
08/11/2003	80.6	79.9	82.4	82.8

**Table 4.4-3: Average column and row pixel location of the star spot on the photometer CCD during the occultation**

	FP1 (column/row)	FP2 (column/row)
11/09/2002	11/4	5/5
01/01/2003	10/4	6/4.9
02/08/2003	10/4	6/5
08/11/2003	10/4	6/5

Table 4.4-4: Location of the star signal on the CCD's

Pixel Column	LUT (Pixel line)	Calibration on 10-APR-2004	Calibration on 04-DEC-2004	Calibration on 27-NOV-2005	Calibration on 19-FEB-2006	Calibration on 14-MAY-2006 and 11-JUN-2006
0	80.59	80.80	80.67	80.93	80.67	80.85
20	80.46	80.60	80.44	80.32	80.43	80.49
449	80.42	80.50	80.42	80.40	80.53	80.56
450	79.25	79.39	79.30	79.16	79.30	79.35
900	79.50	79.63	79.57	79.36	79.45	79.61
1415	79.70	79.76	79.76	80.00	79.81	79.93
1416	82.64	82.80	82.88	82.95	82.76	82.81
1500	82.31	82.60	82.66	82.63	82.58	82.55
1600	82.12	82.22	82.30	82.35	82.41	82.20
1700	81.97	82.04	82.08	82.09	82.05	82.06
1750	81.89	81.98	82.03	82.00	81.92	81.97
1800	81.78	81.91	81.96	81.93	81.83	81.98
1835	81.68	81.88	81.94	81.96	81.79	81.91
1836	82.98	83.10	83.10	83.27	83.17	83.08
2000	82.78	82.90	82.94	83.04	82.83	82.93
2100	82.33	82.70	82.73	82.82	82.83	82.67
2150	82.17	82.40	82.54	82.79	82.70	82.49
2350	81.83	82.00	82.00	82.68	81.96	82.11

## 4.5 *Electronic Performance*

### 4.5.1 DARK CHARGE EVOLUTION AND TREND

The trend of Dark Charge (DC) is of crucial importance for the final quality of the products, and is therefore subject to intense monitoring. As part of the DC there is:

- “Hot pixels”, a pixel is “hot” when its dark charge exceeds its value measured on ground, at the same temperature, by a significant amount.
- RTS phenomenon (Random Telegraphic Signal), it is an abrupt change (positive or negative) of the CCD pixel signal, random in time, affecting only the DC part of the signal and not the photon generated signal.

The temperature dependence of the DC would make this parameter a good indicator of the DC behaviour, but the hot pixels and the RTS are producing a continuous increase of the DC (see trend in fig. 4.5-1 and 4.5-2). To take into account these phenomena, since version GOMOS/4.00 (the current one is GOMOS/5.00) a DC map per orbit is extracted from a Dark Sky Area (DSA) observation performed around ANX (full dark conditions). For every level 1b product (occultation), the actual thermistor temperature of the CCD is used to convert the DC map measured around ANX into an estimate of the DC at the time (and different temperature) of the actual occultation. When the DSA observation is not available, the DC map inside the calibration product that was measured at a given thermistor reference temperature is used; again, the actual thermistor temperature of the CCD is used to compute the actual map. Table 4.5-1 reports the list of products that used the DC maps inside the calibration file due to the non-availability of DSA observation. A “CAL DC map with no T dep.” means that, as the temperature information was not available for that occultation, the DC map used is exactly the one inside the Calibration product.

The “quality ranking” of the products depending on DC correction performed is as follows:

- Best quality: products with DC correction using DSA observation inside the orbit
- Less quality than previous ones: products with DC correction using the map inside the calibration product, thermal corrected ('DC map used' in table 4.5-1)
- Less quality than previous ones: products with DC correction using the map inside the calibration product, no thermal corrected ('DC map with no T dep.' in table 4.5-1)

**Table 4.5-1: Table of level 1b products that used the Calibration DC maps instead of the DSA observation. (Complete table in APPENDIX A)**

Product name	DC information
GOM_TRA_1PNPDE20090803_195313_000000592081_00200_38831_6836.N1	DC map used
GOM_TRA_1PNPDE20090803_195800_000000622081_00200_38831_6837.N1	DC map used
GOM_TRA_1PNPDE20090803_200453_000000502081_00200_38831_6838.N1	DC map used
GOM_TRA_1PNPDE20090803_200855_000000492081_00200_38831_6839.N1	DC map used
GOM_TRA_1PNPDE20090803_202119_000000412081_00200_38831_6840.N1	DC map used
GOM_TRA_1PNPDE20090803_202428_000000382081_00200_38831_6841.N1	DC map used
GOM_TRA_1PNPDE20090803_202954_000000402081_00200_38831_6842.N1	DC map used
GOM_TRA_1PNPDE20090803_203606_000000522081_00200_38831_6843.N1	DC map used
GOM_TRA_1PNPDE20090803_203808_000000512081_00200_38831_6844.N1	DC map used
...	...
...	...
...	...
GOM_TRA_1PNPDE20090803_204443_000000492081_00200_38831_6845.N1	DC map used
GOM_TRA_1PNPDE20090803_204626_000000712081_00200_38831_6846.N1	DC map used
GOM_TRA_1PNPDE20090803_204840_000000442081_00200_38831_6847.N1	DC map used
GOM_TRA_1PNPDE20090828_204804_000000472082_00057_39189_1424.N1	DC map used
GOM_TRA_1PNPDE20090828_210227_000000592082_00057_39189_1425.N1	DC map used
GOM_TRA_1PNPDE20090828_210558_000000472082_00057_39189_1426.N1	DC map used
GOM_TRA_1PNPDE20090831_211143_000000492082_00100_39232_5505.N1	DC map used
GOM_TRA_1PNPDE20090831_211651_000000542082_00100_39232_5506.N1	DC map used
GOM_TRA_1PNPDE20090831_211837_000000492082_00100_39232_5507.N1	DC map used
GOM_TRA_1PNPDE20090831_212312_000000722082_00100_39232_5508.N1	DC map used

The average DC inserted by the processor into the level 1b data products for the spectrometers SPA1 and SPB2 (per band: upper, central and lower) is plotted in fig. 4.5-1 and 4.5-2. The abnormal decreases observed sometimes in all detectors are due to the temperature decreases that occur after GOMOS switch off periods. The same DC values are plotted in fig. 4.5-3 but for the ESRIN occultations belonging only to the reporting month.






Figure 4.5-1: Mean DC evolution on SPA1 since 15<sup>th</sup> December 2002 until the end of the reporting period




Figure 4.5-2: Mean DC evolution on SPB2 from 15<sup>th</sup> December 2002 until the end of the reporting period




Figure 4.5-3: Mean Dark Charge of spectrometers during the reporting period

### 4.5.2 SIGNAL MODULATION

A parasitic signal was found to be systematically present, added to the useful signal, for the spectrometers A and B (fig. 4.5-4). The modulation is corrected in the data processing for spectrometers A1 and A2, for spectrometer B it has much smaller amplitude and so it is not corrected.

The values of the modulation (fig. 4.5-4) are daily extracted and plotted; they should not be very different from the ones coded into the processor: 1.40 ADU for SPA1 and 0.76 ADU for SPA2.

Fig. 4.5-4 shows high values during summer time for the ESRIN data, it now being confirmed that the South Atlantic Anomaly is the cause of these unexpected peaks. The quality of ESRIN data, in particular over the SAA zone, is impacted but the measure of this impact is under investigation. However, in the second half of the months of October for all years (2004-2008) the peaks are smaller because the DSA zone where the data are taken for this analysis is moving towards the Northern Hemisphere. At the end of October the DSA zone is definitely chosen by the planning system in the Northern Hemisphere (to fill the criteria ‘DSA in full dark limb conditions’) and the high peaks disappear.




Figure 4.5-4: Modulation signal

### 4.5.3 ELECTRONIC CHAIN GAIN AND OFFSET

No new electronic chain gain and offset calibration has been done during the reporting period. The routine monitoring of the ADC offset is a good indicator of the ageing of the instrument electronics. The fig. 4.5-5 presents the evolution of the calibrated ADC offset for each spectrometer electronic chain. The unexpected increase of this offset seems to be due to an external contribution. In the ADC offset calibration procedure, linearity observations are used with two integration times of 0.25 and 0.50 seconds to extrapolate to an integration time of 0 seconds that gives the complete chain offset and not only the ADC offset. The complete offset contains any possible offsets, and especially the static dark charge (i.e. the dark charge that does not depend on the spectrometer integration time). The presence of vertical lines visible in the measurement maps in spatial spread monitoring mode confirms that the memory area of the CCD is affected by the generation of hot pixels. These new hot pixels are one contributor to the increase observed in fig. 4.5-5.




Figure 4.5-5: ADC offset evolution for each spectrometer electronic chain

A current QWG task consists in completing the analysis to confirm that the offset increase is also due to the expected dark charge increase in the memory area due to ageing. This can be proven by the study of the noise due to the increased dark charge. The increase of ADC offset will be assumed to be equal to the increase of ‘static dark charge’ and the corresponding noise will be computed and compared to the increase of the residual of the signal variance.

If we keep the ADC offset constant, as it is also used to compute the dark charge at band level (which is used to correct the samples in the level 1b processing), the increase of the static dark charge - not taken into account in the ADC offset - is compensated by an artificial increase of the calibrated dark charge. So, the star and limb spectra are correctly corrected for dark charge. A small bias can be added to the instrument noise due to the incorrect dark charge level. Anyway, this quantity is not large enough to require a modification of the ADC offset value.

## 4.6 Acquisition, Detection and Pointing Performance

### 4.6.1 SATU NOISE EQUIVALENT ANGLE

The Star Acquisition and Tracking Unit (SATU) noise equivalent angle (SATU NEA) consists of the statistical angular variation of the SATU data above the atmosphere. The mean of the standard deviation (STD over the 50 values per measurement) above 105 km are computed for every occultation, giving two values per occultation: one in the ‘X’ direction, one in the ‘Y’ direction. A mean value per day in every direction and limb is calculated and monitored in order to assess instrument performance in terms of star pointing (fig. 4.6-1). Also monthly averages are calculated and plotted (fig. 4.6-2). The thresholds are 2 and 3 micro radians in ‘X’ and ‘Y’ directions respectively. Before May 2003, data above 90 km have been considered (instead of 105 km) but from May 2003 on, data taken in the mesospheric oxygen layer (located around 100 km altitude) have been avoided because they could cause fluctuations on the SATU data. Also the products with errors (error flag set) are discarded from May 2003 onwards.




Figure 4.6-1: Average value per day of SATU NEA STD above 105 km

Different anomalies have affected the SATU during the mission:

- **Sudden increase on September 2005:** as can be seen in fig. 4-6.1, the SATU NEA had a sudden increase on 8<sup>th</sup> September 2005 mainly in ‘Y’ axis. These values remained high, fluctuating between 1 and 1.8 microrad until December 2005 when they came back to the values they used to be before the increase of September. The reason why there was higher noise in the data causing the jump in daily SATU average is not known.
- **Gradual increase on mid April 2006:** a different problem was present since mid April 2006 until October 2007. A gradual increase of the daily SATU ‘Y’ mean was observed. This increase was due to fluctuations of the SATU ‘Y’ data observed at the beginning of nominal occultations (starting at 130 km that corresponds to an elevation angle of around 65°). The decrease of the start elevation angle of the occultation has no impact on the amplitude of the SATU ‘Y’ fluctuations. Investigations carried out by the ESL, ESA and industry pointed to a problem on the SFM (mechanical or electrical) and not to a problem on the SATU itself. Since October 2007 the fluctuations have disappeared and as a consequence the daily SATU ‘Y’ average has come back below the threshold set to 3 micro radians.
- **Current anomaly: sudden increase on December 2008:** similarly to the anomaly happened on April 2006, the SATU NEA had an increase on 29<sup>th</sup> December 2008 due to fluctuations of the SATU ‘Y’ data. The difference with respect to the previous anomaly is that this time, the increase was quite sudden and the fluctuations are present during the whole occultation, not only at the beginning of the occultation. The most critical effect of this anomaly is the loss of the star measurement high in the atmosphere, which means that many times the corresponding ozone profiles do not include the ozone peak present at around 25-30 km. After the increase of the

elevation filter from 100 to 150 on 29<sup>th</sup> June 2009, the abortion of the star measurements is now deeper in the atmosphere but still premature. A new increase on 6<sup>th</sup> July (from 150 to 200) did not produce an immediate improvement but it seems that after that date a gradual improve took place.

The results for some occultations belonging to previous months (monthly averages) are presented in fig. 4.6-2, where the change in trend in September 2005, May 2006 and December 2008, mainly for the ‘Y’ axis is visible.




Figure 4.6-2: Average value per month of SATU NEA STD above 105 km


### 4.6.2 TRACKING LOSS INFORMATION

This verification consists of the monitoring of the tangent altitude at which the star is lost. It is an indicator of the pointing performance although it is to be considered that star tracking is also lost due to the presence of clouds and hence not only due to deficiencies in the pointing performance. Therefore, only the detection of any systematic long-term trend is the main purpose of this monitoring. The recent results are presented in fig. 4.6-3 and 4.6-4:


- The dependence of the altitude at which tracking is lost on the magnitude of the star is very small because the tracking is mainly lost due to the refraction and the scintillation that depend on the atmospheric conditions.
- The altitude at which the stars are lost is now almost nominal after the two gain increases performed (first one on 29<sup>th</sup> June and second one on 6<sup>th</sup> July 2009).
- In bright limb it is not expected that the stars are lost at very low altitudes due to the amount of light arriving to the pointing system mainly when the refraction effects start to be important. We see from fig. 4.6-4 that there are some stars lost at altitudes around 4 km. This occurs when the pointing system is not able to point to the star anymore but, instead of finishing the occultation, it continues to track light until the planned duration is reached.
- Daily statistics are given in fig. 4.6-5 (calculated using 50 products per day). The high peaks in standard deviation before 25<sup>th</sup> January 2005 are due to the long lasting occultations or partial

occultations (the entire occultation is included within the following orbit data). The ones during June/July/August 2005 are due to the tests performed for anomaly investigation. On 2009 (until July) the peaks are due to the elevation anomaly.

- Monthly statistics are given in fig. 4.6-6 (calculated using 50 products per day) where the change in trend, mainly for dark limb, is visible for the period of GOMOS elevation anomaly on 2009.



**Figure 4.6-3: Last tangent altitude of the occultation (dark limb), point at which the star is lost**



**Figure 4.6-4: Last tangent altitude of the occultation (bright limb), point at which the star is lost**




Figure 4.6-5: Daily average and STD of tangent altitude loss for the reporting period




Figure 4.6-6: Monthly mean tangent altitude (and STD) at which the star is lost since January 2003




### 4.6.3 MOST ILLUMINATED PIXEL (MIP)

The MIP (Most Illuminated Pixel) is the star position on the SATU CCD in detection mode and it is recorded in the housekeeping data. The nominal centre of the SATU is pixel number **145** in elevation and number **205** in azimuth. The detection of the stars should not be far from this centre. As it can be seen in fig. 4.6-7 the **azimuth MIP** was within the threshold (table 4.6-1) since September 2002 until the occurrence of the anomaly on January 2005, even if a small variation is present. The reason for the change in trend observed after the anomaly is, at the moment, not understood. The **elevation MIP** had a significant variation (see the *note* below) until 12<sup>th</sup> December 2003 when a new PSO algorithm was activated in order to reduce the deviations of the ENVISAT platform attitude with respect to the nominal one. Similarly to the azimuth, after the anomaly of January 2005 the Elevation MIP has a drift that has no explanation. Investigations are ongoing to try to understand this behavior of the MIP as, although it does not impact the data quality or the star location on the CCD array during the measurements, it may invalidate attitude monitoring by GOMOS and could represent a hidden anomaly.

*Note:* A MIP variation onto the SATU CCD of 50 pixels corresponds to a de-pointing of 0.1 degrees

**Table 4.6-1: MIP Thresholds**


<b>MIP X</b>	<b>Mean delta Az</b>	[198 - 210]
	<b>Std delta Az</b>	7
<b>MIP Y</b>	<b>Mean delta El</b>	[140 – 150]
	<b>Std delta El</b>	4



**Figure 4.6-7: Mean values of MIP for some orbits since 1<sup>st</sup> September 2002 (see table 4.6-1)**

Fig. 4.6-8 shows the standard deviation of azimuth and elevation MIP that should be within the thresholds of table 4.6-1. The peaks observed mean that one (or more) stars were detected very far from

the SATU detection point and, in this case, the stars were lost during the centering phase (see section 3.2 for stars lost in centering).



**Figure 4.6-8: Standard deviation of MIP Azimuth and Elevation for some orbits since 1<sup>st</sup> September 2002 until end of reporting period (see table 4.6-1)**

## 5 LEVEL 1 PRODUCT QUALITY MONITORING

### 5.1 Processor Configuration

#### 5.1.1 VERSION

Around 40% of near real time GOM\_TRA\_1P products have been received by the IDEAS team for routine quality control and long term trend quality monitoring. The current level 1-processor software version for the operational ground segment is GOMOS/5.00 since 8<sup>th</sup> August 2006 (see table 5.1-1). The product specification is PO-RS-MDA-GS2009\_10\_3I. This processor has been cleared for level 1 data release, with a disclaimer for known artefacts (<http://envisat.esa.int/dataproducts/availability/disclaimers>) that are currently being resolved and will be implemented in following releases of the processor (<http://envisat.esa.int/dataproducts/availability>).

Users are also supplied with 2002 - 4<sup>th</sup> July 2006 data sets reprocessed by the last prototype processor GOPR\_6.0c\_6.0f developed and operated by ACRI. See table 5.1-2 for prototype level 1b versions and modifications. The current GOMOS operational ground segment version GOMOS/5.00 is in line with the prototype version used for this second reprocessing.

**Table 5.1-1: PDS level 1b product version and main modifications implemented**

Date	Version	Description of changes
08-AUG-2006	Level 1b version 5.00 at PDHS-E, PDHS-K	Algorithm baseline level 1b DPM 6.3 <ul style="list-style-type: none"> <li>• Correction of FP unfolding algorithm</li> <li>• Background correction of SPB in full dark limb</li> <li>• Modification of the computation of the incidence angle</li> <li>• Correction of the flat-field correction equations</li> <li>• Star spectrum location on CCD modified for SPB</li> <li>• Provide SFA and SATU angles in degrees</li> <li>• Elevation angle dependency of the reflectivity LUT added in the algorithms</li> <li>• Ratio upper/star signal added (FLAGUC)</li> <li>• Add Dark Charge used for dark charge correction (per band)</li> <li>• Flag for illumination condition (PCDillum)</li> <li>• Minimum sample value for which the cosmic rays detection processing is applied (Crmin) is a function of gain index</li> <li>• Logic for computation of the flags attached to the reference star spectrum (Flref) modified</li> <li>• Add the computation of the sun direction in the inertial geocentric frame to be written in the level 1b and limb products.</li> <li>• Spectrometer effective sampling time added</li> </ul>
23-JUL-2006	Level 1b version 5.00 at LRAC	Change in configuration at the time of switch over: <ul style="list-style-type: none"> <li>○ Use of new reflectivity LUT (GOM_CAL_AX)</li> <li>○ New wavelength assignment for SPA1, A2, B1 (GOM_CAL_AX)</li> <li>○ Location of star spectrum projection on the CCD arrays (GOM_CAL_AX)</li> <li>○ Spatial PSF of SPB modified (GOM_INS_AX)</li> <li>○ Some universal constants (GOM_PR1_AX)</li> </ul>
23-MAR-2004	Level 1b version 4.02 at PDHS-E and PDHS-K	Algorithm baseline level 1b DPM 6.0 <ul style="list-style-type: none"> <li>• Adding a new calibration parameters (these values are hard coded at the moment)</li> <li>• Removal of redundancy chain from code</li> <li>• Modifications in the processing to apply new configuration and calibration parameter</li> <li>• New algorithm to determine between dark, twilight and bright limb and to handle data accordingly</li> <li>• Added handling of source packages with invalid packet header</li> <li>• Added enumerations for all configuration flags</li> </ul>
31-MAY-2003	Level 1b version 4.00 at PDHS-E and PDHS-K	Algorithm baseline level 1b DPM 5.4: <ul style="list-style-type: none"> <li>• Modulation correction step added after the cosmic rays detection processing</li> <li>• Inversion of the non-linearity and offset corrections</li> <li>• Modification of the computation of the estimated background signal measured by the photometers: use the spectrometer radiometric sensitivity curve and the photometer transfer function.</li> <li>• Use of the dark charge map at orbit level computed from the DSA (dark sky area) if any in the level 0 product</li> <li>• Implementation of a new unfolding algorithm for the photometer samples</li> </ul>
21-NOV-2002	Level 1b version 3.61 at PDHS-E and PDHS-K	Algorithm baseline DPM 5.3: <ul style="list-style-type: none"> <li>• Review of some default values</li> <li>• New definition of one PCD flag (atmosphere)</li> <li>• Temporal interpolation of ECMWF data</li> </ul>

**Table 5.1-2: GOPR level 1b product version and main modifications implemented**

Date	Version	Description of changes
22-JUL-2005	GOPR_6.0c	Level 1b: <ul style="list-style-type: none"> <li>• Correction of FP unfolding algorithm</li> <li>• Background correction of SPB in full dark limb</li> <li>• Modification of the computation of the incidence angle</li> <li>• Correction of the flat-field correction equations</li> <li>• Star spectrum location on CCD modified for SPB</li> </ul> Configuration for second reprocessing: <ul style="list-style-type: none"> <li>○ Use of new reflectivity LUT</li> <li>○ New wavelength assignment for SPA1, A2, B1</li> <li>○ Spatial PSF of SPB modified</li> </ul>
17-MAR-2004	GOPR 6.0a	<ul style="list-style-type: none"> <li>• Provide SFA and SATU angles in degrees</li> <li>• Elevation angle dependency of the reflectivity LUT added in the algorithms</li> <li>• Ratio upper/star signal added (FLAGUC)</li> <li>• Add Dark Charge used for dark charge correction (per band)</li> <li>• Flag for illumination condition (PCDillum)</li> <li>• Minimum sample value for which the cosmic rays detection processing is applied (Crmin) is a function of gain index</li> <li>• Logic for computation of the flags attached to the reference star spectrum (Flref) modified</li> <li>• Add the computation of the sun direction in the inertial geocentric frame to be written in the level 1b and limb products.</li> <li>• Spectrometer effective sampling time added</li> </ul>
25-JUL-2003	GOPR 5.4f	<ul style="list-style-type: none"> <li>• The demodulation process is applied only in full dark limb and twilight limb conditions.</li> </ul>
17-JUL-2003	GOPR 5.4e	<ul style="list-style-type: none"> <li>• Sun zenith angle is computed in the geolocation process. The occultation is now classified into (0) full dark limb condition, (1) bright limb condition and (2) twilight limb condition.</li> <li>• No background correction applied in full dark limb condition. The location of the image of the star spectrum on the CCD array is no more aligned with the CCD lines.</li> </ul>
02-JUL2003	GOPR 5.4d	<ul style="list-style-type: none"> <li>• The maximum number of measurements is set to 509 (instead of 510) in the GOPR prototype.</li> </ul>
17-MAR-2003	GOPR 5.4c	<ul style="list-style-type: none"> <li>• Modification of the CAL ADFs (update of the limb radiometric LUT). The products are affected only if the limb spectra are converted into physical units</li> <li>• Modifications to allow compatibility with ACRI computational cluster (no modifications of the results)</li> <li>• Modification of the logic to handle dark charge map refresh at orbit level (DSA data is now directly processed by the level 1b processor if available in the level 0 product). No impact on the results</li> </ul>
21-FEB-2003	GOPR 5.4b	<ul style="list-style-type: none"> <li>• DC map values are rounded when written in the level 1b product</li> <li>• Modification of the CAL ADFs (update of the wavelength assignment of SPB1 and SPB2)</li> <li>• Modify the computation of flag_mod in the modulation correction routine</li> </ul>
17-JAN-2003	GOPR 5.4a	<ul style="list-style-type: none"> <li>• use the start and stop dates of the occultation when calling the CFI Interpol instead of start and stop dates of the level 0 product</li> <li>• modify the ECMWF filename information in the SPH of the level 1b and limb products</li> </ul>

### 5.1.2 AUXILIARY DATA FILES (ADF)

The ADF's files in tables 5.1-3, 5.1-4, 5.1-5, 5.1-6 and 5.1-7 have been disseminated to the PDS during the whole mission. Note that the files outlined in yellow are the set of auxiliary files used during the reporting period. For every type of file, the validity runs from the start validity time until the start validity time of the following one, but if an ADF file has been disseminated after the start validity time, it is obvious that it will be used by the PDHS-E and PDHS-K PDS only after the dissemination time (this happens the majority of the time). Just like the other ADF's, the calibration auxiliary file (GOM\_CAL\_AX) has been updated several times in the past (table 5.1-7) but the difference is that now it is updated in a weekly basis with new DC maps and new wavelength assignment (routine weekly wavelength calibration was activated on 14<sup>th</sup> December 2007), and that is why the files used during reporting period are reported in a separate table (table 5.1-8) that changes from report to report.

**Table 5.1-3: Historic GOM\_PR1\_AX files used by PDS for level 1b products generation. The GOM\_PR1\_AX is a file containing the configuration parameters used for processing from level 0 to level 1b products**

Used by PDS for Level 1b products generation during	GOM_PR1_AX (GOMOS processing level 1b configuration file)
01-MAR-2002 → 29-MAR-2002	<b>GOM_PR1_AXVIEC20020121_165314_20020101_000000_20200101_000000</b> <ul style="list-style-type: none"> <li>• Pre-launch configuration</li> </ul>
30-MAR-2002 → 14-NOV-2002	<b>GOM_PR1_AXVIEC20020329_115921_20020324_200000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Changed num_grid_upper, thr_conv and max_iter in the atmospheric GADS</li> </ul>
Not used	<b>GOM_PR1_AXVIEC20020729_083756_20020301_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Cosmic Ray mode + threshold</li> <li>• DC correction based on maps</li> <li>• Non-linearity correction disabled</li> </ul>
Not used	<b>GOM_PR1_AXVIEC20021112_170331_20020301_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Central background estimation by linear interpolation + associated thresholds</li> </ul>
15-NOV-2002 → 26-MAR-2003	<b>GOM_PR1_AXVIEC20021114_153119_20020324_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Same content as GOM_PR1_AXVIEC20021112_170331_20020301_000000_20100101_000000 but validity start updated so as to supersede according to the PDS file selection rules</li> <li>• GOM_PR1_AXVIEC20020329_115921_20020324_200000_20100101_000000</li> </ul>
27-MAR-2003 → 19-MAR-2004	<b>GOM_PR1_AXVIEC20030326_085805_20020324_200000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Same content as GOM_PR1_AXVIEC20021112_170331_20020301_000000_20100101_000000 but validity start updated so as to supersede according to the PDS file selection rules</li> <li>• GOM_PR1_AXVIEC20020329_115921_20020324_200000_20100101_000000</li> </ul>
20-MAR-2004 → 22-MAR-2004	<b>GOM_PR1_AXVIEC20040319_134932_20020324_200000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Ray tracing parameter changed: convergence criteria set to 0.1 microrad</li> </ul>
23-MAR-2004 → 01-APR-2004 <i>Notes:</i> <ul style="list-style-type: none"> <li>• This file was constructed from GOM_PR1_AXVIEC20030326_085805_20020324_200000_20100101_000000</li> </ul>	<b>GOM_PR1_AXVIEC20040316_144850_20020324_200000_20100101_000000</b> GOM_PR1 ADF for version GOMOS/4.02, changes: <ul style="list-style-type: none"> <li>• The central band estimation mode</li> <li>• Atmosphere thickness</li> <li>• Altitude discretisation</li> </ul>

(so without the ray tracing parameter changed)	
<ul style="list-style-type: none"> <li>This file was used by the GOMOS/4.02 processors before the IECF dissemination. The dissemination was done on 25<sup>th</sup> March 2004</li> </ul>	
02-APR-2004 → 07-AUG-2006	<b>GOM_PR1_AXVIEC20040401_083133_20020324_200000_20100101_000000</b> <ul style="list-style-type: none"> <li>Ray tracing parameter changed: convergence criteria set to 0.1 microrad</li> </ul>
08-AUG-2006 Used at the time of switching over GOMOS/5.00	<b>GOM_PR1_AKNIEC20050627_151042_20020301_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>Change of some universal constants</li> </ul>

**Table 5.1-4: Historic GOM\_INS\_AX files used by PDS for level 1b products generation. The GOM\_INS\_AX is a file containing the characteristics of the instrument and it is used for processing from level 0 to level 1b products and from level 1b to level 2 products**

Used by PDS for Level 1b products generation during	GOM_INS_AX (GOMOS instrument characteristics file)
01-MAR-2002 → 29-JUL-2002	<b>GOM_INS_AXVIEC20020121_165107_20020101_000000_20200101_000000</b> <ul style="list-style-type: none"> <li>Pre-launch configuration</li> </ul>
30-JUL-2002 → 12-NOV-2002	<b>GOM_INS_AXVIEC20020729_083625_20020301_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>Factors for the conversion of the SFA angles from SFM axes to GOMOS axes</li> </ul>
13-NOV-2002 → 16-JUL-2003	<b>GOM_INS_AXVIEC20021112_170146_20020301_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>No more invalid spectral range</li> </ul>
Not used	<b>GOM_INS_AXVIEC20030716_080112_20030711_120000_20100101_000000</b> <ul style="list-style-type: none"> <li>New value for SFM elevation zero offset for redundant chain: 10004</li> </ul>
17-JUL-2003 → 07-AUG-2006	<b>GOM_INS_AXVIEC20030716_105425_20030716_120000_20100101_000000</b> <ul style="list-style-type: none"> <li>Bias induct azimuth redundant value set to -0.0084 rad (-0.4813 deg)</li> </ul>
08-AUG-2006 Used at the time of switching over GOMOS/5.00	<b>GOM_INS_AKNIEC20050627_150713_20030716_120000_20100101_000000</b> <ul style="list-style-type: none"> <li>The spatial PSF of SPB</li> </ul>

**Table 5.1-5: Historic GOM\_CAT\_AX files used by PDS for level 1b products generation. The GOM\_CAT\_AX is a file holding the star catalogue used for processing from level 0 to level 1b products**

Used by PDS for Level 1b products generation during	GOM_CAT_AX (GOMOS Stat Catalogue file)
01-MAR-2002	<b>GOM_CAT_AXVIEC20020121_161009_20020101_000000_20200101_000000</b> <ul style="list-style-type: none"> <li>Pre-launch configuration</li> </ul>

**Table 5.1-6: Historic GOM\_STS\_AX files used by PDS for level 1b products generation. The GOM\_STS\_AX is a file containing star spectra used for processing from level 0 to level 1b products**

Used by PDS for Level 1b products generation during	GOM_STS_AX (GOMOS Star Spectra file)
01-MAR-2002 → 07-AUG-2006	<b>GOM_STS_AXVIEC20020121_165822_20020101_000000_20200101_000000</b> <ul style="list-style-type: none"> <li>Pre-launch configuration</li> </ul>
08-AUG-2006 Used at the time of switching over GOMOS/5.00	<b>GOM_STS_AKNIEC20040308_103538_20020101_160000_20100101_000000</b> <ul style="list-style-type: none"> <li>Wavelength assignment GADS has been suppressed from the product</li> <li>Wavelength assignment vector has been added to the star spectrum</li> </ul>

**Table 5.1-7: Historic GOM\_CAL\_AX files used by PDS for level 1b products generation. The GOM\_CAL\_AX is a file containing the calibration parameters used for processing from level 0 to level 1b products**

Used by PDS for Level 1b products generation during	GOM_CAL_AX (GOMOS Calibration file)
01-MAR-2002 → 29-JUL-2002	<b>GOM_CAL_AXVIEC20020121_164808_20020101_000000_20200101_000000</b> <ul style="list-style-type: none"> <li>• Pre-launch configuration</li> </ul>
Not used	<b>GOM_CAL_AXVIEC20020121_142519_20020101_000000_20200101_000000</b> <ul style="list-style-type: none"> <li>• Pre-launch configuration</li> </ul>
30-JUL-2002 → 12-NOV-2002	<b>GOM_CAL_AXVIEC20020729_082426_20020717_193500_20100101_000000</b> <ul style="list-style-type: none"> <li>• Band setting information</li> <li>• Wavelength assignment</li> <li>• Spectral dispersion LUT</li> <li>• ADC offset for Spectrometers</li> <li>• PRNU maps</li> <li>• Thermistor coding LUT</li> <li>• DC maps</li> </ul>
Not used	<b>GOM_CAL_AXVIEC20021112_165603_20020914_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Band setting information</li> <li>• DC maps</li> <li>• PRNU maps</li> <li>• Wavelength assignment</li> <li>• Spectral dispersion LUT</li> <li>• Radiometric sensitivity LUT (star and limb)</li> <li>• SP-FP intercalibration LUT</li> <li>• Vignetting LUT</li> <li>• Reflectivity LUT</li> <li>• ADC offset</li> </ul>
13-NOV-2002 → 30-JAN-2003	<b>GOM_CAL_AXVIEC20021112_165948_20021019_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Only DC maps updated</li> </ul>
31-JAN-2003 → 11-APR-2003	<b>GOM_CAL_AXVIEC20030130_133032_20030101_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Only DC maps updated (using DSA of orbit 04541)</li> </ul>
12-APR-2003 → 02-JUN-2003	<b>GOM_CAL_AXVIEC20030411_065739_20030407_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Modification of the radiometric sensitivity curve for the limb spectra. Note that the modification of this LUT has no impact on the GOMOS processing. The LUT is just copied into the level 1b limb product for user conversion purpose.</li> <li>• Updated DC map only (using DSA of orbit 05762).</li> </ul>
03-JUN-2003: from this date onwards, mainly updates to DC maps are done. Every month, the table of new GOM_CAL files with <b>only</b> DC maps updated is provided (table 5.1-8). Eventual changes to this file not corresponding only to DC maps updates will be reported in this table.	<b>GOM_CAL_AXVIEC20030602_094748_20030531_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Updated DC maps only (using DSA of orbit 06530)</li> </ul>
13-FEB-2004 → 23-FEB-2004	<b>GOM_CAL_AXVIEC20040212_103916_20040209_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>• Update of the reflectivity LUT</li> <li>• Updated DC maps (Orbit 10194, date 11-FEB-2004)</li> </ul>
08-AUG-2006 Used at the time of switching over GOMOS/5.00	<b>GOM_CAL_AXNIEC20050704_110915_20050125_224800_20100101_000000</b> <ul style="list-style-type: none"> <li>• Reflectivity LUT updated</li> <li>• Location of the star spectrum projection on the CCD arrays</li> <li>• Wavelength assignment of the spectra updated</li> <li>• The spatial LSF of SPB updated</li> <li>• Updated DC maps (orbit 15200, date 25 JAN 2005)</li> </ul>

**Table 5.1-8: Calibration ADF for reporting period. These files are updated (only with new DC maps and wavelength calibrated) in a 8-10 days basis**

Used by PDS for Level 1b products generation during	GOM_CAL_AX (GOMOS Calibration file)
31-JUL-2009 → 06-AUG-2009	<b>GOM_CAL_AXVIEC20090730_093840_20090728_000000_20100101_000000</b> (orbit 38755, date 29 JUL 2009)
07-AUG-2009 → 11-AUG-2009	<b>GOM_CAL_AXVIEC20090806_083422_20090804_000000_20100101_000000</b> (orbit 38859, date 05 AUG 2009)
12-AUG-2009 → 18-AUG-2009	<b>GOM_CAL_AXVIEC20090811_154222_20090809_000000_20100101_000000</b> (orbit 38928, date 10-AUG-2009)
19-AUG-2009 → 31-AUG-2009	<b>GOM_CAL_AXVIEC20090818_144743_20090816_000000_20100101_000000</b> (orbit 39024, date 16-AUG-2009)

## 5.2 Quality Flags Monitoring

In this section, the results of monitoring some Product Quality information stored in level 1b products that did not have a fatal error (MPH error flag not set) are discussed. The products with fatal errors were around 1% of the products received during the reporting month for the quality monitoring.

On the one hand, for every product we have information of the **number of measurements** where a given problem was detected (i.e. number of invalid measurements, number of measurements containing saturated samples, number of measurements with demodulation flag set...). On the other hand, there are **flags** that indicate problems within the product (i.e. flag set to one if the reference spectrum was computed from DB, flag set to zero if SATU data were not used...).



For the information on the number of measurements a plot of percentages with respect to time is provided in fig. 5.2-1. The most relevant part of this information is also plotted in a world map as a function of ENVISAT position: % of cosmic ray hits per profile, % of datation errors per profile, % of star falling outside the central band per profile and % of saturation errors per profile (fig.5-2.2a).

It can be seen from fig. 5.2-1 that the cosmic rays hits occurred several times for the 99% of the measurements of the products. Looking at fig. 5.2-2a it can be clearly observed that this high percentage occurred when the satellite crossed the South Atlantic Anomaly (SAA) zone. Also the percentage of saturation errors per profile shows an increase over the SAA zone.

Another observation from fig. 5.2-1 is that for several products, 30-35% of the measurements have the star signal falling outside the central band. In fig. 5.2-2a it is observed that this percentage occurred mainly during twilight/dark conditions (roughly ascending) while in bright conditions the percentage is around 10% (fig.5.2-2a). This is because during the night the stars are lost deeper within the atmosphere and the turbulence phenomena becomes more important, producing the star to be less ‘focused’ on the spectrometers central band.

The other values (% of invalid measurements per product, % of measurements per product with datation errors...) are quite low.





The QWG has requested to perform a different plot of the cosmic rays in order to have a clear picture on the geographical position of the hits: count the cosmic rays detected in every product and when they are more than 100 then consider that cosmic rays have been detected. This plot is in fig. 5.2-2b. The products in bright limb have not been considered because the cosmic rays detection is not activated when processing products in bright.




Figure 5.2-2b: Count every time a cosmic ray has been detected. When it is > 100, then cosmic rays detected (yellow in the plot)

The flag information is given in table 5.2-1. The percentage of the products that have at least one measurement with demodulation flag set is also reported.

Table 5.2-1: Percentage of products during the reporting period with:

At least one measurement with demodulation flag set:	31 %
Reference spectrum computed from DB:	0.0 %
Reference spectrum with small number of measurements:	0.0 %
SATU data not used:	0.0 %

### 5.2.1 QUALITY FLAGS MONITORING (EXTRACTED FROM LEVEL 2 PRODUCTS)

In this section, the Product Quality information coming from the level 1 processing that is also stored in the level 2 products is plotted. Only products that did not have a fatal error (MPH error flag not set) are considered. The purpose of using the level 2 data is simply that the percentage of level 2 products arriving to the IDEAS team for the quality monitoring is much higher. For the reporting month, 100% of the archived products have been received. The plots are very similar to fig. 5.2-1 and 5.2-2a (demodulation flag information is not included) but separating ascending from descending passes. Since new version of the processor (GOMOS/5.00) there is no correspondence between illumination condition and latitude range when separating the passages (ascending and descending). Now, in the geo-location

process, the sun zenith angle is computed and the occultation is then flagged accordingly (dark, bright, twilight, straylight, twilight+straylight). You can see in fig. 5.2-3 the location of the occultations and their limb for the reporting month.

Fig. 5.2-4 and 5.2-5 present some quality information as a function of the time whereas in fig. 5.2-6 and 5.2-7 the plot is respect to the satellite position at the beginning of the occultations.

The percentage of measurements “where a problem occurred during the full transmission” per product ranges between 2 and 30 % (fig. 5.2-4, 5.2-5). The high values are due to the saturation that occurs mainly in bright limb. In dark limb the saturation occurs over the SAA zone but it is quite low elsewhere. From fig. 5.2-4 and 5.2-5 you can see also that there are a variable percentage of the measurements that have the star signal falling outside the central band. This is because in dark the stars are lost deeper within the atmosphere and the turbulence phenomena become more important, resulting in the star being less ‘focused’ on the spectrometers central band.

In ascending (fig. 5.2-6) the SAA is perfectly localized by the high percentage of cosmic ray hits per product (upper left panel). It is not the same if we look at fig. 5.2-7, because in descending most of the occultations in that world region are in bright limb conditions and the cosmic rays detection processing is not activated.




Figure 5.2-3: Position of the occultations based on illumination conditions




Figure 5.2-4: Level 1b product quality monitoring with respect to time ASCENDING ENVISAT passes






Figure 5.2-5: Level 1b product quality monitoring with respect to time DESCENDING ENVISAT passes



**Figure 5.2-6: Level 1b product quality monitoring with respect to satellite geo-location for ASCENDING ENVISAT passes**



**Figure 5.2-7: Level 1b product quality monitoring with respect to satellite geo-location for DESCENDING ENVISAT passes**

### 5.3 Spectral Performance

Every pixel of the spectrometers has a wavelength assigned. This assignment has been monitored through the mission by calculating, for given stars, the spectral shift corresponding to a maximum correlation between the reference star spectrum and the one of the occultation.

In order to have the wavelength well calibrated during the second reprocessing activity, the QWG performed a study to correct the spectral shift that was detected during the routine spectral performance monitoring (see fig. 5.3-1). A linear regression using data from stars 1 and 2 has been used to calibrate the wavelength for each needed orbit (one value for each calibration ADF used for the second reprocessing). This linear law took into account the ageing of the instrument. During the QWG #13, it has been decided to perform a wavelength calibration routinely with an extrapolation of this law and introducing also an extension to a second order law taking into account the seasonal variations. This routine calibration has been implemented on 14<sup>th</sup> December 2007 and is performed once a week at the same time of the DC maps calibration.

With this implementation the monitoring curve presented in fig. 5.3-1 should show small wavelength shifts since 14<sup>th</sup> December 2007. At least, the values should be smaller than the warning value set to 0.07 nm but, as it can be seen, the values have an unexpected variation (exceeding the threshold for a given period) that is currently being investigated by the QWG.




Figure 5.3-1: SPA2 wavelength monitoring since 12<sup>th</sup> November 2002: for every star ID (1, 2, 9, 18, 25) it is plotted the spectral shift for which a maximum correlation has been found between the reference spectrum and the one of the occultation

### 5.4 Radiometric Performance

#### 5.4.1 RADIOMETRIC SENSITIVITY

The monitoring performed consists of the calculation of the radiometric sensitivity of each CCD by computing the ratio between parts of the reference spectrum using specific stars (fig. 5.4-1).

The parts of the spectrum used are:

- UV: 250–300 nm;
- Yellow: 500–550 nm
- Red: 640–690 nm
- Ir1: 761-770 nm
- Ir2: 935-944 nm

For the spectrometers the ratios are with respect to the ‘yellow’ spectral range. For the photometers, the ratios are calculated by dividing the mean photometer signal above the atmosphere (115 km) by the ‘yellow’ spectral range (for PH1) or by the ‘red’ spectral range (for PH2). The variation of the ratio should be within a given threshold which is set to 10% (see table 5.4-1 that corresponds to fig. 5.4-1). For every star, this variation is calculated as the difference between the maximum (or minimum) ratio, and the mean over the 15 first values (if there were not 15 values computed yet, all values would be used).

**Table 5.4-1: Variation of RS for the different ratios (corresponds to fig. 5.4-1). Should be less than 10%**

Star Id	% Variation of UV ratio	% Variation of Red ratio	% Variation of IR1 ratio	% Variation of IR2 ratio	% Variation of Ph1 ratio	% Variation of Ph2 ratio
1	6.6	1.5	0.6	0.3	11.8	30.2
2	1.6	1.8	0.7	0.4	8.8	12.4
4	1.0	2.4	1.5	1.3	8.1	23.5
9	22.2	1.4	0.8	0.6	11.1	9.2
18	6.3	2.6	1.6	1.8	14.8	300.0
25	50.7	4.1	1.9	1.7	28.1	147.4

For star 9 and 25 the UV ratio is greater than the threshold 10%. It is clear (fig. 5.4-1) that there is a global decrease of UV ratios for all the stars. This confirms the expected degradation suffered by the UV optics that is, anyway, very small considering also the small variation for the rest of the stars (table 5.4-1).

By looking at the photometers radiometric sensitivity ratios of fig. 5.4-1, it can be seen that every star has a variation that seems to be annual. The variation is significant for stars 25 and 18. After some investigations performed by the QWG that exclude an inaccurate reflectivity correction LUT, it seems that the PH1/2 radiometric sensitivity variations could come from the fact that the spectrometers and the photometers are not illuminated the same way when the straylight appears (seasonal effect).




Figure 5.4-1: Radiometric sensitivity ratios since December 2002



### 5.4.2 PIXEL RESPONSE NON UNIFORMITY

No new PRNU calibration has been performed during the reporting period. This means that the PRNU maps inside the ADF remain as they are without any change for the moment.

## 5.5 Other Calibration Results

Future reports will address other calibration results, when available.

## 6 LEVEL 2 PRODUCT QUALITY MONITORING

### 6.1 Processor Configuration

#### 6.1.1 VERSION

Level 2 products from the operational ground segment have been disseminated during the reporting period to the users. Around 100% of GOM\_NL\_2P products have been received by the IDEAS team for routine quality control and long term trend monitoring. The current level 2-processor software version for the operational ground segment is GOMOS/5.00 since 8<sup>th</sup> August 2006 (see table 6.1-1). The product specification is PO-RS-MDA-GS2009\_10\_3I. Users are also supplied with 2002 - 4<sup>th</sup> July 2006 data sets reprocessed by the last prototype processor GOPR\_6.0c\_6.0f (developed and operated by ACRI) which is in line with the current GOMOS operational ground segment version GOMOS/5.00

**Table 6.1-1: PDS level 2 product version and main modifications implemented**

Date	Version	Description of changes
08-AUG-2006	Level 2 version 5.00 at PDHS-E and PDHS-K	Algorithm baseline level 2 DPM 6.2: <ul style="list-style-type: none"> <li>• The optimisation of the DOAS iterations</li> <li>• Negative column densities and local densities not flagged anymore</li> <li>• Suppress the setting of maximum error in case of negative local densities</li> <li>• Correction of HRTP discrepancies, and error estimates fixed</li> <li>• Rename Turbulence MDS into High Resolution Temperature MDS (HRTP)</li> <li>• Add vertical resolution per species in local densities MDS</li> <li>• Add Solar zenith angle at tangent point and at satellite level in geolocation ADS</li> <li>• Add "tangent point density from external</li> </ul>

<p>23-JUL-2006</p>	<p>Level 2 version 5.00 at FinCoPAC</p>	<ul style="list-style-type: none"> <li>• model" in geolocation ADS</li> <li>• Suppress contribution of "tangent point density from external model" in "local air density from GOMOS atmospheric profile" in geolocation ADS</li> </ul> <p>Change in configuration at the time of the switch over:</p> <ul style="list-style-type: none"> <li>○ 2<sup>nd</sup> order polynomial for aerosol</li> <li>○ Air fixed to ECMWF (local density set to 0 in the L2 products)</li> <li>○ Orphal cross-sections for O<sub>3</sub></li> <li>○ GOMOS cross-sections for other species</li> <li>○ Covariance matrix terms linked to air set to 0</li> <li>○ Air and NO<sub>2</sub> additional errors set to 0</li> </ul>
<p>23-MAR-2003</p>	<p>Level 2 version 4.02 at PDHS-E and PDHS-K</p>	<p>Algorithm baseline level 2 DPM 5.5:</p> <p>Section 3</p> <ul style="list-style-type: none"> <li>• Add references to technical notes on Tikhonov regularization</li> <li>• Change High level breakdown of modules: SMO/PFG</li> <li>• Change parameter: NFS in I2 ADF</li> <li>• Change parameter <math>\sigma_G</math> in I2 ADF (Table 3.4.1.1-II)</li> <li>• Change content of Level 2/res products – GAP</li> <li>• Change time sampling discretisation</li> <li>• Add covariance matrix explanation</li> </ul> <p>Section 5</p> <ul style="list-style-type: none"> <li>• Replace SMO by PFG VER-1/2: Depending on NFS, Apply either a Gaussian filter or a Tikhonov regularization to the vertical inversion matrix</li> <li>• Unit conversion applied on kernel matrix</li> <li>• Suppress VER-3</li> </ul> <p>Section 6</p> <ul style="list-style-type: none"> <li>• GOMOS Atmospheric Profile (GAP): not used in this version</li> <li>• Time sampling in equation (6.5.3.7-73)</li> </ul>
<p>31-MAY-2003</p>	<p>Level 2 version 4.00 at PDHS-E and PDHS-K</p>	<p>Algorithm baseline level 2 DPM 5.4:</p> <ul style="list-style-type: none"> <li>• Revision of some default values</li> <li>• Add a new parameter</li> <li>• Transmission model computation: suppress tests on valid pixels and species</li> <li>• Apply a Gaussian filter to the vertical inversion matrix</li> <li>• Very low signal values are substituted by threshold value</li> </ul>
<p>21-NOV-2002</p>	<p>Level 2 version 3.61 at PDHS-E and PDHS-K</p>	<p>Algorithm baseline level 2 DPM 5.3a:</p> <ul style="list-style-type: none"> <li>• Revision of some default values</li> <li>• Wording of test T11</li> <li>• Dilution term computation of jend</li> <li>• Covariance computation scaling applied before and after</li> </ul>

**Table 6.1-2: GOPR level 2 product version and main modifications implemented**

Date	Version	Description of changes
14-OCT-2005	GOPR_6.0f	<ul style="list-style-type: none"> <li>The optimisation of the DOAS iterations</li> <li>Negative column densities and local densities not flagged anymore</li> <li>Suppress the setting of maximum error in case of negative local densities</li> <li>Correction of H RTP discrepancies, and error estimates fixed</li> </ul> Configuration for second reprocessing: <ul style="list-style-type: none"> <li>2<sup>nd</sup> order polynomial for aerosol</li> <li>Air fixed to ECMWF (local density set to 0 in the L2 products)</li> <li>Orphal cross-sections for O<sub>3</sub></li> <li>GOMOS cross-sections for other species</li> <li>Covariance matrix terms linked to air set to 0</li> <li>Air and NO<sub>2</sub> additional errors set to 0</li> </ul>
17-MAR-2004	GOPR 6.0a	<ul style="list-style-type: none"> <li>Rename Turbulence MDS into High Resolution Temperature MDS (H RTP)</li> <li>Add vertical resolution per species in local densities MDS</li> <li>Add Solar zenith angle at tangent point and at satellite level in geolocation ADS</li> <li>Add "tangent point density from external model" in geolocation ADS</li> <li>Suppress contribution of "tangent point density from external model" in "local air density from GOMOS atmospheric profile" in geolocation ADS</li> </ul>
18-AUG-2003	GOPR 5.4d	<ul style="list-style-type: none"> <li>Tikhonov regularisation is implemented</li> </ul>
18-MAR-2003	GOPR 5.4b	<ul style="list-style-type: none"> <li>Modification to implement the computation of Tmodel for spectrometer B (in version 5.4b, the Tmodel for SPB is still set to 1)</li> </ul>
30-JAN-2003	GOPR 5.4a	<ul style="list-style-type: none"> <li>Modifications for ACRI internal use only. No impact on level 2 products.</li> </ul>

### 6.1.2 AUXILIARY DATA FILES (ADF)

The ADF's files in table 6.1-3 and 6.1-4 are used by the PDS to process the data from level 1 to level 2. For every type of file, the validity runs from the start validity time until the start validity time of the following one, but if an ADF file has been disseminated after the start validity time, it is obvious that it will be used by the PDHS-E and PDHS-K PDS only after the dissemination time (this happens the majority of the time). Note that the files outlined in yellow are the set of auxiliary files used during the reporting period.

**Table 6.1-3: Historic GOM\_PR2\_AX files used by PDS for level 2 products generation. The GOM\_PR2\_AX is a file containing the configuration parameters used for processing from level 1b to level 2 products**

Used by PDS for Level 2 products generation during	GOM_PR2_AX (GOMOS Processing level 2 configuration file)
01-MAR-2002 → 29-JUL-2002	<b>GOM_PR2_AXVIEC20020121_165624_20020101_000000_20200101_000000</b> <ul style="list-style-type: none"> <li>Pre-launch configuration</li> </ul>
30-JUL-2002 → 02-SEP-2002	<b>GOM_PR2_AXVIEC20020729_083851_20020301_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>Maximum value of chi2 before a warning flag is raised (set to 5)</li> <li>Maximum number of iterations for the main loop (set to 1)</li> </ul>
03-SEP-2002 → 12-NOV-2003	<b>GOM_PR2_AXVIEC20020902_151029_20020301_000000_20100101_000000</b> <ul style="list-style-type: none"> <li>Maximum value of chi2 before a warning flag is raised (set to 100)</li> </ul>

13-NOV-2003 → 22-MAR-2004	<p><b>GOM_PR2_AXVIEC20021112_170458_20020301_000000_20100101_000000</b></p> <ul style="list-style-type: none"> <li>• Smoothing mode</li> <li>• Hanning filter</li> <li>• Number of iterations</li> <li>• Spectral windows to suppress the O2 absorption in the high spectral range of SPA2</li> </ul>
<p>23-MAR-2004</p> <p><i>Note:</i> this file was used by the GOMOS/4.02 processors before the IECF dissemination. The dissemination was done on 25<sup>th</sup> March 2004</p>	<p><b>GOM_PR2_AXVIEC20040316_145613_20020301_000000_20100101_000000</b></p> <ul style="list-style-type: none"> <li>• Pressure at the top of the atmosphere</li> <li>• Number of GOMOS sources data (used in GAP)</li> <li>• Activation flag for GOMOS sources data (GAP)</li> <li>• Smoothing mode (after the spectral inversion)</li> <li>• Atmosphere thickness</li> </ul>
<p>08-AUG-2006</p> <p>Used at the time of switching over GOMOS/5.00</p>	<p><b>GOM_PR2_AXVIEC20051021_081111_20020301_000000_20100101_000000</b></p> <ul style="list-style-type: none"> <li>• Several level 2 processing configuration parameters</li> </ul>

**Table 6.1-4: Historic GOM\_CRS\_AX files used by PDS for level 2 products generation. The GOM\_CRS\_AX is a file containing the cross sections used for processing from level 1b to level 2 products**

Used by PDS for Level 2 products generation during	GOM_CRS_AX (GOMOS Cross Sections file)
01-MAR-2002 → 08-MAR-2002	<p><b>GOM_CRS_AXVIEC20020121_164026_20020101_000000_20200101_000000</b></p> <ul style="list-style-type: none"> <li>• Pre-launch configuration</li> </ul>
09-MAR-2003 → 29-JUL-2002	<p><b>GOM_CRS_AXVIEC20020308_185417_20020101_000000_20200101_000000</b></p> <ul style="list-style-type: none"> <li>• Corrected NUM_DSD in MPH - was 14 and is now 19 - and corrected spare DSD format by replacing last spare by carriage returns in file GOM_CRS_AXVIEC20020121_164026_20020101_000000_20200101_000000</li> </ul>
30-JUL-2002 → 25-MAR-2004	<p><b>GOM_CRS_AXVIEC20020729_082931_20020301_000000_20100101_000000</b></p> <ul style="list-style-type: none"> <li>• O3 cross-sections summary description (SPA)</li> <li>• NO3 cross-sections summary description</li> <li>• O2 transmissions summary description</li> <li>• H2O transmissions summary description</li> <li>• O3 cross sections (SPA)</li> </ul>
<p>26-MAR-2004</p> <p><i>Note:</i> the file was disseminated on 27 Jan 2004 but could not be used by PDS until version GOMOS/4.02 was in operation</p>	<p><b>GOM_CRS_AXVIEC20040127_150241_20020301_000000_20100101_000000</b></p> <ul style="list-style-type: none"> <li>• Update of the O2 and H2O transmissions (S.A input)</li> <li>• Extension by continuity of the O3 cross-section for SPB</li> </ul>
<p>08-AUG-2006</p> <p>Used at the time of switching over GOMOS/5.00</p>	<p><b>GOM_CRS_AXVIEC20051021_080452_20020301_000000_20100101_000000</b></p> <ul style="list-style-type: none"> <li>• Updated O<sub>3</sub> cross-sections</li> </ul>

### 6.1.3 RE-PROCESSING STATUS

The improvement of the GOMOS processing chain is a continuous on-going activity, not only for the processing algorithm but also for the instrument characterization data. In order to provide the best quality products to the users and due to the normal delay between algorithm specification and implementation in the operational PDS, it was decided to reprocess the GOMOS data using the Gopr prototype.

The second reprocessing activity covering years 2002-2006 (until 4<sup>th</sup> July 2006) using the prototype Gopr\_6.0c\_6.0f is completed. All reprocessed data can be retrieved via web query from

<http://www.enviport.org/gomos/index.jsp>. FTP access to bulk reprocessing results (one tar file of GOMOS products per day) is allowed from the D-PAC: <ftp://gomo2usr@ftp-ops.de.envisat.esa.int>.

## 6.2 Quality Flags Monitoring

In this section, some information contained in the Quality Summary data set of the level 2 products arrived during reporting period is shown. In particular, the percentage of flagged points per profile for the local species O<sub>3</sub>, H<sub>2</sub>O, NO<sub>2</sub> and NO<sub>3</sub> is depicted (fig. 6.2-1). Only products in dark limb illumination conditions and without fatal errors (error flag in the MPH set to "0") are used.




Figure 6.2-1: Percentage of flagged points per profile

The flagging strategy for GOMOS version GOMOS/5.00 foresees that a profile point is flagged when:

- The local density is greater than a given maximum value
- The line density is not valid. And it occurs when:
  - The acquisition from level 1b is not valid
  - There is no acquisition used for reference star spectrum
  - The line density is greater than a given maximum value

Only for species: air, aerosol, O<sub>3</sub>, NO<sub>2</sub>, NO<sub>3</sub>, OCIO

- No convergence after a given number of LMA iterations
- $\chi^2$  out of LMA is bigger than  $\chi^2$
- Failure of inversion

Only for species: O<sub>2</sub>, H<sub>2</sub>O

- Spectro B only: no convergence

- Spectro B only: data not available
- Spectro B only: covariance not available

There are points mainly between -60° and 0° latitude (fig. 6.2-1) because in this period of the year full dark illumination condition occultations (only those products have been used for these plots) are geo-located on that region. In summer, full dark illumination data are mainly in the Southern Hemisphere while in winter it is the contrary: full dark illumination occultations are found mainly in the Northern Hemisphere.

Looking at fig. 6.2-1, the most evident characteristic that can be observed is the high percentage of flagged points per profile for some H<sub>2</sub>O profiles. Users should be careful in using these data as the quality is only guaranteed for few stars. As a consequence of the new flagging strategy the percentage of flagged points per profile for O<sub>3</sub>, NO<sub>2</sub> and NO<sub>3</sub> is around 10-15%. It can be seen also that there are latitudinal bands with almost the same color (same percentages) mainly for H<sub>2</sub>O. This means that the percentages of flagged points per profile have a dependence on the stars that have been observed: a given star is always observed at the same latitude but at different longitude.

### 6.3 Other Level 2 Performance Issues

#### 6.3.1 MONTHLY OZONE AVERAGE

The plot presented in fig. 6.3-1 is the average of the Ozone values during the reporting month in a grid of 0.5 degrees in latitude per 1 km in altitude. Only occultations in dark limb have been used.

The latitude reduction is due to the azimuth window fixed to [25, 50] degrees. In this range it seems that there are less VCCS anomaly occurrences.





Figure 6.3-1: Average GOMOS O<sub>3</sub> profile during the reporting month: average in a grid of 0.5° latitude x 1 km altitude

### 6.3.2 OZONE DISPERSION MONITORING

This section is the output of a QWG request for the monitoring of the dispersion around the equator  $[-30^\circ, 30^\circ]$  using the brightest star of the day and with temperature greater or equal than 7000 k. This request includes the plot of daily median ozone, daily averaged  $\chi^2$ , daily averaged estimated errors and daily dispersion (defined as  $STD/Mean$  in %). The first step is the interpolation to given altitude layers (20, 25, 30, 40, 50, 60, 70, 80 and 90 km) and afterwards the daily average is performed. More than 5 profiles per day should be used for the average, if for a given day the number of profiles is less than 5 (for the brightest star) then the following star in increasing magnitude is chosen. The data above the SAA have not been used because those data produce unwanted fluctuations in the monitoring curves. The numbers below the lower curve are the star ID of the stars used for the statistics whilst the numbers above the upper curve are the number of profiles used.

Fig. 6.3-2 shows the daily ozone median for the reporting month. The daily averaged  $\chi^2$  is shown in fig. 6.3-3 while fig. 6.3-4 and fig. 6.3-5 show the daily average of estimated errors and the dispersion, respectively.



**Figure 6.3-2: Daily ozone average at different altitude layers for the reporting month**




Figure 6.3-3: Daily chi2 average at different altitude layers for the reporting month




Figure 6.3-4: Daily average of the estimated errors at different altitudes






Figure 6.3-5: Daily dispersion defined as STD/Mean in %

## 7 VALIDATION ACTIVITIES AND RESULTS

### 7.1 GOMOS-ECMWF Comparisons (Rossana Dragani, ECMWF input)

The full ECMWF validation report is available at the following link:

[http://earth.esa.int/pcs/envisat/calval\\_res/2009/ecmwf\\_gomos\\_monthly\\_200908\\_all.pdf](http://earth.esa.int/pcs/envisat/calval_res/2009/ecmwf_gomos_monthly_200908_all.pdf)

A summary of the report is reported in the following paragraph:

- The amount of observations measured by GOMOS in August 2009 was the highest received since January 2009, although it is not yet at the levels measured in the same period of 2008.
- The comparisons between the stratospheric temperature in the GOMOS BUFR files and the ECMWF temperature first-guess and analyses showed that in the global average and in the mean over the mid and high latitudes in the SH, the first-guess and analysis departures were within  $\pm 1\%$  (about  $\pm 2K$ ). In the tropics, the stratospheric first guess and analysis departures were about  $-0.5\%$  (about  $-1K$ ) at all levels. At mesospheric levels, the first guess and analysis departures were typically within  $-2$  and  $-4\%$  (within  $-4$  and  $-8K$ ) at all latitudes. The mean standard deviation of the first-guess and analysis departures were within  $0.5\%$  and  $4\%$  at all latitudes and levels.
- The ozone first guess and analysis departures were within  $-10$  and  $+20\%$  at all verticals levels in the global average and in the mean over the tropics and mid-latitudes in the SH, as well as in the mean in the upper stratosphere ( $p < 20$  hPa) and mesosphere at high latitudes in the SH. Larger departures ( $> 50\%$  in places) were found elsewhere. The standard deviations of the departures were larger than  $15\%$  at all levels.

- The comparisons between the GOMOS water vapour retrievals and the ECMWF water vapour first guess and analyses showed a poor level of agreement also for August 2009, with GOMOS water vapour observations being from one to three orders of magnitude larger than their model equivalent at all available vertical levels and latitudes.
- The monitoring statistics for August were produced with the operational ECMWF model, CY35R2.

## APPENDIX A

List of level 1b products that used the Calibration DC maps instead of the DSA observation (only ESRIN production):

Product name	DC information
GOM_TRA_1PNPDE20090803_195313_000000592081_00200_38831_6836.N1	DC map used
GOM_TRA_1PNPDE20090803_195800_000000622081_00200_38831_6837.N1	DC map used
GOM_TRA_1PNPDE20090803_200453_000000502081_00200_38831_6838.N1	DC map used
GOM_TRA_1PNPDE20090803_200855_000000492081_00200_38831_6839.N1	DC map used
GOM_TRA_1PNPDE20090803_202119_000000412081_00200_38831_6840.N1	DC map used
GOM_TRA_1PNPDE20090803_202428_000000382081_00200_38831_6841.N1	DC map used
GOM_TRA_1PNPDE20090803_202954_000000402081_00200_38831_6842.N1	DC map used
GOM_TRA_1PNPDE20090803_203606_000000522081_00200_38831_6843.N1	DC map used
GOM_TRA_1PNPDE20090803_203808_000000512081_00200_38831_6844.N1	DC map used
GOM_TRA_1PNPDE20090803_204443_000000492081_00200_38831_6845.N1	DC map used
GOM_TRA_1PNPDE20090803_204626_000000712081_00200_38831_6846.N1	DC map used
GOM_TRA_1PNPDE20090803_204840_000000442081_00200_38831_6847.N1	DC map used
GOM_TRA_1PNPDE20090803_205033_000000582081_00200_38831_6848.N1	DC map used
GOM_TRA_1PNPDE20090803_205347_000000572081_00200_38831_6849.N1	DC map used
GOM_TRA_1PNPDE20090803_205845_000000572081_00200_38831_6850.N1	DC map used
GOM_TRA_1PNPDE20090804_210708_000000612081_00215_38846_8180.N1	DC map used
GOM_TRA_1PNPDE20090804_211354_000000472081_00215_38846_8181.N1	DC map used
GOM_TRA_1PNPDE20090804_211752_000000492081_00215_38846_8182.N1	DC map used
GOM_TRA_1PNPDE20090804_213011_000000412081_00215_38846_8183.N1	DC map used
GOM_TRA_1PNPDE20090804_213321_000000422081_00215_38846_8184.N1	DC map used
GOM_TRA_1PNPDE20090804_213847_000000402081_00215_38846_8185.N1	DC map used
GOM_TRA_1PNPDE20090804_214453_000000492081_00215_38846_8186.N1	DC map used
GOM_TRA_1PNPDE20090804_214656_000000512081_00215_38846_8187.N1	DC map used
GOM_TRA_1PNPDE20090804_215336_000000502081_00215_38846_8188.N1	DC map used
GOM_TRA_1PNPDE20090804_215625_000000482081_00215_38846_8189.N1	DC map used
GOM_TRA_1PNPDE20090804_220302_000000472081_00215_38846_8190.N1	DC map used
GOM_TRA_1PNPDE20090804_220751_000000672081_00215_38846_8191.N1	DC map used
GOM_TRA_1PNPDE20090804_221337_000000442081_00215_38846_8192.N1	DC map used
GOM_TRA_1PNPDE20090805_203052_000000612081_00229_38860_9372.N1	DC map with no T dep.
GOM_TRA_1PNPDE20090805_203539_000000632081_00229_38860_9373.N1	DC map used
GOM_TRA_1PNPDE20090805_204218_000000482081_00229_38860_9374.N1	DC map used
GOM_TRA_1PNPDE20090805_204613_000000492081_00229_38860_9375.N1	DC map used
GOM_TRA_1PNPDE20090805_205828_000000412081_00229_38860_9376.N1	DC map used
GOM_TRA_1PNPDE20090805_210138_000000392081_00229_38860_9377.N1	DC map used
GOM_TRA_1PNPDE20090805_210705_000000412081_00229_38860_9378.N1	DC map used
GOM_TRA_1PNPDE20090805_211304_000000492081_00229_38860_9379.N1	DC map used
GOM_TRA_1PNPDE20090805_211509_000000512081_00229_38860_9380.N1	DC map used
GOM_TRA_1PNPDE20090805_212153_000000722081_00229_38860_9381.N1	DC map used
GOM_TRA_1PNPDE20090805_212339_000000532081_00229_38860_9382.N1	DC map used
GOM_TRA_1PNPDE20090805_212602_000000472081_00229_38860_9383.N1	DC map used
GOM_TRA_1PNPDE20090805_212753_000000512081_00229_38860_9384.N1	DC map used
GOM_TRA_1PNPDE20090805_213128_000000482081_00229_38860_9385.N1	DC map used
GOM_TRA_1PNPDE20090805_213620_000000682081_00229_38860_9386.N1	DC map used
GOM_TRA_1PNPDE20090805_214208_000000432081_00229_38860_9387.N1	DC map used
GOM_TRA_1PNPDE20090806_195922_000000582081_00243_38874_0231.N1	DC map used
GOM_TRA_1PNPDE20090806_200410_000000612081_00243_38874_0232.N1	DC map used
GOM_TRA_1PNPDE20090806_201042_000000472081_00243_38874_0233.N1	DC map used
GOM_TRA_1PNPDE20090806_201434_000000492081_00243_38874_0234.N1	DC map used

GOM_TRA_1PNPDE20090806_202646_000000402081_00243_38874_0235.NI	DC map used
GOM_TRA_1PNPDE20090806_202808_000000572081_00243_38874_0236.NI	DC map used
GOM_TRA_1PNPDE20090806_202956_000000412081_00243_38874_0237.NI	DC map used
GOM_TRA_1PNPDE20090806_203522_000000402081_00243_38874_0238.NI	DC map used
GOM_TRA_1PNPDE20090806_204116_000000502081_00243_38874_0239.NI	DC map used
GOM_TRA_1PNPDE20090806_204322_000000482081_00243_38874_0240.NI	DC map used
GOM_TRA_1PNPDE20090806_205010_000000722081_00243_38874_0241.NI	DC map used
GOM_TRA_1PNPDE20090806_205157_000000542081_00243_38874_0242.NI	DC map used
GOM_TRA_1PNPDE20090806_205425_000000452081_00243_38874_0243.NI	DC map used
GOM_TRA_1PNPDE20090806_205615_000000472081_00243_38874_0244.NI	DC map used
GOM_TRA_1PNPDE20090806_205953_000000482081_00243_38874_0245.NI	DC map used
GOM_TRA_1PNPDE20090806_210450_000000702081_00243_38874_0246.NI	DC map used
GOM_TRA_1PNPDE20090807_055924_000000542081_00249_38880_1152.NI	DC map used
GOM_TRA_1PNPDE20090807_060301_000000472081_00249_38880_1153.NI	DC map used
GOM_TRA_1PNPDE20090807_060749_000000612081_00249_38880_1154.NI	DC map used
GOM_TRA_1PNPDE20090807_061419_000000462081_00249_38880_1155.NI	DC map used
GOM_TRA_1PNPDE20090807_061809_000000482081_00249_38880_1156.NI	DC map used
GOM_TRA_1PNPDE20090807_062409_000000352081_00249_38880_1157.NI	DC map used
GOM_TRA_1PNPDE20090807_063019_000000422081_00249_38880_1158.NI	DC map used
GOM_TRA_1PNPDE20090807_063141_000000392081_00249_38880_1159.NI	DC map used
GOM_TRA_1PNPDE20090807_063329_000000422081_00249_38880_1160.NI	DC map used
GOM_TRA_1PNPDE20090807_063856_000000392081_00249_38880_1161.NI	DC map used
GOM_TRA_1PNPDE20090807_064447_000000492081_00249_38880_1162.NI	DC map used
GOM_TRA_1PNPDE20090807_064653_000000502081_00249_38880_1163.NI	DC map used
GOM_TRA_1PNPDE20090807_211318_000000612081_00258_38889_1671.NI	DC map used
GOM_TRA_1PNPDE20090807_211943_000000462081_00258_38889_1672.NI	DC map used
GOM_TRA_1PNPDE20090807_212331_000000672081_00258_38889_1673.NI	DC map used
GOM_TRA_1PNPDE20090807_212928_000000502081_00258_38889_1674.NI	DC map used
GOM_TRA_1PNPDE20090807_213539_000000392081_00258_38889_1675.NI	DC map used
GOM_TRA_1PNPDE20090807_213703_000000402081_00258_38889_1676.NI	DC map used
GOM_TRA_1PNPDE20090807_213850_000000402081_00258_38889_1677.NI	DC map used
GOM_TRA_1PNPDE20090807_214416_000000402081_00258_38889_1678.NI	DC map used
GOM_TRA_1PNPDE20090807_215003_000000512081_00258_38889_1679.NI	DC map used
GOM_TRA_1PNPDE20090807_215210_000000502081_00258_38889_1680.NI	DC map used
GOM_TRA_1PNPDE20090807_215902_000000712081_00258_38889_1681.NI	DC map used
GOM_TRA_1PNPDE20090807_220050_000000502081_00258_38889_1682.NI	DC map used
GOM_TRA_1PNPDE20090807_220323_000000492081_00258_38889_1683.NI	DC map used
GOM_TRA_1PNPDE20090807_220513_000000522081_00258_38889_1684.NI	DC map used
GOM_TRA_1PNPDE20090807_220855_000000482081_00258_38889_1685.NI	DC map used
GOM_TRA_1PNPDE20090807_221105_000000422081_00258_38889_1686.NI	DC map used
GOM_TRA_1PNPDE20090807_221356_000000622081_00258_38889_1687.NI	DC map used
GOM_TRA_1PNPDE20090807_221551_000000462081_00258_38889_1688.NI	DC map used
GOM_TRA_1PNPDE20090808_204149_000000592081_00272_38903_2938.NI	DC map used
GOM_TRA_1PNPDE20090808_204807_000000472081_00272_38903_2939.NI	DC map used
GOM_TRA_1PNPDE20090808_205153_000000462081_00272_38903_2940.NI	DC map used
GOM_TRA_1PNPDE20090808_205743_000000492081_00272_38903_2941.NI	DC map used
GOM_TRA_1PNPDE20090808_210357_000000392081_00272_38903_2942.NI	DC map used
GOM_TRA_1PNPDE20090808_210708_000000402081_00272_38903_2943.NI	DC map used
GOM_TRA_1PNPDE20090808_211234_000000402081_00272_38903_2944.NI	DC map used
GOM_TRA_1PNPDE20090808_211814_000000502081_00272_38903_2945.NI	DC map used
GOM_TRA_1PNPDE20090808_212023_000000472081_00272_38903_2946.NI	DC map used
GOM_TRA_1PNPDE20090808_212718_000000742081_00272_38903_2947.NI	DC map used
GOM_TRA_1PNPDE20090808_212908_000000572081_00272_38903_2948.NI	DC map used
GOM_TRA_1PNPDE20090808_213146_000000462081_00272_38903_2949.NI	DC map used
GOM_TRA_1PNPDE20090808_213334_000000582081_00272_38903_2950.NI	DC map used
GOM_TRA_1PNPDE20090808_213720_000000492081_00272_38903_2951.NI	DC map used

GOM_TRA_1PNPDE20090808_213932_000000442081_00272_38903_2952.NI	DC map used
GOM_TRA_1PNPDE20090808_214226_000000722081_00272_38903_2953.NI	DC map used
GOM_TRA_1PNPDE20090808_214420_000000462081_00272_38903_2954.NI	DC map used
GOM_TRA_1PNPDE20090809_200530_000000582081_00286_38917_4222.NI	DC map with no T dep.
GOM_TRA_1PNPDE20090809_201019_000000572081_00286_38917_4223.NI	DC map used
GOM_TRA_1PNPDE20090809_202014_000000462081_00286_38917_4224.NI	DC map used
GOM_TRA_1PNPDE20090809_202559_000000482081_00286_38917_4225.NI	DC map used
GOM_TRA_1PNPDE20090809_203215_000000402081_00286_38917_4226.NI	DC map used
GOM_TRA_1PNPDE20090809_203527_000000402081_00286_38917_4227.NI	DC map used
GOM_TRA_1PNPDE20090809_204052_000000402081_00286_38917_4228.NI	DC map used
GOM_TRA_1PNPDE20090809_204626_000000502081_00286_38917_4229.NI	DC map used
GOM_TRA_1PNPDE20090809_204835_000000482081_00286_38917_4230.NI	DC map used
GOM_TRA_1PNPDE20090809_205535_000000512081_00286_38917_4231.NI	DC map used
GOM_TRA_1PNPDE20090809_205726_000000562081_00286_38917_4232.NI	DC map used
GOM_TRA_1PNPDE20090809_210008_000000462081_00286_38917_4233.NI	DC map used
GOM_TRA_1PNPDE20090809_210156_000000532081_00286_38917_4234.NI	DC map used
GOM_TRA_1PNPDE20090809_210546_000000492081_00286_38917_4235.NI	DC map used
GOM_TRA_1PNPDE20090809_210759_000000432081_00286_38917_4236.NI	DC map used
GOM_TRA_1PNPDE20090809_211057_000000682081_00286_38917_4237.NI	DC map used
GOM_TRA_1PNPDE20090809_211249_000000472081_00286_38917_4238.NI	DC map used
GOM_TRA_1PNPDE20090809_211436_000000452081_00286_38917_4239.NI	DC map used
GOM_TRA_1PNPDE20090810_193402_000000572081_00300_38931_5500.NI	DC map used
GOM_TRA_1PNPDE20090810_193849_000000572081_00300_38931_5501.NI	DC map used
GOM_TRA_1PNPDE20090810_194456_000000472081_00300_38931_5502.NI	DC map used
GOM_TRA_1PNPDE20090810_194836_000000452081_00300_38931_5503.NI	DC map used
GOM_TRA_1PNPDE20090810_195057_000000312081_00300_38931_5504.NI	DC map used
GOM_TRA_1PNPDE20090810_195415_000000472081_00300_38931_5505.NI	DC map used
GOM_TRA_1PNPDE20090810_200034_000000392081_00300_38931_5506.NI	DC map used
GOM_TRA_1PNPDE20090810_200346_000000382081_00300_38931_5507.NI	DC map used
GOM_TRA_1PNPDE20090810_200911_000000382081_00300_38931_5508.NI	DC map used
GOM_TRA_1PNPDE20090810_201438_000000492081_00300_38931_5509.NI	DC map used
GOM_TRA_1PNPDE20090810_201648_000000492081_00300_38931_5510.NI	DC map used
GOM_TRA_1PNPDE20090810_202351_000000602081_00300_38931_5511.NI	DC map used
GOM_TRA_1PNPDE20090810_202543_000000562081_00300_38931_5512.NI	DC map used
GOM_TRA_1PNPDE20090810_202831_000000452081_00300_38931_5513.NI	DC map used
GOM_TRA_1PNPDE20090810_203017_000000322081_00300_38931_5514.NI	DC map used
GOM_TRA_1PNPDE20090810_203411_000000472081_00300_38931_5515.NI	DC map used
GOM_TRA_1PNPDE20090810_203625_000000452081_00300_38931_5516.NI	DC map used
GOM_TRA_1PNPDE20090810_203928_000000652081_00300_38931_5517.NI	DC map used
GOM_TRA_1PNPDE20090810_204119_000000512081_00300_38931_5518.NI	DC map used
GOM_TRA_1PNPDE20090811_204755_000000562081_00315_38946_6845.NI	DC map used
GOM_TRA_1PNPDE20090811_205356_000000542081_00315_38946_6846.NI	DC map used
GOM_TRA_1PNPDE20090811_205734_000000442081_00315_38946_6847.NI	DC map used
GOM_TRA_1PNPDE20090811_205951_000000522081_00315_38946_6848.NI	DC map used
GOM_TRA_1PNPDE20090811_210308_000000472081_00315_38946_6849.NI	DC map used
GOM_TRA_1PNPDE20090811_210928_000000412081_00315_38946_6850.NI	DC map used
GOM_TRA_1PNPDE20090811_211240_000000392081_00315_38946_6851.NI	DC map used
GOM_TRA_1PNPDE20090811_211805_000000392081_00315_38946_6852.NI	DC map used
GOM_TRA_1PNPDE20090811_212325_000000502081_00315_38946_6853.NI	DC map used
GOM_TRA_1PNPDE20090811_212536_000000482081_00315_38946_6854.NI	DC map used
GOM_TRA_1PNPDE20090811_213242_000000602081_00315_38946_6855.NI	DC map used
GOM_TRA_1PNPDE20090811_213435_000000422081_00315_38946_6856.NI	DC map used
GOM_TRA_1PNPDE20090811_213549_000000502081_00315_38946_6857.NI	DC map used
GOM_TRA_1PNPDE20090811_214313_000000492081_00315_38946_6858.NI	DC map used
GOM_TRA_1PNPDE20090811_214528_000000442081_00315_38946_6859.NI	DC map used
GOM_TRA_1PNPDE20090811_214835_000000702081_00315_38946_6860.NI	DC map used

GOM_TRA_1PNPDE20090811_215025_000000502081_00315_38946_6861.NI	DC map used
GOM_TRA_1PNPDE20090811_215221_000000472081_00315_38946_6862.NI	DC map used
GOM_TRA_1PNPDE20090811_215344_000000442081_00315_38946_6863.NI	DC map used
GOM_TRA_1PNPDE20090812_201625_000000562081_00329_38960_8165.NI	DC map used
GOM_TRA_1PNPDE20090812_202221_000000452081_00329_38960_8166.NI	DC map used
GOM_TRA_1PNPDE20090812_202556_000000432081_00329_38960_8167.NI	DC map used
GOM_TRA_1PNPDE20090812_202810_000000502081_00329_38960_8168.NI	DC map used
GOM_TRA_1PNPDE20090812_203125_000000452081_00329_38960_8169.NI	DC map used
GOM_TRA_1PNPDE20090812_203748_000000382081_00329_38960_8170.NI	DC map used
GOM_TRA_1PNPDE20090812_204624_000000402081_00329_38960_8171.NI	DC map used
GOM_TRA_1PNPDE20090812_205136_000000502081_00329_38960_8172.NI	DC map used
GOM_TRA_1PNPDE20090812_205348_000000492081_00329_38960_8173.NI	DC map used
GOM_TRA_1PNPDE20090812_210057_000000542081_00329_38960_8174.NI	DC map used
GOM_TRA_1PNPDE20090812_210408_000000482081_00329_38960_8175.NI	DC map used
GOM_TRA_1PNPDE20090812_210551_000000512081_00329_38960_8176.NI	DC map used
GOM_TRA_1PNPDE20090812_211138_000000522081_00329_38960_8177.NI	DC map used
GOM_TRA_1PNPDE20090812_211355_000000472081_00329_38960_8178.NI	DC map used
GOM_TRA_1PNPDE20090812_211707_000000772081_00329_38960_8179.NI	DC map used
GOM_TRA_1PNPDE20090812_211855_000000472081_00329_38960_8180.NI	DC map used
GOM_TRA_1PNPDE20090812_212051_000000462081_00329_38960_8181.NI	DC map used
GOM_TRA_1PNPDE20090812_212215_000000452081_00329_38960_8182.NI	DC map used
GOM_TRA_1PNPDE20090813_194454_000000542081_00343_38974_9599.NI	DC map used
GOM_TRA_1PNPDE20090813_195045_000000452081_00343_38974_9600.NI	DC map used
GOM_TRA_1PNPDE20090813_195418_000000442081_00343_38974_9601.NI	DC map used
GOM_TRA_1PNPDE20090813_195629_000000502081_00343_38974_9602.NI	DC map used
GOM_TRA_1PNPDE20090813_195942_000000672081_00343_38974_9603.NI	DC map used
GOM_TRA_1PNPDE20090813_200607_000000372081_00343_38974_9604.NI	DC map used
GOM_TRA_1PNPDE20090813_201442_000000392081_00343_38974_9605.NI	DC map used
GOM_TRA_1PNPDE20090813_201948_000000492081_00343_38974_9606.NI	DC map used
GOM_TRA_1PNPDE20090813_202201_000000492081_00343_38974_9607.NI	DC map used
GOM_TRA_1PNPDE20090813_202912_000000422081_00343_38974_9608.NI	DC map used
GOM_TRA_1PNPDE20090813_203225_000000392081_00343_38974_9609.NI	DC map used
GOM_TRA_1PNPDE20090813_203413_000000612081_00343_38974_9610.NI	DC map used
GOM_TRA_1PNPDE20090813_203702_000000392081_00343_38974_9611.NI	DC map used
GOM_TRA_1PNPDE20090813_204004_000000512081_00343_38974_9612.NI	DC map used
GOM_TRA_1PNPDE20090813_204222_000000472081_00343_38974_9613.NI	DC map used
GOM_TRA_1PNPDE20090813_204539_000000552081_00343_38974_9614.NI	DC map used
GOM_TRA_1PNPDE20090813_204726_000000502081_00343_38974_9615.NI	DC map used
GOM_TRA_1PNPDE20090814_053051_000000502081_00349_38980_0556.NI	DC map used
GOM_TRA_1PNPDE20090814_054050_000000542081_00349_38980_0557.NI	DC map used
GOM_TRA_1PNPDE20090814_054833_000000552081_00349_38980_0558.NI	DC map used
GOM_TRA_1PNPDE20090814_055422_000000452081_00349_38980_0559.NI	DC map used
GOM_TRA_1PNPDE20090814_055753_000000412081_00349_38980_0560.NI	DC map used
GOM_TRA_1PNPDE20090814_060002_000000502081_00349_38980_0561.NI	DC map used
GOM_TRA_1PNPDE20090814_060316_000000662081_00349_38980_0562.NI	DC map used
GOM_TRA_1PNPDE20090814_060941_000000372081_00349_38980_0563.NI	DC map used
GOM_TRA_1PNPDE20090814_061816_000000392081_00349_38980_0564.NI	DC map used
GOM_TRA_1PNPDE20090814_062319_000000492081_00349_38980_0565.NI	DC map used
GOM_TRA_1PNPDE20090814_062532_000000492081_00349_38980_0566.NI	DC map used
GOM_TRA_1PNPDE20090814_205400_000000532081_00358_38989_1005.NI	DC map used
GOM_TRA_1PNPDE20090814_205946_000000412081_00358_38989_1006.NI	DC map used
GOM_TRA_1PNPDE20090814_210316_000000422081_00358_38989_1007.NI	DC map used
GOM_TRA_1PNPDE20090814_210523_000000492081_00358_38989_1008.NI	DC map used
GOM_TRA_1PNPDE20090814_210836_000000472081_00358_38989_1009.NI	DC map used
GOM_TRA_1PNPDE20090814_211507_000000332081_00358_38989_1010.NI	DC map used
GOM_TRA_1PNPDE20090814_212337_000000392081_00358_38989_1011.NI	DC map used

GOM_TRA_1PNPDE20090814_212836_000000502081_00358_38989_1012.NI	DC map used
GOM_TRA_1PNPDE20090814_213048_000000482081_00358_38989_1013.NI	DC map used
GOM_TRA_1PNPDE20090814_214311_000000502081_00358_38989_1014.NI	DC map used
GOM_TRA_1PNPDE20090814_214602_000000422081_00358_38989_1015.NI	DC map used
GOM_TRA_1PNPDE20090814_214906_000000532081_00358_38989_1016.NI	DC map used
GOM_TRA_1PNPDE20090814_215125_000000462081_00358_38989_1017.NI	DC map used
GOM_TRA_1PNPDE20090814_215521_000000512081_00358_38989_1018.NI	DC map used
GOM_TRA_1PNPDE20090814_215829_000000482081_00358_38989_1019.NI	DC map used
GOM_TRA_1PNPDE20090814_215954_000000462081_00358_38989_1020.NI	DC map used
GOM_TRA_1PNPDE20090815_201456_000000662081_00372_39003_2203.NI	DC map with no T dep.
GOM_TRA_1PNPDE20090815_202229_000000542081_00372_39003_2204.NI	DC map used
GOM_TRA_1PNPDE20090815_202810_000000602081_00372_39003_2205.NI	DC map used
GOM_TRA_1PNPDE20090815_203139_000000422081_00372_39003_2206.NI	DC map used
GOM_TRA_1PNPDE20090815_203343_000000472081_00372_39003_2207.NI	DC map used
GOM_TRA_1PNPDE20090815_203654_000000462081_00372_39003_2208.NI	DC map used
GOM_TRA_1PNPDE20090815_205901_000000472081_00372_39003_2209.NI	DC map used
GOM_TRA_1PNPDE20090815_211133_000000622081_00372_39003_2210.NI	DC map used
GOM_TRA_1PNPDE20090815_211426_000000412081_00372_39003_2211.NI	DC map used
GOM_TRA_1PNPDE20090815_211953_000000472081_00372_39003_2212.NI	DC map used
GOM_TRA_1PNPDE20090815_212349_000000492081_00372_39003_2213.NI	DC map used
GOM_TRA_1PNPDE20090815_212700_000000502081_00372_39003_2214.NI	DC map used
GOM_TRA_1PNPDE20090815_212825_000000462081_00372_39003_2215.NI	DC map used
GOM_TRA_1PNPDE20090816_195058_000000532081_00386_39017_3849.NI	DC map used
GOM_TRA_1PNPDE20090816_195634_000000602081_00386_39017_3850.NI	DC map used
GOM_TRA_1PNPDE20090816_200001_000000432081_00386_39017_3851.NI	DC map used
GOM_TRA_1PNPDE20090816_200202_000000472081_00386_39017_3852.NI	DC map used
GOM_TRA_1PNPDE20090816_200513_000000442081_00386_39017_3853.NI	DC map used
GOM_TRA_1PNPDE20090816_202713_000000502081_00386_39017_3854.NI	DC map used
GOM_TRA_1PNPDE20090816_203954_000000472081_00386_39017_3855.NI	DC map used
GOM_TRA_1PNPDE20090816_204250_000000392081_00386_39017_3856.NI	DC map used
GOM_TRA_1PNPDE20090816_204557_000000502081_00386_39017_3857.NI	DC map used
GOM_TRA_1PNPDE20090816_204820_000000472081_00386_39017_3858.NI	DC map used
GOM_TRA_1PNPDE20090817_210003_000000532081_00401_39032_5158.NI	DC map used
GOM_TRA_1PNPDE20090817_210535_000000592081_00401_39032_5159.NI	DC map used
GOM_TRA_1PNPDE20090817_210900_000000582081_00401_39032_5160.NI	DC map used
GOM_TRA_1PNPDE20090817_211058_000000472081_00401_39032_5161.NI	DC map used
GOM_TRA_1PNPDE20090817_211407_000000442081_00401_39032_5162.NI	DC map used
GOM_TRA_1PNPDE20090817_213347_000000492081_00401_39032_5163.NI	DC map used
GOM_TRA_1PNPDE20090817_213518_000000272081_00401_39032_5164.NI	DC map used
GOM_TRA_1PNPDE20090817_213723_000000462081_00401_39032_5165.NI	DC map used
GOM_TRA_1PNPDE20090817_214852_000000632081_00401_39032_5166.NI	DC map used
GOM_TRA_1PNPDE20090817_215150_000000412081_00401_39032_5167.NI	DC map used
GOM_TRA_1PNPDE20090817_215459_000000512081_00401_39032_5168.NI	DC map used
GOM_TRA_1PNPDE20090817_215723_000000482081_00401_39032_5169.NI	DC map used
GOM_TRA_1PNPDE20090817_220124_000000542081_00401_39032_5170.NI	DC map used
GOM_TRA_1PNPDE20090817_220440_000000492081_00401_39032_5171.NI	DC map used
GOM_TRA_1PNPDE20090818_202116_000000612081_00415_39046_6444.NI	DC map with no T dep.
GOM_TRA_1PNPDE20090818_202831_000000512081_00415_39046_6445.NI	DC map used
GOM_TRA_1PNPDE20090818_203400_000000442081_00415_39046_6446.NI	DC map used
GOM_TRA_1PNPDE20090818_203723_000000582081_00415_39046_6447.NI	DC map used
GOM_TRA_1PNPDE20090818_203918_000000472081_00415_39046_6448.NI	DC map used
GOM_TRA_1PNPDE20090818_204226_000000432081_00415_39046_6449.NI	DC map used
GOM_TRA_1PNPDE20090818_210159_000000492081_00415_39046_6450.NI	DC map used
GOM_TRA_1PNPDE20090818_210328_000000382081_00415_39046_6451.NI	DC map used
GOM_TRA_1PNPDE20090818_210535_000000502081_00415_39046_6452.NI	DC map used
GOM_TRA_1PNPDE20090818_211713_000000642081_00415_39046_6453.NI	DC map used

GOM_TRA_1PNPDE20090818_212014_000000392081_00415_39046_6454.NI	DC map used
GOM_TRA_1PNPDE20090818_212325_000000542081_00415_39046_6455.NI	DC map used
GOM_TRA_1PNPDE20090818_212551_000000492081_00415_39046_6456.NI	DC map used
GOM_TRA_1PNPDE20090818_212954_000000552081_00415_39046_6457.NI	DC map used
GOM_TRA_1PNPDE20090818_213115_000000532081_00415_39046_6458.NI	DC map used
GOM_TRA_1PNPDE20090818_213312_000000512081_00415_39046_6459.NI	DC map used
GOM_TRA_1PNPDE20090818_213439_000000462081_00415_39046_6460.NI	DC map used
GOM_TRA_1PNPDE20090819_194950_000000602081_00429_39060_7759.NI	DC map with no T dep.
GOM_TRA_1PNPDE20090819_195700_000000502081_00429_39060_7760.NI	DC map used
GOM_TRA_1PNPDE20090819_200224_000000422081_00429_39060_7761.NI	DC map used
GOM_TRA_1PNPDE20090819_200546_000000412081_00429_39060_7762.NI	DC map used
GOM_TRA_1PNPDE20090819_200738_000000462081_00429_39060_7763.NI	DC map used
GOM_TRA_1PNPDE20090819_201046_000000432081_00429_39060_7764.NI	DC map used
GOM_TRA_1PNPDE20090819_203011_000000482081_00429_39060_7765.NI	DC map used
GOM_TRA_1PNPDE20090819_203347_000000482081_00429_39060_7766.NI	DC map used
GOM_TRA_1PNPDE20090819_204535_000000552081_00429_39060_7767.NI	DC map used
GOM_TRA_1PNPDE20090819_204837_000000462081_00429_39060_7768.NI	DC map used
GOM_TRA_1PNPDE20090819_205151_000000532081_00429_39060_7769.NI	DC map used
GOM_TRA_1PNPDE20090819_205419_000000512081_00429_39060_7770.NI	DC map used
GOM_TRA_1PNPDE20090820_210604_000000502081_00444_39075_9213.NI	DC map used
GOM_TRA_1PNPDE20090820_211125_000000412081_00444_39075_9214.NI	DC map used
GOM_TRA_1PNPDE20090820_211445_000000382081_00444_39075_9215.NI	DC map used
GOM_TRA_1PNPDE20090820_211635_000000442081_00444_39075_9216.NI	DC map used
GOM_TRA_1PNPDE20090820_211941_000000432081_00444_39075_9217.NI	DC map used
GOM_TRA_1PNPDE20090820_212812_000000462081_00444_39075_9218.NI	DC map used
GOM_TRA_1PNPDE20090820_213859_000000482081_00444_39075_9219.NI	DC map used
GOM_TRA_1PNPDE20090820_214113_000000472081_00444_39075_9220.NI	DC map used
GOM_TRA_1PNPDE20090820_215432_000000512081_00444_39075_9221.NI	DC map used
GOM_TRA_1PNPDE20090820_215737_000000432081_00444_39075_9222.NI	DC map used
GOM_TRA_1PNPDE20090820_220053_000000562081_00444_39075_9223.NI	DC map used
GOM_TRA_1PNPDE20090820_220323_000000522081_00444_39075_9224.NI	DC map used
GOM_TRA_1PNPDE20090820_220730_000000632081_00444_39075_9225.NI	DC map used
GOM_TRA_1PNPDE20090820_220857_000000592081_00444_39075_9226.NI	DC map used
GOM_TRA_1PNPDE20090820_221055_000000562081_00444_39075_9227.NI	DC map used
GOM_TRA_1PNPDE20090821_051243_000000532081_00449_39080_0084.NI	DC map used
GOM_TRA_1PNPDE20090821_052204_000000512081_00449_39080_0085.NI	DC map used
GOM_TRA_1PNPDE20090821_052906_000000522081_00449_39080_0086.NI	DC map used
GOM_TRA_1PNPDE20090821_053425_000000432081_00449_39080_0087.NI	DC map used
GOM_TRA_1PNPDE20090821_053744_000000532081_00449_39080_0088.NI	DC map used
GOM_TRA_1PNPDE20090821_053934_000000432081_00449_39080_0089.NI	DC map used
GOM_TRA_1PNPDE20090821_054240_000000402081_00449_39080_0090.NI	DC map used
GOM_TRA_1PNPDE20090821_055107_000000502081_00449_39080_0091.NI	DC map used
GOM_TRA_1PNPDE20090821_060155_000000472081_00449_39080_0092.NI	DC map used
GOM_TRA_1PNPDE20090821_060409_000000472081_00449_39080_0093.NI	DC map used
GOM_TRA_1PNPDE20090821_202734_000000602081_00458_39089_0640.NI	DC map used
GOM_TRA_1PNPDE20090821_203432_000000502081_00458_39089_0641.NI	DC map used
GOM_TRA_1PNPDE20090821_203950_000000422081_00458_39089_0642.NI	DC map used
GOM_TRA_1PNPDE20090821_204456_000000432081_00458_39089_0643.NI	DC map used
GOM_TRA_1PNPDE20090821_204801_000000442081_00458_39089_0644.NI	DC map used
GOM_TRA_1PNPDE20090821_205623_000000482081_00458_39089_0645.NI	DC map used
GOM_TRA_1PNPDE20090821_210712_000000452081_00458_39089_0646.NI	DC map used
GOM_TRA_1PNPDE20090821_210926_000000492081_00458_39089_0647.NI	DC map used
GOM_TRA_1PNPDE20090821_212253_000000562081_00458_39089_0648.NI	DC map used
GOM_TRA_1PNPDE20090821_212601_000000432081_00458_39089_0649.NI	DC map used
GOM_TRA_1PNPDE20090821_212748_000000592081_00458_39089_0650.NI	DC map used
GOM_TRA_1PNPDE20090821_212919_000000572081_00458_39089_0651.NI	DC map used



GOM_TRA_1PNPDE20090821_213151_000000532081_00458_39089_0652.NI	DC map used
GOM_TRA_1PNPDE20090821_213600_000000572081_00458_39089_0653.NI	DC map used
GOM_TRA_1PNPDE20090821_213731_000000592081_00458_39089_0654.NI	DC map used
GOM_TRA_1PNPDE20090822_200300_000000492081_00472_39103_2027.NI	DC map used
GOM_TRA_1PNPDE20090822_201316_000000602081_00472_39103_2028.NI	DC map used
GOM_TRA_1PNPDE20090822_201621_000000432081_00472_39103_2029.NI	DC map used
GOM_TRA_1PNPDE20090822_202435_000000492081_00472_39103_2030.NI	DC map used
GOM_TRA_1PNPDE20090822_203524_000000452081_00472_39103_2031.NI	DC map used
GOM_TRA_1PNPDE20090822_203738_000000492081_00472_39103_2032.NI	DC map used
GOM_TRA_1PNPDE20090822_205113_000000662081_00472_39103_2033.NI	DC map used
GOM_TRA_1PNPDE20090822_205425_000000432081_00472_39103_2034.NI	DC map used
GOM_TRA_1PNPDE20090822_205745_000000532081_00472_39103_2035.NI	DC map used
GOM_TRA_1PNPDE20090822_210019_000000492081_00472_39103_2036.NI	DC map used
GOM_TRA_1PNPDE20090825_200222_000000472082_00014_39146_7366.NI	DC map with no T dep.
GOM_TRA_1PNPDE20090825_200859_000000482082_00014_39146_7367.NI	DC map used
GOM_TRA_1PNPDE20090825_201857_000000532082_00014_39146_7368.NI	DC map used
GOM_TRA_1PNPDE20090825_202159_000000572082_00014_39146_7369.NI	DC map used
GOM_TRA_1PNPDE20090825_202948_000000462082_00014_39146_7370.NI	DC map used
GOM_TRA_1PNPDE20090825_204039_000000462082_00014_39146_7371.NI	DC map used
GOM_TRA_1PNPDE20090825_204251_000000472082_00014_39146_7372.NI	DC map used
GOM_TRA_1PNPDE20090825_205651_000000562082_00014_39146_7373.NI	DC map used
GOM_TRA_1PNPDE20090825_210012_000000452082_00014_39146_7374.NI	DC map used
GOM_TRA_1PNPDE20090825_210339_000000602082_00014_39146_7375.NI	DC map used
GOM_TRA_1PNPDE20090825_210637_000000442082_00014_39146_7376.NI	DC map used
GOM_TRA_1PNPDE20090825_211041_000000602082_00014_39146_7377.NI	DC map used
GOM_TRA_1PNPDE20090826_211131_000000532082_00029_39161_8776.NI	DC map used
GOM_TRA_1PNPDE20090826_211803_000000492082_00029_39161_8777.NI	DC map used
GOM_TRA_1PNPDE20090826_212754_000000592082_00029_39161_8778.NI	DC map used
GOM_TRA_1PNPDE20090826_213056_000000392082_00029_39161_8779.NI	DC map used
GOM_TRA_1PNPDE20090826_213838_000000442082_00029_39161_8780.NI	DC map used
GOM_TRA_1PNPDE20090826_214927_000000462082_00029_39161_8781.NI	DC map used
GOM_TRA_1PNPDE20090826_215139_000000482082_00029_39161_8782.NI	DC map used
GOM_TRA_1PNPDE20090826_220547_000000592082_00029_39161_8783.NI	DC map used
GOM_TRA_1PNPDE20090826_220911_000000452082_00029_39161_8784.NI	DC map used
GOM_TRA_1PNPDE20090826_221242_000000612082_00029_39161_8785.NI	DC map used
GOM_TRA_1PNPDE20090826_221541_000000462082_00029_39161_8786.NI	DC map used
GOM_TRA_1PNPDE20090826_221949_000000612082_00029_39161_8787.NI	DC map used
GOM_TRA_1PNPDE20090826_222138_000000642082_00029_39161_8788.NI	DC map used
GOM_TRA_1PNPDE20090827_204003_000000512082_00043_39175_0078.NI	DC map with no T dep.
GOM_TRA_1PNPDE20090827_204630_000000472082_00043_39175_0079.NI	DC map used
GOM_TRA_1PNPDE20090827_205616_000000412082_00043_39175_0080.NI	DC map used
GOM_TRA_1PNPDE20090827_205918_000000562082_00043_39175_0081.NI	DC map used
GOM_TRA_1PNPDE20090827_210244_000000252082_00043_39175_0082.NI	DC map used
GOM_TRA_1PNPDE20090827_210652_000000452082_00043_39175_0083.NI	DC map used
GOM_TRA_1PNPDE20090827_211740_000000462082_00043_39175_0084.NI	DC map used
GOM_TRA_1PNPDE20090827_211952_000000442082_00043_39175_0085.NI	DC map used
GOM_TRA_1PNPDE20090827_213407_000000602082_00043_39175_0086.NI	DC map used
GOM_TRA_1PNPDE20090827_213734_000000472082_00043_39175_0087.NI	DC map used
GOM_TRA_1PNPDE20090827_214108_000000632082_00043_39175_0088.NI	DC map used
GOM_TRA_1PNPDE20090827_214408_000000432082_00043_39175_0089.NI	DC map used
GOM_TRA_1PNPDE20090827_214822_000000712082_00043_39175_0090.NI	DC map used
GOM_TRA_1PNPDE20090828_045438_000000552082_00048_39180_0890.NI	DC map used
GOM_TRA_1PNPDE20090828_050306_000000492082_00048_39180_0891.NI	DC map used
GOM_TRA_1PNPDE20090828_050931_000000482082_00048_39180_0892.NI	DC map used
GOM_TRA_1PNPDE20090828_051915_000000402082_00048_39180_0893.NI	DC map used
GOM_TRA_1PNPDE20090828_052540_000000452082_00048_39180_0894.NI	DC map used

GOM_TRA_1PNPDE20090828_052948_000000672082_00048_39180_0895.NI	DC map used
GOM_TRA_1PNPDE20090828_054036_000000462082_00048_39180_0896.NI	DC map used
GOM_TRA_1PNPDE20090828_054248_000000462082_00048_39180_0897.NI	DC map used
GOM_TRA_1PNPDE20090828_200835_000000492082_00057_39189_1418.NI	DC map with no T dep.
GOM_TRA_1PNPDE20090828_201457_000000492082_00057_39189_1419.NI	DC map used
GOM_TRA_1PNPDE20090828_202438_000000412082_00057_39189_1420.NI	DC map used
GOM_TRA_1PNPDE20090828_203057_000000522082_00057_39189_1421.NI	DC map used
GOM_TRA_1PNPDE20090828_203506_000000442082_00057_39189_1422.NI	DC map used
GOM_TRA_1PNPDE20090828_204554_000000462082_00057_39189_1423.NI	DC map used
GOM_TRA_1PNPDE20090828_204804_000000472082_00057_39189_1424.NI	DC map used
GOM_TRA_1PNPDE20090828_210227_000000592082_00057_39189_1425.NI	DC map used
GOM_TRA_1PNPDE20090828_210558_000000472082_00057_39189_1426.NI	DC map used
GOM_TRA_1PNPDE20090828_211053_000000532082_00057_39189_1427.NI	DC map used
GOM_TRA_1PNPDE20090828_211236_000000462082_00057_39189_1428.NI	DC map used
GOM_TRA_1PNPDE20090828_211654_000000722082_00057_39189_1429.NI	DC map used
GOM_TRA_1PNPDE20090829_193707_000000552082_00071_39203_2826.NI	DC map used
GOM_TRA_1PNPDE20090829_194324_000000452082_00071_39203_2827.NI	DC map used
GOM_TRA_1PNPDE20090829_195300_000000392082_00071_39203_2828.NI	DC map used
GOM_TRA_1PNPDE20090829_195911_000000492082_00071_39203_2829.NI	DC map used
GOM_TRA_1PNPDE20090829_200321_000000442082_00071_39203_2830.NI	DC map used
GOM_TRA_1PNPDE20090829_201407_000000452082_00071_39203_2831.NI	DC map used
GOM_TRA_1PNPDE20090829_201617_000000472082_00071_39203_2832.NI	DC map used
GOM_TRA_1PNPDE20090829_203047_000000552082_00071_39203_2833.NI	DC map used
GOM_TRA_1PNPDE20090829_203421_000000512082_00071_39203_2834.NI	DC map used
GOM_TRA_1PNPDE20090829_203920_000000542082_00071_39203_2835.NI	DC map used
GOM_TRA_1PNPDE20090829_204104_000000502082_00071_39203_2836.NI	DC map used
GOM_TRA_1PNPDE20090830_205227_000000472082_00086_39218_4236.NI	DC map used
GOM_TRA_1PNPDE20090830_210159_000000392082_00086_39218_4237.NI	DC map used
GOM_TRA_1PNPDE20090830_210802_000000482082_00086_39218_4238.NI	DC map used
GOM_TRA_1PNPDE20090830_211212_000000432082_00086_39218_4239.NI	DC map used
GOM_TRA_1PNPDE20090830_212256_000000452082_00086_39218_4240.NI	DC map used
GOM_TRA_1PNPDE20090830_212505_000000452082_00086_39218_4241.NI	DC map used
GOM_TRA_1PNPDE20090830_213942_000000552082_00086_39218_4242.NI	DC map used
GOM_TRA_1PNPDE20090830_214320_000000492082_00086_39218_4243.NI	DC map used
GOM_TRA_1PNPDE20090830_214824_000000552082_00086_39218_4244.NI	DC map used
GOM_TRA_1PNPDE20090830_215009_000000472082_00086_39218_4245.NI	DC map used
GOM_TRA_1PNPDE20090831_202054_000000462082_00100_39232_5499.NI	DC map used
GOM_TRA_1PNPDE20090831_203617_000000472082_00100_39232_5500.NI	DC map used
GOM_TRA_1PNPDE20090831_204028_000000412082_00100_39232_5501.NI	DC map used
GOM_TRA_1PNPDE20090831_205110_000000432082_00100_39232_5502.NI	DC map used
GOM_TRA_1PNPDE20090831_205319_000000422082_00100_39232_5503.NI	DC map used
GOM_TRA_1PNPDE20090831_210802_000000612082_00100_39232_5504.NI	DC map used
GOM_TRA_1PNPDE20090831_211143_000000492082_00100_39232_5505.NI	DC map used
GOM_TRA_1PNPDE20090831_211651_000000542082_00100_39232_5506.NI	DC map used
GOM_TRA_1PNPDE20090831_211837_000000492082_00100_39232_5507.NI	DC map used
GOM_TRA_1PNPDE20090831_212312_000000722082_00100_39232_5508.NI	DC map used