



MERIS CYCLIC REPORT 54<sup>th</sup>  
 18<sup>th</sup> December 2006 - 22<sup>nd</sup> January 2007



Lake Urmia (Persian: دریاچه ارومیه) - MERIS full resolution scene - Salt lake in northwestern Iran

---

prepared by/*préparé par* DPQC MERIS Team and QWG  
 reference/*référence*  
 issue/*édition* 1  
 revision/*révision* 0  
 date of issue/*date d'édition* Friday 2<sup>nd</sup> February, 2007  
 status/*état*  
 Document type/*type de document* MERIS Cyclic Report  
 Distribution/*distribution*

---

**A P P R O V A L**

---

Title <i>titre</i>	MERIS Cyclic Report - Cycle 54 <sup>th</sup>	issue 1 revision 0 <i>issue révision</i>
author <i>auteur</i>	B. Josso	date 02/02/07 <i>date</i>
approved by <i>approuvé par</i>	L. Bourg	date 02/02/07 <i>date</i>

---

**C H A N G E L O G**

---

reason for change/ <i>raison du changement</i>	issue/ <i>issue</i>	revision/ <i>révision</i>	date/ <i>date</i>
------------------------------------------------	---------------------	---------------------------	-------------------

**C H A N G E R E C O R D**

---

reason for change/ <i>raison du changement</i>	page(s)/ <i>page(s)</i>	paragraph(s)/ <i>para- graphe(s)</i>
------------------------------------------------	-------------------------	------------------------------------------

## Contents

<b>1</b>	<b>INTRODUCTION</b>	<b>1</b>
1.1	Acronyms and abbreviations . . . . .	1
<b>2</b>	<b>SUMMARY</b>	<b>3</b>
<b>3</b>	<b>PROCESSOR VERSION AND PROCESSING CONFIGURATION</b>	<b>4</b>
3.1	MERIS Processor Release . . . . .	4
3.2	Auxiliary data files (ADF) . . . . .	4
3.3	Level 1/Level 2 Configuration (SciHiO2) . . . . .	5
3.4	Configuration Table Interface (CTI) . . . . .	5
3.5	Level 1 / Level 2 RR or FR products . . . . .	5
<b>4</b>	<b>PDS STATUS</b>	<b>7</b>
4.1	MERIS Level 0 products availability . . . . .	7
4.2	MERIS FR acquisitions . . . . .	9
4.3	MER_CA__0P Products . . . . .	13
<b>5</b>	<b>INSTRUMENT/DATA UNAVAILABILITY</b>	<b>13</b>
5.1	Instrument Unavailability . . . . .	13
5.2	Data Unavailability . . . . .	15
<b>6</b>	<b>CALIBRATION AND INSTRUMENT CHARACTERIZATION</b>	<b>16</b>
6.1	Calibration . . . . .	16
6.1.1	Radiometric calibration . . . . .	16
6.1.2	Spectral calibration . . . . .	16
6.1.3	Geolocation . . . . .	16
6.1.4	VEU Temperature Analysis . . . . .	16
6.1.5	Vicarious calibration results . . . . .	19
6.2	Instrument Characterization . . . . .	20
6.2.1	Instrument degradation . . . . .	20
6.2.2	Diffuser ageing . . . . .	20
6.2.3	Smile Effect . . . . .	20
6.2.4	Spectral evolution from erbium measurements . . . . .	20
<b>7</b>	<b>DATA QUALITY CONTROL</b>	<b>20</b>
7.1	MERIS products quality status . . . . .	20
7.2	Anomalies and Software Problem Reporting (SPR) . . . . .	21
<b>8</b>	<b>FIRST 2003 MERIS ARCHIVE REPROCESSING</b>	<b>21</b>
<b>9</b>	<b>SECOND 2005 MERIS ARCHIVE REPROCESSING</b>	<b>21</b>
<b>10</b>	<b>MERIS PROCESSOR EVOLUTION</b>	<b>21</b>
<b>11</b>	<b>VALIDATION ACTIVITIES AND RESULTS</b>	<b>21</b>

<b>12 WATER VAPOUR AND BROWSE MAPS</b>	<b>21</b>
<b>13 HOW TO GET MERIS DATA</b>	<b>22</b>
<b>14 GENERAL INFORMATION</b>	<b>22</b>

## List of Tables

1	Cycle Characteristics . . . . .	3
2	MERIS processor parameters - version 5.02 . . . . .	4
3	Auxiliary Data Files in use for the cycle #54 . . . . .	4
4	MERIS Level 1 Auxiliary Data Files . . . . .	5
5	MERIS Level 2 Auxiliary Data Files . . . . .	5
6	Reduced Resolution Level 0 products availability . . . . .	7
7	Full Resolution Level 0 products availability . . . . .	8
8	Available calibration files . . . . .	13
10	Specific sites results . . . . .	20

## List of Figures

1	MER_RR__0P generated/missing by the ground segment during cycle #54 .	7
2	MER_FR__0P generated/missing by the ground segment during cycle #54 .	8
3	MERIS Full Resolution Level 0 acquisitions - Part #1 - 19/12/2006 - 23/12/2006	9
4	MERIS Full Resolution Level 0 acquisitions - Part #2 - 24/12/2006 - 28/12/2006	10
5	MERIS Full Resolution Level 0 acquisitions - Part #3 - 29/12/2006 - 02/01/2007	10
6	MERIS Full Resolution Level 0 acquisitions - Part #4 - 03/01/2007 - 07/01/2007	11
7	MERIS Full Resolution Level 0 acquisitions - Part #5 - 08/01/2007 - 12/01/2007	11
8	MERIS Full Resolution Level 0 acquisitions - Part #6 - 13/01/2007 - 17/01/2007	12
9	MERIS Full Resolution Level 0 acquisitions - Part #7 - 18/01/2007 - 22/01/2007	12
10	VEU Temperature during cycle #54 . . . . .	17
11	24/12/2006 VEU temperature increase . . . . .	18
12	METRIC calibration site map . . . . .	19

# 1 INTRODUCTION

The MERIS Cyclic Report is distributed by ESRIN- DPQC (Data Processing Quality Control) to keep the MERIS Community informed of any modification regarding the processor, updates of auxiliary products, anomalies of the instrument behaviour, data acquisition and processing, and finally the status of the calibration, validation, and quality control activities. The Cyclic Report collects the inputs coming from different groups involved in MERIS data exploitation:

- ESRIN- Product Control Facility (PCF)
- Quality Working Group (QWG)
- MERIS/AATSR validation team (MAVT)
- Brockmann Consult (BC)
- ACRI-st
- Laboratoire d'Océanographie de Villefranche (LOV)
- Centre National d'Études Spatiales (CNES)
- Frei Universitat Berlin (FUB)
- Laboratoire Interdisciplinaire en Sciences de l'Environnement (LISE)

The main objective of the Cyclic Report is to provide the users community with useful information regarding the instrument performances, the data production chain, the results of calibration activities and validation campaigns, at the end of each ENVISAT cycle, which represents 501 orbits, about 35 days.

## 1.1 Acronyms and abbreviations

ADF	Auxiliary Data File
ADS	Auxiliary Data Server
ARF	Archiving Facility (PDS)
CNES	Centre National d'Études Spatiales
CTI	Configuration Table Interface
CR	Cyclic Report
DAC	Diffuser Ageing Calibration
DMOP	Detailed Mission Operation Plan
DOY	Day Of Year
DS	Data Server
DSD	Data Set Descriptor
EDAC	Error Detection And Correction
FR	Full Resolution
FUB	Freie Universitat Berlin

---

GS	Ground Segment
IAT	Interactive Analysis Tool
IDL	Interactive Data Language
IECF	Instrument Engineering and Calibration Facilities
IPF	Instrument Processing Facilities (PDS)
INV	Inventory Facilities (PDS)
JRC	Joint Research Centre
LAN	Local Area Network
LISE	Laboratoire Interdisciplinaire en Sciences de l'Environnement
LOV	Laboratoire d'Océanographie de Villefranche-sur-mer
MERIS	Medium Resolution Image Spectrometer
MPH	Main Product Header
OP	Operational Phase of ENVISAT
OCL	Offset Control Loop
PAC	Processing and Archiving Centre (PDS)
PDCC	Payload Data Control Centre (PDS)
PDHS	Payload Data Handling Station (PDS)
PDS	Payload Data Segment
PEP	Payload Exploitation Plan
QC	Quality Control
QWG	Quality Control Working Group
QUARC	Quality Analysis and Reporting Computer
RGC	Radiometric Gain Calibration
RR	Reduced Resolution
SEU	Single Event Upset
SPH	Specific Product Header
SQADS	Summary Quality ADS
WV1	Wavelength type 1 calibration
WV2	Wavelength type 2 calibration

## 2 SUMMARY

Cycle #54 starts on the 18<sup>th</sup> December 2006 and stops on the 22<sup>nd</sup> January 2007 .

- A diffuser Ageing calibration, a Wavelength Type 1 calibration as well as two Radiometric calibrations were planned and successfully executed during cycle #54.
- No any new auxiliary files were disseminated during the reporting period.
- Three instrument unavailability periods occurred during the reporting period. The longest period of unavailability lasted more than six hours.
- One data unavailability period occurred during the reporting period.

Details about the start and stop of the cycle can be found in [Table 1](#).

Cycle number	54
Start time	18 <sup>th</sup> December 2006 , 21:59:29
Stop time	22 <sup>nd</sup> January 2007 , 21:59:29
Start orbit	25105
Stop orbit	25605

Table 1: Cycle Characteristics

## 3 PROCESSOR VERSION AND PROCESSING CONFIGURATION

### 3.1 MERIS Processor Release

No changes in the IPF have been performed during cycle #54. The current MERIS processor configuration is described in [Table 2](#).

IPF Version	Validity	Reference Documents
5.02	8 <sup>th</sup> May 2006 08:00 UTC Orbit # 21890 →	<ol style="list-style-type: none"> <li>1. ENVISAT Product Specification [Iss_5_Rev_A]</li> <li>2. MERIS Input/Output Data Definition [Iss_7_Rev_3a]</li> <li>3. MERIS Level 1b Detailed Processing Model [Iss_7_Rev_0a]</li> <li>4. MERIS Level 2b Detailed Processing Model [Iss_7_Rev_2a]</li> </ol>

Table 2: MERIS processor parameters - version 5.02

### 3.2 Auxiliary data files (ADF)

Auxiliary Data Files in use for the cycle #54 are listed [Table 3](#). Note: the other files not

Product description	Product name	Comment
<b>Level 1 aux files</b>		
Instrument Characterization Data	MER_INS	No change
Processing Level 1 Control Parameters data	MER_CP1	No change
Radiometric Calibration data	MER_RAC	No change
Digital Roughness Model	MER_DRM	No change
Digital Elevation Model	AUX_DEM	No change
Land Surface Map	AUX_LSM	No change
Attitude data file	AUX_ATT	No change
<b>Level 2 aux files</b>		
Aerosol Climatology data	MER_AER	No change
Atmosphere Parameter data	MER_ATP	No change
Cloud Measurement Parameters data	MER_CMP	No change
Processing Level-2 Control Parameters data	MER_CP2	No change
Land Aerosols Parameters data	MER_LAP	No change
Land Vegetation Index parameters data	MER_LVI	No change
Ocean Aerosols Parameters data	MER_OAP	No change
Ocean I parameters data	MER_OC1	No change
Ocean II parameters data	MER_OC2	No change
Water Vapour Parameters	MER_WVP	No change

Table 3: Auxiliary Data Files in use for the cycle #54

included into the list change every time (ECMWF).



### 3.3 Level 1/Level 2 Configuration (SciHiO2)

The current operational ADF files, used in the processing from Level 0 data to Level 1b or Level 2 products, are listed in [Table 4](#) and [Table 5](#).

- Level 1 ADF configuration:

Product name	Start Validity
AUX_ATT_AXVIEC20020924_131534_20020703_120000_20781231_235959	03/07/2002
AUX_DEM_AXVIEC20031201_000000_20031201_000000_20200101_000000	01/12/2003
AUX_LSM_AXVIEC20020123_141228_20020101_000000_20200101_000000	01/01/2002
MER_CP1_AXVIEC20050607_065745_20020321_193100_20120321_193100	21/03/2002
MER_DRM_AXVIEC20020122_083343_20020101_000000_20200101_000000	01/01/2002
MER_INS_AXVIEC20050708_134312_20050101_000000_20150101_000000	01/01/2005
MER_RAC_AXVIEC20061009_084736_20061009_220000_20161009_220000	09/10/2006

Table 4: MERIS Level 1 Auxiliary Data Files

- Level 2 ADF configuration:

Product name	Start Validity
MER_AER_AXVIEC20040407_174356_20020321_193100_20120321_193100	21/03/2002
MER_ATP_AXVIEC20050628_123340_20021224_121445_20121224_121445	24/12/2002
MER_CMP_AXVIEC20040407_180835_20021224_121445_20121224_121445	24/12/2002
MER_CP2_AXVIEC20050704_065814_20021224_121445_20121224_121445	24/12/2002
MER_LAP_AXVIEC20050628_124246_20020321_193100_20120321_193100	21/03/2002
MER_LVI_AXVIEC20050704_145357_20020321_193100_20120321_193100	21/03/2002
MER_OAP_AXVIEC20050704_145633_20020321_193100_20120321_193100	21/03/2002
MER_OC1_AXVIEC20050704_145802_20020321_193100_20120321_193100	21/03/2002
MER_OC2_AXVIEC20050628_123950_20020321_193100_20120321_193100	21/03/2002
MER_SCM_AXVIEC20030620_120000_20020321_193100_20110725_103844	21/03/2002
MER_WVP_AXVIEC20040407_181941_20020321_193100_20120321_193100	21/03/2002

Table 5: MERIS Level 2 Auxiliary Data Files

### 3.4 Configuration Table Interface (CTI)

No any new CTI was disseminated during cycle#54.

### 3.5 Level 1 / Level 2 RR or FR products

During cycle #54 neither format changes nor algorithm modifications regarding MERIS RR and FR products were implemented into the operational processor.

REMINDER:

In the middle of cycle #47, some format changes or algorithm modifications regarding MERIS

RR and FR products were implemented during the operational processor upgrade from v4.10 to 5.02. The data changes decided within the Data Quality Working Group are listed below:

- ▷ New Chlorophyll 1 polynomial characterisation from LOV (Laboratoire d'Océanologie de Villefranche - France)
- ▷ Chlorophyll 1 validity range set to [0.01, 30.], no PCD raise when out of range
- ▷ Troposphere-free MAR99 replaces BLUE –  $\leq 1.5$  (from previous BOMEM runs)
- ▷ Gothic R Look Up Table from LOV (Laboratoire d'Océanologie de Villefranche - France)
- ▷ Chlorophyll 2 conversion factors from GKSS (revised with latest Neural Network delivery)
- ▷ Yellow Substance coding offset and scaling factor changes (linear to log scale, same range)
- ▷ Chlorophyll coding range changes ( $[-2, 2]$  in  $\log_{10}$  scale instead of  $[-3, 3]$  previously)
- ▷ Whitecaps threshold set to  $10 \text{ m.s}^{-1}$
- ▷ New Case 2 Neural Network from GKSS (with and without linear reflectances as input)
- ▷ White scatterer threshold set to 4.8
- ▷ MTCI threshold on B13-B8 difference set to 0.05, on B10-B8 to  $1e^{-6}$  (numerical purpose only), ceiling for B8 set to 0.3, floor for B9 to 0.1
- ▷ Preliminary version of LARS Look Up Tables from Hygeos

For further details concerning the changes, please refer to the documentation available at: [http://earth.esa.int/pcs/envisat/meris/documentation/MERIS\\_IPF\\_evolution.pdf](http://earth.esa.int/pcs/envisat/meris/documentation/MERIS_IPF_evolution.pdf)

## 4 PDS STATUS

The statistics resulting from the query to the PDS inventory facility (INV) for the MERIS products availability are presented in the following paragraphs.

### 4.1 MERIS Level 0 products availability

Table 6 shows the statistics regarding the RR L0 availability (compared to the planned production).

Week	MER_RR__OP %	
	Inventoried	Missing
From 18/12 to 25/12	92.85	7.15
From 25/12 to 01/01	97.74	2.15
From 01/01 to 08/01	77.00	23.00
From 08/01 to 15/01	99.43	0.57
From 15/01 to 22/01	99.98	0.02

Table 6: Reduced Resolution Level 0 products availability




Figure 1: MER\_RR\_\_OP generated/missing by the ground segment during cycle #54

Table 7 shows the statistics regarding the FR L0 availability (compared to the planned production).

Week	MER_RR_OP %	
	Inventoried	Missing
From 18/12 to 25/12	89.97	10.03
From 25/12 to 01/01	87.58	12.23
From 01/01 to 08/01	91.74	8.26
From 08/01 to 15/01	98.73	1.27
From 15/01 to 22/01	99.08	0.92

Table 7: Full Resolution Level 0 products availability




Figure 2: MER\_FR\_OP generated/missing by the ground segment during cycle #54

## 4.2 MERIS FR acquisitions

The pictures below (Figure 3, Figure 4, Figure 5, Figure 6, Figure 7, Figure 8 and Figure 9) show the MERIS Full Resolution global coverage for the reporting period. As specified for this type of MERIS products, all lands and coastal areas are covered by MERIS FR acquisitions.




Figure 3: MERIS Full Resolution Level 0 acquisitions - Part #1 - 19/12/2006 - 23/12/2006




Figure 4: MERIS Full Resolution Level 0 acquisitions - Part #2 - 24/12/2006 - 28/12/2006




Figure 5: MERIS Full Resolution Level 0 acquisitions - Part #3 - 29/12/2006 - 02/01/2007






Figure 6: MERIS Full Resolution Level 0 acquisitions - Part #4 - 03/01/2007 - 07/01/2007




Figure 7: MERIS Full Resolution Level 0 acquisitions - Part #5 - 08/01/2007 - 12/01/2007




Figure 8: MERIS Full Resolution Level 0 acquisitions - Part #6 - 13/01/2007 - 17/01/2007




Figure 9: MERIS Full Resolution Level 0 acquisitions - Part #7 - 18/01/2007 - 22/01/2007



### 4.3 MER\_CA\_\_0P Products

During the Cycle #54, the following calibration campaigns were successfully executed:

- ▷ A Diffuser Ageing calibration was executed in orbits #25258-25259 (29<sup>th</sup> of December).
- ▷ A Wavelength Type 1 calibration was executed in orbits #25260 to #25261 (29<sup>th</sup> of December). This calibration required MERIS in Stabilization Mode during Region 1.
- ▷ A Radiometric Calibration was executed in orbit #25263 (29<sup>th</sup> of December 2006).
- ▷ A Radiometric Calibration was executed on doy 012 (Friday the 12<sup>th</sup> of January 2007) at 23.32.11z during orbit #25463.

The list of corresponding available calibration files is reported in [Table 8](#).

MER_CA__0PNPDK20061229_155040_000001772054_00154_25258_0068.N1	DAC
MER_CA__0PNPDK20061229_173117_000001772054_00155_25259_0069.N1	DAC
MER_CA__0PNPDK20061229_191149_000001782054_00156_25260_0070.N1	WV1
MER_CA__0PNPDE20061229_205226_000001772054_00157_25261_0040.N1	WV1
MER_CA__0PNPDE20061230_001339_000001772054_00159_25263_0042.N1	RC
MER_CA__0PNPDE20070112_233251_000001772054_00359_25463_0043.N1	RC

Table 8: Available calibration files

## 5 INSTRUMENT/DATA UNAVAILABILITY

### 5.1 Instrument Unavailability

The following instrument unavailability has been reported during cycle #54:

- MERIS SDPSS unexpectedly switched to PAUSE mode From 20<sup>th</sup> Jan 2007 20:28:53.000. It came back in acquisition at 20.29.25. (Day of Year = 020 Orbit = 25576 Anx Offset = 2843 To 20 Jan 2007 20:29:25.000 Day of Year = 020 Orbit = 25576 Anx Offset = 2875). As a result 32" of data loss occurred.

In the reporting period, 60 EDAC-corrected SEU occurred. The dates, times and geolocation of these events are given in the table below:

Date/Time year.day.hr.mn.sc	Lon.	Lat.
2006.352.00.06.25	28.5628° W	19.3060° S
2006.353.12.21.44	39.4254° W	25.1940° S
2006.353.14.02.00	64.2791° W	24.0187° S
2006.353.15.41.05	88.1220° W	18.6461° S
...	...	...

Date/Time year.day.hr.mn.sc	Lon.	Lat.
2006.354.00.14.29	134.0097° E	55.1818° S
2006.354.13.30.19	56.3188° W	23.7952° S
2006.354.22.32.51	5.9252° W	14.7110° S
2006.355.10.08.28	173.4914° W	44.9737° S
2006.355.12.59.02	48.7142° W	24.9890° S
2006.355.13.00.35	50.1520° W	30.4702° S
2006.356.12.27.39	41.0232° W	25.8290° S
2006.358.23.32.03	0.4983° W	68.6500° S
2006.360.00.54.00	39.6985° W	23.7256° S
2006.361.11.29.51	26.3864° W	24.7936° S
2006.361.13.08.11	49.5910° W	16.7610° S
2006.362.11.16.39	114.2251° W	81.4669° S
2006.362.19.40.34	99.3698° E	80.5966° S
2006.363.01.00.53	42.1290° W	19.6458° S
2006.363.02.38.02	64.1471° W	31.8526° S
2006.363.02.41.46	67.5195° W	18.6346° S
2006.363.12.11.29	40.0118° W	39.8843° S
2006.363.13.46.04	59.3726° W	18.6380° S
2006.363.22.40.09	0.1121° E	49.6079° S
2006.364.00.29.40	34.5663° W	18.2060° S
2006.364.02.04.53	54.6659° W	37.2358° S
2006.365.03.19.47	77.4121° W	16.2798° S
2007.001.12.11.14	35.6861° W	18.7583° S
2007.002.00.35.50	36.3645° W	16.6561° S
2007.003.00.03.35	27.9353° W	18.8819° S
2007.003.01.41.24	50.6189° W	28.7375° S
2007.004.12.16.48	36.9835° W	18.1810° S
2007.005.23.56.59	12.9116° W	61.9735° S
2007.006.12.57.59	49.7731° W	31.7470° S
2007.006.14.33.42	70.6042° W	14.4537° S
2007.010.10.46.54	14.0344° W	15.2538° S
2007.010.14.12.55	68.6279° W	32.3258° S
2007.011.11.58.31	33.5292° W	24.6145° S
2007.011.13.32.50	53.5272° W	2.2922° S
2007.011.13.34.14	54.6273° W	7.2760° S
2007.012.03.42.24	82.8607° W	17.6026° S
2007.012.13.03.25	47.3616° W	10.1356° S
2007.012.13.23.29	97.0654° W	78.3110° S
2007.014.02.36.27	64.6881° W	27.2051° S
...	...	...

Date/Time year.day.hr.mn.sc	Lon.	Lat.
2007.014.05.37.02	28.4057° E	76.1707° S
2007.015.11.29.38	24.5183° W	13.9689° S
2007.015.13.13.59	52.8932° W	27.2739° S
2007.015.23.54.33	25.4456° W	20.3348° S
2007.016.12.38.34	41.7277° W	13.8103° S
2007.017.02.44.54	68.4934° W	17.5900° S
2007.017.13.17.23	66.8869° E	80.5475° N
2007.018.04.16.09	109.6839° W	61.0460° N
2007.018.13.15.13	50.4980° W	11.2958° S
2007.018.14.57.35	77.0924° W	17.5787° S
2007.018.15.01.53	80.9791° W	32.8008° S
2007.018.16.55.03	138.5879° W	75.3479° S
2007.018.23.59.47	26.4498° W	22.1203° S
2007.019.01.43.11	53.9342° W	12.1734° S
2007.019.12.48.11	46.5040° W	27.5703° S
2007.019.14.24.17	67.8150° W	11.6117° S
2007.020.10.33.29	11.2685° W	18.7800° S

## 5.2 Data Unavailability

The following data Unavailability related either to Envisat or Artemis were reported during cycle #54

- At 2006.362.00.07.17z, ARTEMIS triggered a spurious anomaly Level2A, which disconnected the APC from the bus. The APC has been reset and put back operative at 2006.362.00:22:56z (without harmonic compensation). At 2006.362.06.31.00z, ARTEMIS returned to his nominal operation with harmonic compensation activated.
- At 2006.362.13.04.25z, ARTEMIS triggered a spurious anomaly on the APC, which went into EMERGENCY mode. The APC has been reset and put back operative at 13:55:00z.
- At 2007.013.02.22.29z, ARTEMIS triggered a spurious anomaly on the APC, which went into READY NO-AUTO mode. The APC has been reset and put back operative at 02:46:58z.

## 6 CALIBRATION AND INSTRUMENT CHARACTERIZATION

### 6.1 Calibration

#### 6.1.1 Radiometric calibration

Cycle #54, potential radiometric calibrations are detailed in [subsection 4.3](#) page 13.

#### 6.1.2 Spectral calibration

Cycle #54, potential spectral calibrations (Wavelength Type 1 or 2) are detailed in [subsection 4.3](#) page 13.

#### 6.1.3 Geolocation

The accuracy specification for MERIS geolocation is 2000 m, with an operational goal of 150 m. The 290 m (nadir) bands 2, 5, 8 are used to estimate the absolute geolocation accuracy. This analysis shows significant improvements since launch, with one major upgrade, which occurred in 2003 DOY (Day of Year) 343. The update of the star tracker has been performed to reduce the systematic offset and improve orientation parameters. Global absolute geolocation error (North and South hemispheres) for the three consecutive periods can be summarized as follow:

- (I) Initially, after the launch, according to results related to the 2002 period, the geolocation accuracy is in the order of  $\pm 135$  metres along-track and  $\pm 207$  metres across-track. The RMS absolute geolocation error stays within the range of  $251.24 \pm 81$  m.
- (II) The 2003 period is characterised by a degradation of the absolute geolocation accuracy where error is around  $\pm 209$  metres along-track and  $\pm 295$  metres across-track. For this period, the RMS absolute geolocation error stays within the range of  $368.39 \pm 67$  m.
- (III) After the update, 2004 period, MERIS geolocation is achieving the goal of 300 m with accuracy of  $\pm 132$  m along-track and  $\pm 165$  m across-track. The RMS absolute geolocation error remains within the range of  $212 \pm 22$  m.

When correcting products from the systematic offset (centred results), for 2004 period the RMS absolute geolocation error stays within the range of  $166 \pm 18$  m. The amount of products located on northern hemisphere is much larger than the one from the Southern hemisphere. Comparison between the two sets of results is not trivial. For the 2004 period, this study demonstrated the temporal stability of the absolute geolocation. More results are now needed to confirm this trend. For more details, refer to the Gael Consultant (Fr) report available on the ESA website: <http://earth.esa.int/pcs/envisat/meris/reports/>

#### 6.1.4 VEU Temperature Analysis

During one of the operation modes of MERIS, Stabilization mode, a thermal regulation of VEU (Video Electronic Unit) unit is performed in order to stabilise its temperature to reach full performances and insure a safe transition towards Observation and Calibration modes. During observation, the VEU Temperature has to remain in the operational acceptance temperature

range  $-10^{\circ}/+50^{\circ}$  in order to meet the image quality requirements. The VEU temperature should be maximum  $\pm 10^{\circ}\text{C}$  different from the last radiometric calibration for optimum performance.

During cycle #54, one could observe two temperature increases, respectively the 24/12/2006 and the 29/12/2006 as seen in Figure 10. The biggest one occurred the 24/12/2006 as shown on the zoom in Figure 11. That day one could register a temperature of nearly  $36^{\circ}\text{C}$ . The fact is that on the 24<sup>th</sup> of December 2006, MERIS stayed in *Dir&Avg* mode<sup>1</sup> during the complete orbit and did not switch to *Heater* mode during eclipse. This anomaly was due to a controlling problem on ENVISAT spacecraft. The consequence was this little VEU temperature increase. According to the spacecraft manufacturer<sup>2</sup>, one cannot foresee any consequences on the MERIS instrument to dwell in acquisition mode if it only lasts the time of a single eclipse. As a matter of consequence one can deduce that the very little temperature increase that occurred the 29/12/2006 is also of no consequences.




Figure 10: VEU Temperature during cycle #54

---

<sup>1</sup>in acquisition

<sup>2</sup>Astrium




Figure 11: 24/12/2006 VEU temperature increase

### 6.1.5 Vicarious calibration results

For absolute calibration of MERIS by vicarious methods, METRIC2.0 tools are used to perform data extraction and spatial compression from MERIS Level1b products over specified sites following site type specific radiometric and geographic criteria. The child L1b products are ordered systematically on the basis of sites definition and mission analysis. Because the list of sites can be over-dimensioned and vary with season, it has a validity period of 3 months. Each L1b child product is submitted to METRIC with the correct version of auxiliary files MER\_INS\_AX and MER\_CP1\_AX used during its generation, and a dedicated resource file which stores all parameters necessary for data filtering (cloud and aerosol screening, distance from coast...). Metric generates one file for each selected site pertaining to the following categories, according to the potential use of the data in the calibration processing: Rayleigh, Glitter, Desert, Snow, and Buoy. Output files have HDF format. During cycle #24 new over-pass tables have been regenerated for all sites of interest updating the relative orbits inside the cycle. The site map is shown in [Figure 12](#).




Figure 12: METRIC calibration site map

During the cycle Metric has generated for specific sites the results presented [Table 10](#).

Latest results from these methods were presented at:

Second working meeting on MERIS and AATSR

Calibration and Geophysical Validation

MAVT-2006 ESRIN, Frascati, Italy

20-24 March 2006

Sites	#Products
DESERT	387
GLITTER	86
RAYLEIGH	0
SNOW	25
BUOY	87

Table 10: Specific sites results

Corresponding presentations can be found at:

[http://envisat.esa.int/workshops/mavt\\_2006/MAVT-2006-0303\\_CTinel.pdf](http://envisat.esa.int/workshops/mavt_2006/MAVT-2006-0303_CTinel.pdf)

[http://envisat.esa.int/workshops/mavt\\_2006/MAVT-2006-0304\\_CTinel.pdf](http://envisat.esa.int/workshops/mavt_2006/MAVT-2006-0304_CTinel.pdf)

Note: in the same Workshop, other results of vicarious calibration for MERIS, not based on METRIC extraction, were also presented.

## 6.2 Instrument Characterization

### 6.2.1 Instrument degradation

No new results to be shown for cycle #54. For the last updates, refer to Cyclic Report #45 that can be found on the above-mentioned MERIS website (see [subsubsection 6.1.3](#) page 16).

### 6.2.2 Diffuser ageing

No new results to be shown for cycle #54. For the last updates, refer to Cyclic Report #45 that can be found on the above-mentioned MERIS website.

### 6.2.3 Smile Effect

No new results to be shown for cycle #54. For the last updates, refer to Cyclic Report #23 that can be found on the above-mentioned MERIS website.

### 6.2.4 Spectral evolution from erbium measurements

No new results to be shown for cycle #54. Please refer to Cyclic Report #23 that can be found on the above-mentioned MERIS website.

## 7 DATA QUALITY CONTROL

### 7.1 MERIS products quality status

IPF version 5.02 did not have any impact on the MERIS products quality; an increase of the Level 2 processing time has been reported - it is linked to the increase of the number of pixels taken into account for the retrieval of aerosols over land, and the more detailed aerosols Look Up Table.



## 7.2 Anomalies and Software Problem Reporting (SPR)

Blank records have been identified in some MERIS products rejected by visual inspections using the AMALFI system. These black lines crossing the track are a nominal behavior of the processor, which replaces missing or corrupted Instrument Source Packets (ISPs) with blank data to preserve the geographical consistency of the scene.

## 8 FIRST 2003 MERIS ARCHIVE REPROCESSING

Information concerning the 1st reprocessing of the 2003 MERIS data archive done spring 2004 can be found on the MERIS website:

[http://earth.esa.int/pcs/envisat/meris/documentation/First\\_2003\\_MERIS\\_Reprocessing.pdf](http://earth.esa.int/pcs/envisat/meris/documentation/First_2003_MERIS_Reprocessing.pdf).

The document explains also how to get the reprocessed data.

## 9 SECOND 2005 MERIS ARCHIVE REPROCESSING

Following the recommendations of the Data Quality Working Group and the Science Advisory Group, improvements to MERIS processing resulted in version 7.4 of the off-line processor MEGS. It is currently being used for a complete reprocessing of the MERIS Reduced Resolution data archive. The corresponding time period extends from June 2002 to June 2005. 2003 and 2004 data will be made available through the MERCI (MERIS Catalogue and Inventory) service by the end of year 2005. For further information see:

<http://envisat.esa.int/services/catalogues.html>

## 10 MERIS PROCESSOR EVOLUTION

A detailed description of the MERIS IPF evolution since March 2002 until present, in terms of data format changes and algorithm modifications, can be found on the MERIS website:

[http://earth.esa.int/pcs/envisat/meris/documentation/MERIS\\_IPF\\_evolution.pdf](http://earth.esa.int/pcs/envisat/meris/documentation/MERIS_IPF_evolution.pdf)

## 11 VALIDATION ACTIVITIES AND RESULTS

The presentations given at the MAVT-2006 yield at ESRIN premises, Frascati, Italy, from 20 to 24 March 2006 are now available at the following address:

[http://envisat.esa.int/workshops/mavt\\_2006/](http://envisat.esa.int/workshops/mavt_2006/)

## 12 WATER VAPOUR AND BROWSE MAPS

Water Vapour data, retrieved from MER\_LRC\_2P products, have been used to generate global coverage maps for each day of the cycle. Maps are available on the ESA website:

<http://earth.esa.int/pcs/envisat/meris/maps/watervapour/>.

MERIS tracks for each day of the cycle have been plotted using Browse products. Maps are available on the ESA website: <http://earth.esa.int/pcs/envisat/meris/maps/browse/>

## 13 HOW TO GET MERIS DATA

Information concerning the different ways to access the MERIS data can be found on the MERIS website:

[http://earth.esa.int/pcs/envisat/meris/documentation/Access\\_to\\_MERIS\\_data.pdf](http://earth.esa.int/pcs/envisat/meris/documentation/Access_to_MERIS_data.pdf)

## 14 GENERAL INFORMATION

1. The European Space Agency organised a joint MERIS and (A)ATSR workshop, held at ESRIN, Frascati, Italy, on 26-30 September 2005. All information about the objectives of the workshop as well as the participants' presentations can be found on ESA's official page: [http://envisat.esa.int/workshops/meris\\_aatsr2005/](http://envisat.esa.int/workshops/meris_aatsr2005/)
2. The European Space Agency organised the second working meeting on MERIS and AATSR Calibration and Geophysical Validation (MAVT-2006) in ESRIN, Frascati, Italy, from 20 to 24 March 2006. All information about the objectives of the workshop as well as the participants' presentations can be found on ESA's official pages: <http://www.congrex.nl/06M07>