

IDEAS+-SER-OQC-REP-2071 Issue 1.0

Customer : ESRIN Document Ref : IDEAS+-SER-OQC-REP-2071

Contract No : 4000111304/14/I-AM Issue Date : 05 October 2015

WP No : 6110 Issue : 1.0



IDEAS+ Swarm Weekly Report 2015/39: 2015/09/21 - 2015/09/27

Abstract : This is the Instrument Data quality Evaluation and Analysis Service Plus

(IDEAS+) Swarm Weekly report on Swarm products quality, covering the period from

21 September to 27 September 2015.

Author : Approval :

Igino Coco, Jan Miedzik and Enkelejda Qamili on behalf of Swarm IDEAS+ Team Lidia Saavedra de Miguel IDEAS+ Science and Ops.

Coordinator

Distribution : ESA/ESRIN EOP-GMQ

ESA/ESRIN EOP-GM Swarm MM

IDEAS+ Leadership Team IDEAS+ subcontractors ESA/ESTEC Swarm PLSO ESA/ESOC Swarm FOS

Copyright © 2015 Telespazio VEGA UK Ltd

All rights reserved.

No part of this work may be disclosed to any third party translated reproduced copied or disseminated in any form or by any means except as defined in the contract or with the written permission of Telespazio Vega UK Ltd.

Telespazio VEGA UK Ltd

350 Capability Green, Luton, Bedfordshire, LU1 3LU, United Kingdom Tel: +44 (0) 1582 399 000 Fax: +44 (0) 1582 728 686 www.telespazio-vega.com



TABLE OF CONTENTS

TABLE OF CONTENTS	2
1. INTRODUCTION	5
1.1 Current Operational configuration of monitored data:	
1.2 Reference documents	
2. SUMMARY OF THE OBSERVATIONS	
2.1 General status of Swarm instruments and Level 1B products qual	
2.2 Plan for operational processor updates	8
2.3 Quality Working Group and Cal/Val Coordination	8
2.4 Summary of observations for 2015, Week 39 (21/09 - 27/09)	8
3. ROUTINE QUALITY CONTROL	10
3.1 Gaps analysis	
3.2 Orbit and Attitude Products	
3.2.1 Position Statistics	
3.2.2 Attitude observations	
3.3 Magnetic Products	
3.3.1 VFM-ASM anomaly	
3.3.2 ASM-VFM difference statistics	
3.3.4 VFM Instrument parameters: VFM temperatures (MAG_CA) .	
3.3.5 Magnetic time series visual inspection	
3.3.6 B _{NEC} vs Chaos4plus model residuals	24
4 ON-DEMAND ANALYSIS	28



This page intentionally left blank.



AMENDMENT POLICY

This document shall be amended by releasing a new edition of the document in its entirety. The Amendment Record Sheet below records the history and issue status of this document.

AMENDMENT RECORD SHEET

ISSUE	DATE	REASON
1.0	05 Oct 2015	First issue



1. INTRODUCTION

This document refers to the activities carried out in the framework of the Sensor Performance, Products and Algorithms (SPPA) Office [RD.1], and as such it reports on work related to:

- Algorithms and Processors Development, Maintenance and Evolution: these include all algorithm and software evolution and maintenance aspects for the different components, for both the Operational processors (OP) and Prototypes processors (PP) of L1 and L2 chains.
- Performance Assessment: these include all Quality Control activities (on-line and offline, systematic or on-demand), for the applicable product levels.
- System Calibration: these include the activities related to calibration, from sensor to system level. They also include aspects like cross calibration and handling of external calibration sources.
- Product validation: these include definition and maintenance of product validation plans.
- End-to-end Sensor Dataset Performance: these include activities related to the
 organisation and coordination of Quality Working Groups and all aspects of the
 Experimental platform. It also covers the product baseline, coordination and
 handling of external communities, and all aspects of ADF handling (both for the
 operational processors and for the prototypes).

This weekly report constitutes a work in progress throughout the mission life time, and new parts and complements will be added while the consolidation of knowledge on Swarm data and instruments will progress.

Section 2.1 always gives an overview of the general quality status of the mission instruments and products, while the main observations of the week are summarized in Section 2.2.

The document also includes information on data quality for the three Swarm spacecraft, inferred from automated HTML quality reports which are produced on daily basis for each product. Please contact the IDEAS+ Swarm team if interested in accessing the reports via web or FTP (all details about interfaces and folder structure available on [RD.2]). Such quality reports represent the core of the Routine Quality Control (Chapter 3). A description of the implemented quality checks is given in [RD.3], and references therein.

Basing on specific findings of the routine quality control, or on-demand from other entities (i.e. Swarm PDGS, FOS, Mission Management, Post-Launch Support Office, Expert Support Laboratories, Quality Working Groups, user community), anomalies can be triggered and preliminary characterisations and investigations of such anomalies are given in Chapter 4.The anomalies documented in the Weekly Reports are tracked in the following way:

- 1. If triggered by ESA Eohelp or within the Service: IDEAS+ action and ticketing system (http://requests-sppa.serco.it/RT3/index.html).
- 2. If triggered by IDEAS+ Swarm team or other entities:
 - 2a. If the observation/analysis leads to an anomaly to be addressed to the processor provider (GMV): SPR on EO ARTS (https://arts.eo.esa.int), SWL1L2DB project;
 - 2.b. If the observation/analysis does not lead to an anomaly or the investigation shall be escalated to other entities (PLSO/industry, ESL, PDGS): Action tracked on EO ARTS, **SW-IDEAS** project, then addressed to the proper tracking system if needed (e.g. JIRA for ESLs, SW-CP-AR project on EO ARTS for PDGS).

Information on Level 1B Swarm products can be found in [RD.4].



1.1 Current Operational configuration of monitored data:

- Processors Version: L1BOP 3.16, L2-Cat2 1.15
- L0 input products baseline: 02
- L1B baseline: MAGNET and PLASMA 04, ORBATT and ACCELE 03 (for definitions and description of the data baseline concept see https://earth.esa.int/web/guest/missions/esa-operational-eo-missions/swarm/data-access/product-baseline-definition)
- Level 2 Cat 2 baseline: EEF and TEC 01, FAC 02
- Input auxiliary files baseline: CCDB 0005 (20/07/2015), ADF 0101
- MPPF-CVQ v.2.14.01

1.2 Reference documents

The following is a list of documents with a direct bearing on the content of this report. Where referenced in the text, these are identified as RD.n, where 'n' is the number in the list below:

- [RD.1] Sensor Performance, Products and Algorithms (SPPA), PGSI-GSOP-EOPG-TN-05-0025. Version 2.3.
- [RD.2] Swarm PDGS External DMC Interface Control Document, SW-ID-DS-GS-0001, Issue 3.2.
- [RD.3] Swarm MPPF-CVQ Monitoring Baseline Document, ST-ESA-SWARM-MBD-0001, Issue 1.7.
- [RD.4] Swarm Level 1B Product Definition, SW-RS-DSC-SY-0007, Issue 5.13.
- [RD.5] Swarm IDEAS Configuration Management Plan, IDEAS-SER-MGT-PLN-1081 v0.14.
- [RD.6] Swarm Quality Control Project Plan, IDEAS-SER-MGT-PLN-1071
- [RD.7] SW_L1BOP_status_20141124_MoM
- [RD.8] Planned Updates for Level 1b, SW-PL-DTU-GS-008, Rev: 1dC.
- [RD.9] IDEAS+ Swarm Weekly Report: 25/08/2014 31/08/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20140825_20140831.pdf (ref. for SWL1L2DB-9)
- [RD.10] IDEAS+ Swarm Weekly Report: 29/09/2014 05/10/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20140929_20141005.pdf (ref. for SW-IDEAS-34)
- [RD.11] IDEAS+ Swarm Weekly Report: 06/10/2014 12/10/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20141006_20141012.pdf (ref. for SW-IDEAS-36)
- [RD.12] IDEAS+ Swarm Weekly Report: 20/10/2014 26/10/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20141020_20141026.pdf (ref. for SW-IDEAS-40, GPS sync loss)
- [RD.13] IDEAS+ Swarm Weekly Report: 15/09/2014 21/09/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20140915_20140921.pdf (ref. for SW-IDEAS-27)
- [RD.14] Swarm L1B 03.15 Validation Report, OSMV-OPMT-SRCO-RP-15-3385, Issue
- [RD.15] IDEAS+ Swarm Weekly Report: 23/03/2015 29/03/2015, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_201513_20150323_20150329.pdf.
- [RD.16] SWARM Weekly Operations Report #76, SW-RP-ESC-FS-6172



IDEAS+-SER-OQC-REP-2071 Issue 1.0

- [RD.17] Olsen, N., H. Luhr, C.C. Finlay, T.J. Sabaka, I. Michaelis, J. Rauberg and L. Tøffner-Clausen, The CHAOS-4 geomagnetic field model, Geophys. J. Int. 197, 815–827, 2014
- [RD.18] IDEAS+-SER-IPF-PLN-2272, Swarm Level 1B Operational Processor Verification Plan, IDEAS+-SER-IPF-PLN-2272_L1BOP_316_v1.5_final.pdf



2. SUMMARY OF THE OBSERVATIONS

2.1 General status of Swarm instruments and Level 1B products quality

Nothing to report.

2.2 Plan for operational processor updates

L1B: After having validated the patches, an update of VFM scale factors in the CCDB has been done on September 30. In this framework, a regeneration of the MAGNET and PLASMA data is therefore needed, starting from 18/07/2015. This will be done during CW 41 (5-11/10/2015). The full mission regeneration is still pending from the resolution of few ORBATT issues.

L2-Cat2: Data distributed by GFZ from beginning of mission for IBI, FAC and TEC will be opened to users during week 41 with baseline/counter 0201. At the same time the operational baseline/counter will be increased to 0202, and the gaps between the GFZ and operational production will be filled by an operational generation, with baseline/counter 0202 as well.

2.3 Quality Working Group and Cal/Val Coordination

Following the decisions of the 5th QWG in Paris, these activities will be carried on:

- The discussion group on the TII operational configuration will start its works on Wednesday 7th October, at 16 CET, through a WebEx teleconference involving the interested parties: ESA, University of Calgary, IRF, GFZ, Finnish Meteorological Institute, Institute of Space Science of Romania. The aim of this group is to develop a "mission planning" concept for the TII which should be sustainable in terms of automated operations and also flexible enough for the most interesting science purposes.
- The Langmuir Probe team at IRF, Uppsala, is preparing two datasets for expert users: 1) electron temperature data from single probe measurements separately and 2) 16 Hz raw faceplate currents data. The dataset will be distributed through the ESA FTP server (/Advanced folder) and will be ready, along with proper descriptions and documentation, by Mid November.
- The University of Calgary is preparing a new dataset for expert users: 16 Hz on board TII moments. The dataset will be distributed through the ESA FTP server (/Advanced folder) and will be ready, along with proper descriptions and documentation, by CW 41 (5-11/10/2015), or 42 at latest.
- The University of Calgary is going as well to update the TII preliminary dataset with new versions passed through more accurate calibration. The timeline for this is about one month.

2.4 Summary of observations for 2015, Week 39 (21/09 - 27/09)

During the monitored week the following events have been found and investigated:

Several few seconds gaps in MAGx_CA_1B products throughout the week.
 Some of them seem not to be associated to gaps in telemetry. Monitoring ongoing.





• Few MAG_HR gap on S/C Alpha, more details in 3.1.



3. ROUTINE QUALITY CONTROL

3.1 Gaps analysis

- MAG_CA_1B gaps throughout the week (SW-IDEAS-63), due at times to gaps in
 the HK telemetry required to compute magnetic stray fields; this is expected from
 the MAGNET algorithms, nevertheless only a portion of such gaps can be
 explained by a telemetry rejection, so analyses are still on-going for
 understanding the reason of the unexplained gaps.
- MAGx_HR_1B gaps on 21/09 and 25/09 for S/C Alpha, at the same times gaps in dB_AOCS and gaps in MAGx_CA_1B, t_csc, t_cdc and t_eu.

Table 1: Gaps in MAGA_HR on 21/09.

Start time gap	Stop time gap	Length gap (s)	Lack VFM temp
17:14:33	17:14:35	1.9999	Yes
18:02:33	18:02:35	1.9999	Yes
18:10:33	18:10:35	1.9999	Yes
20:42:33	20:42:35	1.9999	Yes
21:10:33	21:10:35	1.9999	Yes

Table 2: Gaps in MAGA_HR on 25/09.

Start time gap	Stop time gap	Length gap (s)	Lack VFM temp
03:53:53	03:53:55	1.99989	Yes

3.2 Orbit and Attitude Products

In Table 3 are listed events that have to be reported.

Table 3: List of events related to attitude and orbit products to be reported in the monitoring for 2015, Week 39: 21/09 - 27/09.

Observation ID	Description	Affected parameter	Sect. of Obs. Description	Sect. of Obs. Analysis
SW-IDEAS-63	OBS_ROUTINE: MAGx_CA_1B gaps	MAGx_CA_1B	3.1	3.1

The relevant parameters that have been monitored are:

- Position difference between calculated Medium Accuracy orbits (MODx_SC_1B) and on-board solution (GPSxNAV_0). Threshold values for such differences have not been assessed yet: we have just monitored the average values and maximum variations around the week, and reported in tables in the sections below, along with some example from the HTML daily reports. For the time being we evaluated an anomaly should be raised if one (or more) of the following conditions occurs:
 - The average difference on a given day exceeds the position accuracy requirement for the mission (1.5 m),



IDEAS+-SER-OQC-REP-2071 Issue 1.0

- The variability around the average is quite high: standard deviation threshold has been arbitrarily chosen to be twice the position accuracy requirement for the mission (2-sigma = 3 m).
- At least 4-5 spikes are observed on a given day, exceeding +/- 50 m.
- Visual inspection of Star Tracker characterisation flags (STRxATT_1B)
- Deviation of the quaternion norm from unity (deviation threshold = $\pm 10^{-9}$)
- Visual inspection of Euler Angles derived from quaternions.

3.2.1 Position Statistics

In Table 4, one can see the statistics of the differences between MOD and on-board solution positions for S/C A, B and C respectively. In the third column the maximum differences (maximum negative and maximum positive) are reported. The maximum standard deviation is in the fourth column. Maxima, minima and standard deviations usually refer to the Z component which is often the most disturbed; in case another component is most affected, it will be specified in parentheses.



Table 4: Swarm A, B and C, difference between MOD and on-board solution positions. If not specified maximum difference and maximum standard deviation refers to the Z axis.

	not specified ma	aximum dinere	nce and maxin	Turri Staridard deviation rei	ers to the Z
		Swarm A	, 21/09 - 27/09), Position difference	
Day	Average difference (m)	Maximum difference (m)		Maximum standard deviation (m)	Notes
21/09	0.16	-9.3	12.1	1.41	
22/09	0.1	-7 (Y)	8.2	1.27	
23/09	0.17	-8.4 (Y)	7.8	1.56	
24/09	0.17	-8.2	7.9	1.37	
25/09	0.12	-8	8	1.28	
26/09	0.04	-8.5	12.1 (Y)	1.38	
27/09	0.13	-6.3	6.8	1.26	
		Swarm B	3, 21/09 - 27/09	, Position difference	
Day	Average difference (m)	Maximum difference (m)		Maximum standard deviation (m)	Notes
21/09	0.2	-8.6 (Y)	9.5	1.33	
22/09	0.08	-8.1	13.8	1.34	
23/09	0.12	-9.2	7.1	1.48	
24/09	0.1	-6.1	6.5	1.23	
25/09	0.09	-5.9	7.4	1.22	
26/09	0.06	-10.7	7.1	1.53	
27/09	0.1	-8.2	6.2 (X)	1.33	
		Swarm C	;, 21/09 - 27/09	, Position difference	
Day	Average difference (m)	Maximum di	fference (m)	Maximum standard deviation (m)	Notes
21/09	0.14	-9.5	9.1	1.34	
22/09	0.09	-6	6.4	1.18	
23/09	0.17	-6.8 (Y)	9.2	1.54	
24/09	0.14	-8.2	7.4	1.34	
25/09	0.06	-6.5	9.4	1.25	
26/09	0.06	-7.9	8.6 (Y)	1.35	
27/09	0.12	-5.9	7.3	1.28	

3.2.1.1 Swarm A

Below some plot example follows of MOD-NAV differences, S/C A, taken at the beginning of the week (21/09, Figure 1) in the middle (24/09, Figure 2) and at the end (27/09, Figure 3). From top to bottom the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z axis respectively. The difference between both solutions is given in [m].






Figure 1: Difference MOD-GPSNAV, S/C A, 21/09. From top to bottom: X, Y and Z axis




Figure 2: Difference MOD-GPSNAV, S/C A, 24/09. From top to bottom: X, Y and Z axis






Figure 3: Difference MOD-GPSNAV, S/C A, 27/09. From top to bottom: X, Y and Z axis

3.2.1.2 Swarm B

Below some plot example follows of MOD-NAV differences, S/C B, taken at the beginning of the week (21/09, Figure 4), in the middle (24/09, Figure 5), and at end of the week (27/09, Figure 6). From top to bottom the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z axis respectively. The difference between both solutions is given in [m].






Figure 4: Difference MOD-GPSNAV, S/C B, 21/09. From top to bottom: X, Y and Z axis




Figure 5: Difference MOD-GPSNAV, S/C B, 24/09. From top to bottom: X, Y and Z axis






Figure 6: Difference MOD-GPSNAV, S/C B, 27/09. From top to bottom: X, Y and Z axis

3.2.1.3 Swarm C

Below some plot example of MOD-NAV differences, S/C C, taken at: the beginning of the week (21/09, Figure 7), in the middle (24/09, Figure 8) and at the end (27/09, Figure 9). From top to bottom the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z axis respectively. The difference between both solutions is given in [m].




Figure 7: Difference MOD-GPSNAV, S/C C, 21/09. From top to bottom: X, Y and Z axis






Figure 8: Difference MOD-GPSNAV, S/C C, 24/09. From top to bottom: X, Y and Z axis




Figure 9: Difference MOD-GPSNAV, S/C C, 27/09. From top to bottom: X, Y and Z axis

3.2.2 Attitude observations

3.2.2.1 Swarm A

Nothing to report.



3.2.2.2 Swarm B

Nothing to report.

3.2.2.3 Swarm C

Nothing to report.

3.3 Magnetic Products

For the magnetic products the weekly monitoring consists in:

- ASM instrument monitoring: quartz frequency and ASM temperature
- VFM instrument monitoring: temperatures
- Visual inspection of daily time series of magnetic field intensity F, BNEC and BVFM. Looking for gaps (or zero values in case of MAGx_LR_1B products), outof-threshold values (i.e. exceeding +/- 60000 nT), and other strange features.
- Monitoring of the ASM-VFM known anomaly: visual inspection of |BNEC| F and recording of daily maximum variations. If +/- 2 nT are exceed on a given day, an alert is raised.
- Comparison of magnetic data (B_NEC) with a model (Chaos4plus).

3.3.1 VFM-ASM anomaly

3.3.1.1 ASM-VFM difference statistics

In Table 5, one can see the statistics of the differences between magnetic field absolute value measured by ASM and by VFM. In the second and third column are reported the maximum differences, maximum negative and maximum positive respectively. The maximum standard deviation is in the fourth column.

The ASM-VFM difference is defined as follow: $dF = |B_{NEC}| - F_{ASM}$



Table 5 Swarm A and B, difference between magnetic field absolute value measured by ASM and by VFM.

	7 Civi and by v				
Swarm A, 21/09 - 27/09, ASM-VFM difference					
Day	Max (nT)	Min (nT)	Standard deviation (nT)	Mean (nT)	Notes
21/09	0.81	-47075.26	469.7	-5	MAGxHR gap
22/09	0.7	-0.86	0.18	0.02	
23/09	0.96	-1.22	0.18	0.01	
24/09	2.26	-1.43	0.21	-0.01	
25/09	3.36	-49978.11	170.03	-0.58	MAGxHR gap
26/09	0.96	-0.94	0.27	-0.03	
27/09	1.46	-1.09	0.32	0.09	
		Swarm B, 2	1/09 - 27/09, ASM-VFM	difference	
Day	Max (nT)	Min (nT)	Standard deviation (nT)	Mean (nT)	Notes
21/09	1.07	-0.37	0.16	0.14	
22/09	1.55	-0.58	0.17	0.12	
23/09	1.05	-1.18	0.16	0.13	
24/09	10.32	-0.88	0.17	0.14	
25/09	1.31	-0.53	0.18	0.13	
26/09	1.13	-1.28	0.2	0.13	
27/09	0.98	-0.49	0.19	0.06	




Figure 10: Daily mean and standard deviation values of ASM-VFM residuals (defined as $dF=|B_{VFM}|-F_{ASM}$) for S/C Alpha.






Figure 11: Daily mean and standard deviation values of ASM-VFM residuals (defined as $dF = |B_{VEM}| - F_{ASM}$) for S/C Bravo.

3.3.1.2 Swarm A

The daily peak-to-peak difference around the week stays within [-0.8, 0.8] nT with a few spikes not exceeding 1nT and two spikes of about 3-5nT on 24/09 and 25/09. Below two example plots of such differences for the days: 21/09 (Figure 12), and 27/09 (Figure 13).




Figure 12: ASM-VFM residuals for S/C A, 21/09.




Figure 13: ASM-VFM residuals for S/C A, 27/09.

3.3.1.3 Swarm B

The daily peak-to-peak difference around the week is, on average: [-0.6, 0.6] nT, with a few spikes not exceeding 2 nT and one spike of 10nT on 24/09. Below two example plots follows of such differences: 21/09 (Figure 14), and 27/09 (Figure 15).






Figure 14: ASM-VFM residuals for S/C B, 21/09.




Figure 15: ASM-VFM residuals for S/C B, 27/09.

3.3.1.4 Swarm C

No data because ASM is switched off.

3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC_0)

For S/C Alpha and Bravo the temperature and quartz frequency behaved as expected.

3.3.3 VFM Instrument parameters: VFM temperatures (MAG_CA)

The VFM instrument parameters important for monitoring the instrument health are the VFM sensor temperatures: T_CDC, T_CSC and T_EU.

For S/C Alpha and Bravo for reported period the temperatures behaved as expected.

3.3.4 Magnetic time series visual inspection

3.3.4.1 Swarm A

An example of representative magnetic field time series for S/C A (27/09) can be seen in Figure 16 below.






Figure 16: Time series of the geomagnetic field, for 27/09, S/C A. From top to bottom: magnetic field components in NEC reference frame, magnetic field components in the VFM reference frame, magnetic field intensity (F) from ASM, and location of gaps (if any).

3.3.4.2 Swarm B

An example of representative magnetic field time series for S/C B (27/09) can be seen in Figure 17 below.






Figure 17: Time series of the geomagnetic field for 27/09, S/C B. From top to bottom: magnetic field components in NEC reference frame, magnetic field components in the VFM reference frame, magnetic field intensity (F) from ASM, and location of gaps (if any).

3.3.4.3 Swarm C

An example of magnetic field time series for S/C C (27/09) can be seen in Figure 18.






Figure 18: Time series of the geomagnetic field for 27/09, S/C C. From top to bottom: magnetic field components in NEC reference frame, magnetic field components in the VFM reference frame, magnetic field intensity (F) from ASM (no data here because ASM it is off) and location of gaps.

3.3.5 B_{NEC} vs Chaos4plus model residuals

Figure 19, Figure 21 and Figure 23 show field residuals dB=B_{NEC} - B_{Chaos} (all versus colatitude in degrees), from top to bottom: 1) Br, 2) B θ and 3) B ϕ .

As a general feature one can see the field residuals to be steady and usually below 50 nT at low and middle latitudes, up to |55| - |60| degrees; then the residual increases at high latitudes because the Chaos model does not take into account the contribution from the external field ([RD.17]).

Figure 20, Figure 22 and Figure 24 shows, from top to bottom, the time series on 21/09 of: (1-2-3) residuals of B_{NEC} - B_{CHAOS} by components, related to Swarm Alpha, Bravo and Charlie respectively,

The component most affected by residual spikes and variations is B θ _NEC, i.e. the component which shows the variations of the field wrt to co-latitude. At high latitudes, the order of magnitude of the variability is about +/- 200 nT.



3.3.5.1 Swarm A




Figure 19: Swarm A day 21/09 B_NEC - B_Chaos vs colatitude.




Figure 20: Swarm A day 21/09: time series of B_NEC – B_Chaos residuals.



3.3.5.2 Swarm B




Figure 21: Swarm B day 21/09 B_NEC - B_Chaos difference vs colatitude.




Figure 22: Swarm B day 21/09 time series of B_NEC – B_Chaos residuals.



3.3.5.3 Swarm C




Figure 23: Swarm C day 21/09 B_NEC - B_Chaos difference vs colatitude.





Figure 24: Swarm C day 21/09 time series of B_NEC – B_Chaos residuals.



4. ON-DEMAND ANALYSIS

Nothing to report.





End of Document