

Customer	: ESRIN	Document Ref	:	IDEAS+-SER-OQC-REP-2071
Contract No	: 4000111304/14/I-A	M Issue Date	:	18 November 2015
WP No	: 6110	Issue	:	1.0



# IDEAS+ Swarm Weekly Report 2015/45: 2015/11/02 - 2015/11/08

Abstract : This is the Instrument Data quality Evaluation and Analysis Service Plus (IDEAS+) Swarm Weekly report on Swarm products quality, covering the period from 02 November to 08 November 2015.

Author	:	Approval :
	Igino Coco, Jan Miedzik and Enkelejda Qamili on behalf of Swarm IDEAS+ Team	Lidia Saavedra de Miguel IDEAS+ Science and Ops. Coordinator
Distribution	: ESA/ESRIN EOP-GMQ ESA/ESRIN EOP-GM Swarm M IDEAS+ Leadership Team IDEAS+ subcontractors ESA/ESTEC Swarm PLSO ESA/ESOC Swarm FOS	IM

#### Copyright © 2015 Telespazio VEGA UK Ltd

All rights reserved. No part of this work may be disclosed to any third party translated reproduced copied or disseminated in any form or by any means except as defined in the contract or with the written permission of Telespazio Vega UK Ltd.

Telespazio VEGA UK Ltd 350 Capability Green, Luton, Bedfordshire, LU1 3LU, United Kingdom Tel: +44 (0) 1582 399 000 Fax: +44 (0) 1582 728 686 www.telespazio-vega.com



# TABLE OF CONTENTS

2
5
6
6
8
8
8
8
9
10
10
10
10
14
14
15
15
0)17
17
18
20
24



IDEAS+-SER-OQC-REP-2071 Issue 1.0

This page intentionally left blank.

Telespazio VEGA UK Ltd © 2015



# AMENDMENT POLICY

This document shall be amended by releasing a new edition of the document in its entirety. The Amendment Record Sheet below records the history and issue status of this document.

#### AMENDMENT RECORD SHEET

ISSUE	DATE	REASON
1.0	18 Nov 2015	First issue



# 1. INTRODUCTION

This document refers to the activities carried out in the framework of the Sensor Performance, Products and Algorithms (SPPA) Office [RD.1], and as such, it reports on work related to:

- Algorithms and Processors Development, Maintenance and Evolution: these include all algorithm and software evolution and maintenance aspects for the different components, for both the Operational processors (OP) and Prototypes processors (PP) of L1 and L2 chains.
- Performance Assessment: these include all Quality Control activities (on-line and offline, systematic or on-demand), for the applicable product levels.
- System Calibration: these include the activities related to calibration, from sensor to system level. They also include aspects like cross calibration and handling of external calibration sources.
- Product validation: these include definition and maintenance of product validation plans.
- End-to-end Sensor Dataset Performance: these include activities related to the organisation and coordination of Quality Working Groups and all aspects of the Experimental platform. It also covers the product baseline, coordination and handling of external communities, and all aspects of ADF handling (both for the operational processors and for the prototypes).

This weekly report constitutes a work in progress throughout the mission lifetime, and new parts and complements will be added while the consolidation of knowledge on Swarm data and instruments will progress.

Section 2.1 always gives an overview of the general quality status of the mission instruments and products, while the main observations of the week are summarized in Section 2.1.

The document also includes information on data quality for the three Swarm spacecraft, inferred from automated HTML quality reports, which are produced on daily basis for each product. Please contact the IDEAS+ Swarm team if interested in accessing the reports via web or FTP (all details about interfaces and folder structure available on [RD.2]). Such quality reports represent the core of the Routine Quality Control (Chapter 3). A description of the implemented quality checks is given in [RD.3], and references therein.

Basing on specific findings of the routine quality control, or on-demand from other entities (i.e. Swarm PDGS, FOS, Mission Management, Post-Launch Support Office, Expert Support Laboratories, Quality Working Groups, and user community), anomalies can be triggered. Preliminary characterisations and investigations of such anomalies are given in Chapter 4.The anomalies documented in the Weekly Reports are tracked in the following way:

1. If triggered by ESA Eohelp or within the Service: IDEAS+ action and ticketing system (<u>http://requests-sppa.serco.it/RT3/index.html</u>).

2. If triggered by IDEAS+ Swarm team or other entities:

2a. If the observation/analysis leads to an anomaly to be addressed to the processor provider (GMV): SPR on EO ARTS (<u>https://arts.eo.esa.int</u> ), **SWL1L2DB** project;

2.b. If the observation/analysis does not lead to an anomaly or the investigation shall be escalated to other entities (PLSO/industry, ESL, PDGS): Action tracked on EO ARTS, **SW-IDEAS** project, then addressed to the proper tracking system if needed (e.g. JIRA for ESLs, SW-CP-AR project on EO ARTS for PDGS).

Information on Level 1B Swarm products can be found in [RD.4].



## **1.1** Current Operational configuration of monitored data:

- Processors Version: L1BOP 3.16.p2, L2-Cat2 1.15.p1.
- L0 input products baseline: 02
- L1B baseline: MAGNET and PLASMA 04, ORBATT and ACCELE 03 (for definitions and description of the data baseline concept see <u>https://earth.esa.int/web/guest/missions/esa-operational-eo-missions/swarm/dataaccess/product-baseline-definition</u>)
- Level 2 Cat 2 baseline: EEF 01, IBI, FAC and TEC 02
- Input auxiliary files baseline: CCDB 0005 (20/07/2015), ADF 0101
- MPPF-CVQ v.2.14.01

## **1.2** Reference documents

The following is a list of documents with a direct bearing on the content of this report. Where referenced in the text, these are identified as RD.n, where 'n' is the number in the list below:

- [RD.1] Sensor Performance, Products and Algorithms (SPPA), PGSI-GSOP-EOPG-TN-05-0025. Version 2.3.
- [RD.2] Swarm PDGS External DMC Interface Control Document, SW-ID-DS-GS-0001, Issue 3.2.
- [RD.3] Swarm MPPF-CVQ Monitoring Baseline Document, ST-ESA-SWARM-MBD-0001, Issue 1.7.
- [RD.4] Swarm Level 1B Product Definition, SW-RS-DSC-SY-0007, Issue 5.13.
- [RD.5] Swarm IDEAS Configuration Management Plan, IDEAS-SER-MGT-PLN-1081 v0.14.
- [RD.6] Swarm Quality Control Project Plan, IDEAS-SER-MGT-PLN-1071
- [RD.7] SW\_L1BOP\_status\_20141124\_MoM
- [RD.8] Planned Updates for Level 1b, SW-PL-DTU-GS-008, Rev: 1dC.
- [RD.9] IDEAS+ Swarm Weekly Report: 25/08/2014 31/08/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20140825\_20140831.pdf (ref. for SWL1L2DB-9)
- [RD.10] IDEAS+ Swarm Weekly Report: 29/09/2014 05/10/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20140929\_20141005.pdf (ref. for SW-IDEAS-34)
- [RD.11] IDEAS+ Swarm Weekly Report: 06/10/2014 12/10/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20141006\_20141012.pdf (ref. for SW-IDEAS-36)
- [RD.12] IDEAS+ Swarm Weekly Report: 20/10/2014 26/10/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20141020\_20141026.pdf (ref. for SW-IDEAS-40, GPS sync loss)
- [RD.13] IDEAS+ Swarm Weekly Report: 15/09/2014 21/09/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20140915\_20140921.pdf (ref. for SW-IDEAS-27)
- [RD.14] Swarm L1B 03.15 Validation Report, OSMV-OPMT-SRCO-RP-15-3385, Issue 1.3.
- [RD.15] IDEAS+ Swarm Weekly Report: 23/03/2015 29/03/2015, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_201513\_20150323\_20150329.pdf.
- [RD.16] SWARM Weekly Operations Report #76, SW-RP-ESC-FS-6172



- [RD.17] Olsen, N., H. Luhr, C.C. Finlay, T.J. Sabaka, I. Michaelis, J. Rauberg and L. Tøffner-Clausen, The CHAOS-4 geomagnetic field model, Geophys. J. Int. 197, 815–827, 2014
- [RD.18] IDEAS+-SER-IPF-PLN-2272, Swarm Level 1B Operational Processor Verification Plan, IDEAS+-SER-IPF-PLN-2272\_L1BOP\_316\_v1.5\_final.pdf



# 2. SUMMARY OF THE OBSERVATIONS

## 2.1 General status of Swarm instruments and Level 1B products quality

Nothing to report

## 2.2 Plan for operational processor updates

After the deployment in operations of the L2Cat2 1.15 patch 3 affecting the FAC processor, a partial regeneration have been performed. I.e. L2-Cat2 FAC combined regeneration from 16/7/15 to 3/11/2015 and L2-Cat2 FAC Single regeneration from 16/7/15 to 26/8/2015. The still missing data (02/06-15/07) will be produced by GFZ.

The magnetic data with product baseline 04 we are currently producing in operational are corrected from the scalar residuals that are thought to be predominantly Sun-driven. But one part of the correction is included in VFM calibration parameters, i.e. a scaling of the  $\beta$  Sun incident angle  $s_{\beta}$  in included in the sensitivity vector changing with time  $s_t$ . A change will be introduced in the L1BOP Magnet processor in order to have these two parameters separately. A new patch will be delivered soon by GMV containing this change.

There is a need to update the Euler angles by end of November 2015. As soon as the updated information will be provided, a regeneration of the MAG\_LR\_1B data will be performed using the MAGREP processor. The impact on L2-CAT2 data from Euler angles update has to be investigated once Level-1B regenerated data will be available. The regeneration will be performed only from 16/07/2015, date when the L1BOP v03.16 has been put in operation. The MAG\_LR\_1B data from beginning of the mission till 15/07/2015 will be provided by DTU.

A discrepancy has been found while comparing the Swarm L1B Rinex products (GPS\_RO\_1B) with the same Rinex obtained independently by DLR. This issue has been investigated by GMV which has found the cause of the differences to be due to the pseudo-ranges that were noisier due to the scaling of the Instrument Measurement Time (IMT). This issue was fixed in the ORBATT processor and soon a patch with be delivered.

## 2.3 Quality Working Group and Cal/Val Coordination

Following the decisions of the 5<sup>th</sup> QWG in Paris, these activities will be carried on in order to better understand the origin of the ASM-VFM residuals:

- Investigation on boom alignment error budget (from industry) that could help in developing a thermos-elastic correction approach.
- Investigation on plasma-induced stray field in order to focus on any possible current flow near the VFM (and not the ASM), because the intensity of such a current may be stronger near the VFM due to the accumulation of charged plasma particles near the STR baffles.

In the meantime, DTU will deliver an ASM-VFM residual dataset to be distributed to all the Swarm users through the ESA FTP server (/Advanced folder).



# 2.4 Summary of observations for 2015, Week 45 (02/11 - 08/11)

During the monitored week the following events have been found and investigated:

- Several few seconds gaps in MAGx\_CA\_1B products throughout the week. Some of them seem not to be associated to gaps in telemetry. Monitoring ongoing.
- One MAG\_LR gap of 2 seconds duration on S/C A, more details in 3.1



# 3. ROUTINE QUALITY CONTROL

## 3.1 Gaps analysis

- One MAG\_LR gap on SC A on 07/11 at 04:49:55, 5 seconds long. The gap is a result of L1B processing - sample rejection.
- Several few seconds gaps in MAGx\_CA\_1B products throughout the week. Some of them seem not to be associated to gaps in telemetry. Monitoring ongoing.

# 3.2 Orbit and Attitude Products

In Table 1 are listed events that have to be reported.

 Table 1: List of events related to attitude and orbit products to be reported in the monitoring for 2015, Week 45: 02/11 - 08/11.

Observation ID	Description	Affected parameter	Sect. of Obs. Description	Sect. of Obs. Analysis
SW-IDEAS-63	OBS_ROUTINE: MAGx_CA_1B gaps	MAGx_CA_1B	3.1	3.1

The relevant parameters that have been monitored are:

- Position difference between calculated Medium Accuracy orbits (MODx\_SC\_1B) and on-board solution (GPSxNAV\_0). Threshold values for such differences have not been assessed yet: we have just monitored the average values and maximum variations around the week. They are reported in tables in the sections below. In addition, some example plots are given from the HTML daily reports. For the time being we evaluated an anomaly should be raised if one (or more) of the following conditions occurs:
  - The **average difference** on a given day exceeds the position accuracy requirement for the mission (1.5 m),
  - The variability around the average is quite high: **standard deviation** threshold has been arbitrarily chosen to be twice the position accuracy requirement for the mission (2-sigma = 3 m).
  - At least 4-5 spikes are observed on a given day, exceeding +/- 50 m.
- Visual inspection of Star Tracker characterisation flags (STRxATT\_1B)
- Deviation of the quaternion norm from unity (deviation threshold =  $\pm -10^{-9}$ )
- Visual inspection of Euler Angles derived from quaternions.

## 3.2.1 Position Statistics

In Table 2, one can see the statistics of the differences between MOD and on-board solution positions for S/C A, B and C respectively. In the third column the maximum differences (maximum negative and maximum positive) are reported. The maximum standard deviation is in the fourth column. Maxima, minima and standard deviations usually refer to the Z component that is often the most disturbed; in case another component is most affected, it will be specified in parentheses. Figure 1 shows a cumulative trend of the maximum daily standard deviation for the past 30 days of operations of the MOD-NAV difference, while Figure 2 shows the daily maximum




difference, in absolute value, of the MOD-NAV difference, always for the past 30 days of operations.


**Table 2:** Swarm A, B and C, difference between MOD and on-board solution positions. If not specified maximum difference and maximum standard deviation refers to the Z-axis.

Swarm A, 02/11 - 08/11, Position difference						
Day	Average difference (m)	Maximum difference (m)		Maximum standard deviation (m)	Notes	
02/11	0.06	-9 6.2		1.3		
03/11	0.11	-9.4	8.5 (Y)	1.31		
04/11	0.11	-6.6	7.8 (Y)	1.34		
05/11	0.08	-7.2	7.6	1.25		
06/11	0.02	-6.5	6.8	1.16		
07/11	0.14	-7.2	7.3	1.27		
08/11	0.06	-6.9	7.7	1.28		
		Swarm B	8, 02/11 - 08/11	, Position difference		
Day	Average difference (m)	Maximum di	ifference (m)	Maximum standard deviation (m)	Notes	
02/11	0.13	-5.4	6.6	1.32		
03/11	0.09	-7.8	6 (X)	1.28		
04/11	0.19	-9	6.8	1.4		
05/11	0.13	-5.6	8.7	1.22		
06/11	0.05	-6.1	9.2 (Y)	1.2		
07/11	0.05	-5.9	5.6 (Y)	1.14		
08/11	0.06 -9 (Y) 9.3		1.4			
		Swarm C	<b>;, 02/11 - 08/1</b> 1	, Position difference		
Day	Average difference (m)	Maximum difference (m)		Maximum standard deviation (m)	Notes	
02/11	0.08	-6.9	5.9	1.28		
03/11	0.11	-10.5	7.8	1.24		
04/11	0.11	-7.7	5.9	1.37		
05/11	0.06	-6.4	7.8	1.23		
06/11	0.05	-11.5	11.2 (Y)	1.21		
07/11	0.07	-10.7 (Y)	8 (Y)	1.28		
08/11	0.06	-6.5	7	1.21		





**Figure 1:** Plot of the standard deviation of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.



**Figure 2:** Plot of the maximum difference of the absolute value of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.

#### 3.2.1.1 Swarm A

Below is presented plot example of MOD-NAV differences of S/C A at the beginning of the week (02/11, Figure 3). From top to bottom, the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].






Figure 3: Difference MOD-GPSNAV, S/C A, 02/11. From top to bottom: X, Y and Z-axis

#### 3.2.1.2 Swarm B

Below is presented plot example of MOD-NAV differences of S/C B at the beginning of the week (02/11, Figure 4). From top to bottom, the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].




Figure 4: Difference MOD-GPSNAV, S/C B, 02/11. From top to bottom: X, Y and Z-axis



## 3.2.1.3 Swarm C

Below is presented plot example of MOD-NAV differences of S/C C at the beginning of the week (02/11, Figure 5). From top to bottom, the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].




Figure 5: Difference MOD-GPSNAV, S/C C, 02/11. From top to bottom: X, Y and Z-axis

## 3.2.2 Attitude observations

#### 3.2.2.1 Swarm A

Nothing to report.

#### 3.2.2.2 Swarm B

Nothing to report.

#### 3.2.2.3 Swarm C

Nothing to report.

## 3.3 Magnetic Products

For the magnetic products, the weekly monitoring consists in:

- ASM instrument monitoring: quartz frequency and ASM temperature
- VFM instrument monitoring: temperatures
- Visual inspection of daily time series of magnetic field intensity F, B<sub>NEC</sub> and B<sub>VFM</sub>. Looking for gaps (or zero values in case of MAGx\_LR\_1B products), out-ofthreshold values (i.e. exceeding +/- 60000 nT), and other strange features.



- Monitoring of the ASM-VFM known anomaly: visual inspection of  $|B_{VFM}|$  F and recording of daily maximum variations. If +/- 2 nT are exceed on a given day, an alert is raised.
- Comparison of magnetic data (B<sub>NEC</sub>) with a model (Chaos4plus)

## 3.3.1 VFM-ASM anomaly

## 3.3.2 ASM-VFM difference statistics

In Table 3, one can see the statistics of the differences between magnetic field absolute value measured by ASM and by VFM. In the second and third column are reported the maximum differences, maximum negative and maximum positive respectively. The maximum standard deviation is in the fourth column.

The ASM-VFM difference is defined as follow:

 $dF = |B_{VFM}| - F_{ASM}$ 

Figure 6 and Figure 7 shows the daily mean (circles) and standard deviation (crosses) of dF from the deployment of the new L1B processor on (18/07) for Swarm A and Swarm B respectively. The only remarkable feature is a peak on the residual standard deviation for Swarm B, which reaches a 400 pT maximum around the 20 of August. The experts at DTU claim this is not a cause of concern.

**Table 3** Swarm A and B, difference between magnetic field absolute value measured by ASM and by VFM.

Swarm A, 02/11 - 08/11, ASM-VFM difference					
Day	Max (nT)	Min (nT)	Standard deviation (nT)	Mean (nT)	Notes
02/11	1.34	-0.87	0.23	0.04	
03/11	2.41	-3.25	0.22	0.03	
04/11	6.92	-1.27	0.21	0.02	
05/11	0.93	-5.16	0.27	0.01	
06/11	1.06	-0.96	0.29	0.02	
07/11	46431.54	-5.51	273.56	1.63	Gap in MAG_LR see 3.1
08/11	1.6	-1.24	0.31	0.02	
		Swarm B, 0	2/11 - 08/11, ASM-VFM	difference	•
Day	Max (nT)	Min (nT)	Standard deviation (nT)	Mean (nT)	Notes
02/11	1.22	-1.16	0.17	-0.05	
03/11	8.14	-3.93	0.2	-0.05	
04/11	2.36	-7.48	0.23	-0.02	
05/11	1.6	-0.73	0.22	-0.03	
06/11	5.39	-6.07	0.22	-0.04	
07/11	5.38	-3.65	0.23	-0.03	
08/11	1.97	-2.09	0.22	-0.04	






Figure 6 Daily mean and standard deviation values of ASM-VFM residuals (defined as  $dF=|B_{VFM}|-F_{ASM}$ ) for S/C A.




Figure 7 Daily mean and standard deviation values of ASM-VFM residuals (defined as  $dF=|B_{VFM}|-F_{ASM}$ ) for S/C B.

#### 3.3.2.1 Swarm A

The daily peak-to-peak difference around the week stays within [-0.75, 0.5] nT with a few spikes not exceeding 6nT, most spikes on 07/11. Below two example plots of such differences for the days: 02/11 (Figure 8), and 08/11 (Figure 9).




Figure 8: ASM-VFM residuals for S/C A, 02/11.






Figure 9: ASM-VFM residuals for S/C A, 08/11.

#### 3.3.2.2 Swarm B

The daily peak-to-peak difference around the week is, on average: [-0.5, 0.64] nT, with a few spikes not exceeding 8 nT, most spikes on 03/11 and 07/11. Below two example plots follows of such differences: 02/11 (Figure 10), and 08/11 (Figure 11).




Figure 10: ASM-VFM residuals for S/C B, 02/11.




Figure 11: ASM-VFM residuals for S/C B, 08/11.

#### 3.3.2.3 Swarm C

No data because ASM is switched off.

# 3.3.3 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC\_0)

For S/C A and B, the temperature and quartz frequency behaved as expected.

## 3.3.4 VFM Instrument parameters: VFM temperatures (MAG\_CA)

The VFM instrument parameters important for monitoring the instrument health are the VFM sensor temperatures: T\_CDC, T\_CSC and T\_EU.


For S/C A, B and C, or reported period, the temperatures behaved as expected.



## 3.3.5 Magnetic time series visual inspection

#### 3.3.5.1 Swarm A

An example of representative magnetic field time series for S/C A (08/11) can be seen in Figure 12 below.




**Figure 12:** Time series of the geomagnetic field, for 08/11, S/C A. From top to bottom: magnetic field components in NEC reference frame, magnetic field components in the VFM reference frame, magnetic field intensity (F) from ASM, and location of gaps (if any).

#### 3.3.5.2 Swarm B

An example of representative magnetic field time series for S/C B (08/11) can be seen in Figure 13 below.





**Figure 13:** Time series of the geomagnetic field for 08/11, S/C B. From top to bottom: magnetic field components in NEC reference frame, magnetic field components in the VFM reference frame, magnetic field intensity (F) from ASM, and location of gaps (if any).

#### 3.3.5.3 Swarm C

An example of magnetic field time series for S/C C (08/11) can be seen in Figure 14.




Figure 14: Time series of the geomagnetic field for 08/11, S/C C. From top to bottom: magnetic field components in NEC reference frame, magnetic field components in the



VFM reference frame, magnetic field intensity (F) from ASM (no data here because ASM it is off) and location of gaps.

## 3.3.6 B<sub>NEC</sub> vs Chaos4plus model residuals

Figure 15, Figure 17 and Figure 19 show field residuals dB=B<sub>NEC</sub> - B<sub>Chaos</sub> (all versus colatitude in degrees), from top to bottom: 1) Br, 2) B $\theta$  and 3) B $\phi$ .

As a general feature one can see the field residuals to be steady and usually below 50 nT at low and middle latitudes, up to |55| - |60| degrees; then the residual increases at high latitudes because the Chaos model does not take into account the contribution from the external field ([RD.17]).

Figure 16, Figure 18 and Figure 20 shows, from top to bottom, the time series on 02/11 of: (1-2-3) residuals of  $B_{NEC}$ - $B_{CHAOS}$  by components, related to S/C A, B and C respectively.

The component most affected by residual spikes and variations is B $\theta$ \_NEC, i.e. the component that shows the variations of the field wrt to co-latitude. At high latitudes, the order of magnitude of the variability is about +/- 200 nT.

## 3.3.6.1 Swarm A




Figure 15: Swarm A day 02/11 B\_NEC - B\_Chaos vs colatitude.






Figure 16: Swarm A day 02/11: time series of B\_NEC – B\_Chaos residuals.

## 3.3.6.2 Swarm B




Figure 17: Swarm B day 02/11 B\_NEC - B\_Chaos difference vs colatitude.






Figure 18: Swarm B day 02/11 time series of B\_NEC – B\_Chaos residuals.

## 3.3.6.3 Swarm C




Figure 19: Swarm C day 02/11 B\_NEC - B\_Chaos difference vs colatitude.






Figure 20: Swarm C day 02/11 time series of B\_NEC – B\_Chaos residuals.



IDEAS+-SER-OQC-REP-2071 Issue 1.0

# 4. ON-DEMAND ANALYSIS

Nothing to report.



IDEAS+-SER-OQC-REP-2071 Issue 1.0

End of Document