



Customer	: ESRIN	Document Ref	: IDEAS+-SER-OQC-REP-2071
Contract No	: 4000111304/14/I-AM	Issue Date	: 19 October 2016
WP No	: 6110	Issue	: 1.0



## IDEAS+ Swarm Weekly Report 2016/22: 2016/05/30 - 2016/06/05

**Abstract** : This is the **Instrument Data Quality Evaluation and Analysis Service Plus (IDEAS+)** Swarm Weekly report on Swarm products quality, covering the period from 30 May to 05 June 2016.

**Author** : \_\_\_\_\_ **Approval** : \_\_\_\_\_  
Igino Coco, Jan Miedzik and Marta De Laurentis  
Enkelejda Qamili on behalf of IDEAS+ Science and Ops.  
Swarm IDEAS+ Team Coordinator

**Distribution** : ESA/ESRIN EOP-GMQ ESA/ESTEC Swarm PLSO  
ESA/ESRIN EOP-GM Swarm MM ESA/ESOC Swarm FOS  
IDEAS+ Leadership Team  
IDEAS+ subcontractors

**Copyright © 2015 Telespazio VEGA UK Ltd**

*All rights reserved.*

*No part of this work may be disclosed to any third party translated reproduced copied or disseminated in any form or by any means except as defined in the contract or with the written permission of Telespazio Vega UK Ltd.*

**Telespazio VEGA UK Ltd**  
**350 Capability Green, Luton, Bedfordshire, LU1 3LU, United Kingdom**  
**Tel: +44 (0) 1582 399 000 Fax: +44 (0) 1582 728 686**  
**www.telespazio-vega.com**



## TABLE OF CONTENTS

<b>TABLE OF CONTENTS .....</b>	<b>2</b>
<b>1. INTRODUCTION .....</b>	<b>5</b>
1.1 Current Operational configuration of monitored data: .....	5
1.2 Reference documents .....	6
<b>2. SUMMARY OF THE OBSERVATIONS .....</b>	<b>7</b>
2.1 General status of Swarm instruments and Level 1B products quality .....	7
2.2 Plan for operational processor updates .....	7
2.3 Quality Working Group and Cal/Val Coordination .....	7
2.4 Summary of observations for 2016, Week 22 (30/05 - 05/06) .....	8
<b>3. ROUTINE QUALITY CONTROL .....</b>	<b>9</b>
3.1 Gaps analysis .....	9
3.2 Orbit and Attitude Products .....	9
3.2.1 Position Statistics .....	9
3.2.2 Attitude observations .....	14
3.3 Magnetic Products .....	14
3.3.1 VFM-ASM anomaly .....	15
3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC_0) .....	19
3.3.3 VFM Instrument parameters: VFM temperatures (MAG_CA) .....	19
3.3.4 Magnetic time series visual inspection .....	21
3.3.5 S/C A and C magnetic correlation .....	26
3.3.6 S/C A and C magnetic difference .....	26
3.3.7 $B_{NEC}$ vs Chaos5 model residuals .....	26
3.3.8 Second derivative of $B_{NEC}$ and $B_{VFM}$ .....	30
<b>4. ON-DEMAND ANALYSIS .....</b>	<b>33</b>



*This page intentionally left blank.*



## **AMENDMENT POLICY**

This document shall be amended by releasing a new edition of the document in its entirety. The Amendment Record Sheet below records the history and issue status of this document.

### **AMENDMENT RECORD SHEET**

<b>ISSUE</b>	<b>DATE</b>	<b>REASON</b>
1.0	19 Oct 2016	First issue



## 1. INTRODUCTION

This document refers to the activities carried out in the framework of the Sensor Performance, Products and Algorithms (SPPA) Office [RD.1], and as such, it reports on work related to:

- Algorithms and Processors Development, Maintenance and Evolution: these include all algorithm and software evolution and maintenance aspects for the different components, for both the Operational processors (OP) and Prototypes processors (PP) of L1 and L2 chains.
- Performance Assessment: these include all Quality Control activities (on-line and offline, systematic or on-demand), for the applicable product levels.
- System Calibration: these include the activities related to calibration, from sensor to system level. They also include aspects like cross calibration and handling of external calibration sources.
- Product validation: these include definition and maintenance of product validation plans.
- End-to-end Sensor Dataset Performance: these include activities related to the organisation and coordination of Quality Working Groups and all aspects of the Experimental platform. It also covers the product baseline, coordination and handling of external communities, and all aspects of ADF handling (both for the operational processors and for the prototypes).

This weekly report constitutes a work in progress throughout the mission lifetime, and new parts and complements will be added while the consolidation of knowledge on Swarm data and instruments will progress.

Section 2.1 always gives an overview of the general quality status of the mission instruments and products, while the main observations of the week are summarized in Section 2.1.

The document also includes information on data quality for the three Swarm spacecraft, inferred from automated HTML quality reports, which are produced on daily basis for each product. Please contact the IDEAS+ Swarm team if interested in accessing the reports via web or FTP (all details about interfaces and folder structure available on [RD.2]). Such quality reports represent the core of the Routine Quality Control (Chapter 3). A description of the implemented quality checks is given in [RD.3], and references therein.

Basing on specific findings of the routine quality control, or on-demand from other entities (i.e. Swarm PDGS, FOS, Mission Management, Post-Launch Support Office, Expert Support Laboratories, Quality Working Groups, and user community), anomalies can be triggered. Preliminary characterisations and investigations of such anomalies are given in Chapter 4. The anomalies documented in the Weekly Reports are tracked in the following way:

1. If triggered by ESA Eohelp or within the Service: IDEAS+ action and ticketing system (<http://requests-sppa.serco.it/RT3/index.html> ).
2. If triggered by IDEAS+ Swarm team or other entities:
  - 2a. if the observation/analysis leads to an anomaly to be addressed to the processor provider (GMV): SPR on EO ARTS (<https://arts.eo.esa.int> ), **SWL1L2DB** project;
  - 2b. if the observation/analysis does not lead to an anomaly or the investigation shall be escalated to other entities (PLSO/industry, ESL, and PDGS): Action tracked on EO ARTS, **SW-IDEAS** project, then addressed to the proper tracking system if needed (e.g. JIRA for ESLs, SW-CP-AR project on EO ARTS for PDGS).

Information on Level 1B Swarm products can be found in [RD.4].

### 1.1 Current Operational configuration of monitored data:

- Processors Version: L1BOP 3.17, L2-Cat2 1.15.p5.
- L0 input products baseline: 02



- L1B baseline: MAGNET and PLASMA 04, ORBATT and ACCELE 03 (for definitions and description of the data baseline concept see <https://earth.esa.int/web/guest/missions/esa-operational-eo-missions/swarm/data-access/product-baseline-definition>)
- Level 2 – Cat 2 baseline: EEF 01, IBI, FAC and TEC 02
- Input auxiliary files baseline: S/C A - CCDB 0011 (17/05/2016), S/C B – CCDB 0011 (14/04/2016), S/C C – CCDB 0011 (17/05/2016), ADF 0101
- MPPF-CVQ v.2.14.01

## 1.2 Reference documents

The following is a list of documents with a direct bearing on the content of this report. Where referenced in the text, these are identified as RD.n, where 'n' is the number in the list below:

- [RD.1] Sensor Performance, Products and Algorithms (SPPA), PGSI-GSOP-EOPG-TN-05-0025. Version 2.3.
- [RD.2] Swarm PDGS External DMC Interface Control Document, SW-ID-DS-GS-0001, Issue 3.2.
- [RD.3] Swarm MPPF-CVQ Monitoring Baseline Document, ST-ESA-SWARM-MBD-0001, Issue 1.7.
- [RD.4] Swarm Level 1B Product Definition, SW-RS-DSC-SY-0007, Issue 5.13.
- [RD.5] Swarm IDEAS Configuration Management Plan, IDEAS-SER-MGT-PLN-1081 v0.14.
- [RD.6] Swarm Quality Control Project Plan, IDEAS-SER-MGT-PLN-1071
- [RD.7] SW\_L1BOP\_status\_20141124\_MoM
- [RD.8] Planned Updates for Level 1b, SW-PL-DTU-GS-008, Rev: 1dC.
- [RD.9] IDEAS+ Swarm Weekly Report: 25/08/2014 – 31/08/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20140825\_20140831.pdf (ref. for SWL1L2DB-9)
- [RD.10] IDEAS+ Swarm Weekly Report: 29/09/2014 – 05/10/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20140929\_20141005.pdf (ref. for SW-IDEAS-34)
- [RD.11] IDEAS+ Swarm Weekly Report: 06/10/2014 – 12/10/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20141006\_20141012.pdf (ref. for SW-IDEAS-36)
- [RD.12] IDEAS+ Swarm Weekly Report: 20/10/2014 – 26/10/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20141020\_20141026.pdf (ref. for SW-IDEAS-40, GPS sync loss)
- [RD.13] IDEAS+ Swarm Weekly Report: 15/09/2014 – 21/09/2014, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_20140915\_20140921.pdf (ref. for SW-IDEAS-27)
- [RD.14] Swarm L1B 03.15 Validation Report, OSMV-OPMT-SRCO-RP-15-3385, Issue 1.3.
- [RD.15] IDEAS+ Swarm Weekly Report: 23/03/2015 – 29/03/2015, IDEAS+-SER-OQC-REP-2071\_SPPA\_SwarmWeeklyReport\_201513\_20150323\_20150329.pdf.
- [RD.16] SWARM Weekly Operations Report #76, SW-RP-ESC-FS-6172
- [RD.17] Olsen, N., H. Luhr, C.C. Finlay, T.J. Sabaka, I. Michaelis, J. Rauberg and L. Tøffner-Clausen, The CHAOS-4 geomagnetic field model, Geophys. J. Int. 197, 815–827, 2014
- [RD.18] IDEAS+-SER-IPF-PLN-2272, Swarm Level 1B Operational Processor Verification Plan, IDEAS+-SER-IPF-PLN-2272\_L1BOP\_316\_v1.5\_final.pdf



## 2. SUMMARY OF THE OBSERVATIONS

### 2.1 General status of Swarm instruments and Level 1B products quality

An in-orbit residual test campaign consisting in the activation of heaters near the VFM sensor was performed from 30/05 to 03/06 on Swarm Alpha. Actually, the disturbance field is believed to be the consequence of local heating by the Sun near the VFM sensor, and a likely mechanism is thermo-electric current generation. Thus the aim of the tests was to create temperature gradients near the VFM sensor.

Only in the test using the heater on the Optical Bench Radiator balcony the VFM sensor temperature was impacted by a few degrees (but not sufficient to explain the ASM-VFM scalar residuals). The other test, using the Optical Bench Interface heater, resulted in VFM sensor temperature variations of less than a couple of degrees, deemed insufficient to sign any meaningful thermal gradient near the VFM sensor.

### 2.2 Plan for operational processor updates

**L1BOP:** In the next delivery of the L1BOP after the PDGS evolution activities (not earlier than Sept./Oct. 2016) the following changes will be implemented into the MAGNET processor:

- Change the data resampling used for the interpolation of the ASM measurements to UTC second because the present used in L1B data is removing a significant part of the high frequency content of the ASM data.
- Modification of the F\_error computation
- More robust handling of the leap second occurrence, and modification in timestamp sorting, in order to make the processor complete the run even in case of VFM on-board anomalies, that could imply a packet frequency greater than 1 Hz.

For what concern the ORBATT processor, two changes to be implemented are under discussion.

- A STR Inter-Boresight Angles correction model
- An increase of the frequency of the STR L0 product from 1 Hz to 2 Hz

**L2 Cat-2:** Any other delivery of the L2 Cat-2 OP is postponed after the PDGS evolution activity (Sept./Oct. 2016).

### 2.3 Quality Working Group and Cal/Val Coordination

The former "Task Force" deputed to study the ASM-VFM residuals, has been reshaped in a Magnetic Measurements Expert Group (MMEG). The first MMEG meeting was held in Warsaw (GMV premises) on **10-11/03/2016**. During this meeting the following decisions have been taken:

ASM-VFM Scalar Residual:

- VFM calibration issue:



- Finalize the covariance analysis between parameters in alpha/beta space
- Further analysis directly on non-orthogonalities in-flight and on ground test data and not the computed models.
- Scale factor  $s_{exp}$  shall be further investigated.
- Consolidate the analysis on dependency on  $s$  and  $T_{EU}$ .
- Further investigation on the correlation between ASM-VFM residuals and  $T_{EU}$ .
- Continue the investigation on ASM-VFM residuals using MAGx\_HR\_1B data.
- Consolidate test proposal to use Heater activation to generate T gradient on boom during Dawn/Dusk orbit.
- Continue the investigation on ASM-VFM residuals using the ASM-V data to constrain the disturbance field.
- Further investigation on the possible impact the plasma-induced perturbations have on the VFM/ASM instrument.

STR:

- IBA correction model: to check the impact this model have on the field modelling

Euler Angles:

- Ionospheric/Magnetospheric currents have been found to be responsible for the apparent change of the Euler Angles. Static Euler angles are suggested to be used in L1B processing.

Data Distribution:

- L1B data produced with corrected quaternions. A TDS to be distributed to full community in preparation of the next DQW (Sept 2016)
- Burst mode ASM data delivered by IPGP to be distributed for the day 19/01/2014
- Generate RPRO MAGx\_HR\_1B data for the day 19/01/2014.
- A complete data package containing all available data for the day 19/01/2014 will be distributed.

IPGP have delivered the Burst mode ASM data for the day 19/01/2014 for the benefit of expert users (/Advanced folder). A complete data package containing all available data for the day 19/01/2014 have also been distributed. Data are available in the ESA FTP server under the path:  
~/Advanced/ASM\_Burst\_Mode\_Data/

DTU have just delivered an updated version of the ASM-VFM residuals, i.e., Baseline and File Counter 0408 covering the period from BOM to 17/07/2015. This dataset has been also extended until 30/06/2016. Data are available in the ESA FTP server under the path:  
~/Advanced/ASM\_VFM\_Residuals/

IRF have delivered a new extended plasma dataset from Swarm Langmuir Probes for the benefit of all the users. Data are available in the ESA FTP server under the path:  
~/Advanced/2\_Hz\_Langmuir\_Probe\_Extended\_Dataset/

IRF is about to deliver the Sweep mode derived electron density and temperature and SC potential dataset, for the benefit of expert users (by end of July).

## **2.4 Summary of observations for 2016, Week 22 (30/05 - 05/06)**

During the monitored week the following events have been found and investigated:

- ASM\_VFM residuals on SC A and B are out of threshold for most of the week. More details in Section 3.3.
- During the heater activation tests the temperatures of VFM ( $T_{CDC}$  and  $T_{CSC}$ ) are non-nominal, for more details please see Section 3.3.3.





### 3. ROUTINE QUALITY CONTROL

#### 3.1 Gaps analysis

- **Several few seconds gaps in MAGx\_CA\_1B products** throughout the week. Some of them seem not to be associated to gaps in telemetry. Monitoring on-going.

#### 3.2 Orbit and Attitude Products

In Table 1 are listed events that have to be reported.

**Table 1:** List of events related to attitude and orbit products to be reported in the monitoring for 2016, Week 22: 30/05 - 05/06.

Observation ID	Description	Affected parameter	Sect. of Obs. Description	Sect. of Obs. Analysis

The relevant parameters that have been monitored are:

- Position difference between calculated Medium Accuracy orbits (**MODx\_SC\_1B**) and on-board solution (**GPSxNAV\_0**). Threshold values for such differences have not been assessed yet: we have just monitored the average values and maximum variations around the week. They are reported in tables in the sections below. In addition, some example plots are given from the HTML daily reports. For the time being we evaluated an anomaly should be raised if one (or more) of the following conditions occurs:
  - o The **average difference** on a given day exceeds the position accuracy requirement for the mission (1.5 m),
  - o The variability around the average is quite high: **standard deviation** threshold has been arbitrarily chosen to be twice the position accuracy requirement for the mission (2-sigma = 3 m).
  - o At least 4-5 spikes are observed on a given day, exceeding +/- 50 m.
- Visual inspection of Star Tracker characterisation flags (**STRxATT\_1B**)
- Deviation of the quaternion norm from unity (deviation threshold = +/- 10<sup>-9</sup>)
- Visual inspection of Euler Angles derived from quaternions.


##### 3.2.1 Position Statistics

In Table 2, one can see the statistics of the differences between MOD and on-board solution positions for S/C A, B and C respectively. In the third column the maximum differences (maximum negative and maximum positive) are reported. The maximum standard deviation is in the fourth column. Maxima, minima and standard deviations usually refer to the Z component that is often the most disturbed; in case another component is most affected, it will be specified in parentheses. Figure 1 shows a cumulative trend of the maximum daily standard deviation for the past 30 days of operations of the MOD-NAV difference, while Figure 2 shows the daily maximum difference, in absolute value, of the MOD-NAV difference, always for the past 30 days of operations.




**Table 2:** Swarm A, B and C, difference between MOD and on-board solution positions. If not specified maximum difference and maximum standard deviation refers to the Z-axis.

<b>Swarm A, 30/05 - 05/06, Position difference</b>					
<b>Day</b>	<b>Average difference (m)</b>	<b>Maximum difference (m)</b>		<b>Maximum standard deviation (m)</b>	<b>Notes</b>
30/05	0.08	-6.6 (Y)	7	1.21	
31/05	0.14	-7.1	6.8	1.26	
01/06	0.08	-8.5	9.8	1.34	
02/06	0.19	-5.8	6.2	1.11	
03/06	0.11	-7.8	5.2	1.05	
04/06	0.02	-5	4.8	1.02	
05/06	0.03	-7.6	7	1.28	
<b>Swarm B, 30/05 - 05/06, Position difference</b>					
<b>Day</b>	<b>Average difference (m)</b>	<b>Maximum difference (m)</b>		<b>Maximum standard deviation (m)</b>	<b>Notes</b>
30/05	0.08	-9.3	9.9	1.41	
31/05	0.22	-6.4	8	1.42	
01/06	0.07	-6.6	9.8	1.29	
02/06	0.09	-6	5.5	1.2	
03/06	0.14	-8.6	6.8	1.25	
04/06	0.06	-5.6	7	1.02	
05/06	0.14	-6.5	7.1 (Y)	1.28	
<b>Swarm C, 30/05 - 05/06, Position difference</b>					
<b>Day</b>	<b>Average difference (m)</b>	<b>Maximum difference (m)</b>		<b>Maximum standard deviation (m)</b>	<b>Notes</b>
30/05	0.16	-8.1	7.1	1.21	
31/05	0.13	-8.8	6.5	1.29	
01/06	0.13	-7	8.1	1.35	
02/06	0.15	-6.6	6.3 (X)	1.11	
03/06	0.09	-7.7	7.5 (X)	1.09	
04/06	0.06	-6.3	5.3	1.04	
05/06	0.09	-7.8	8.2	1.26	



**Figure 1:** Plot of the standard deviation of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.




**Figure 2:** Plot of the maximum difference of the absolute value of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.



### 3.2.1.1 Swarm A

Below is presented plot of MOD-NAV differences for S/C A, taken at the beginning of the week (30/05, Figure 3). From top to bottom, the plots show of MOD-NAV differences in ITRF reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].




**Figure 3:** Difference MOD-GPSNAV, S/C A, 30/05. From top to bottom: X, Y and Z-axis.



### 3.2.1.2 Swarm B

Below is presented plot of MOD-NAV differences for S/C B, taken at the beginning of the week (30/05, Figure 4). From top to bottom, the plots show of MOD-NAV differences in ITRF reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].



**Figure 4:** Difference MOD-GPSNAV, S/C B, 30/05. From top to bottom: X, Y and Z-axis.

### 3.2.1.3 Swarm C

Below is presented plot of MOD-NAV differences for S/C C, taken at the beginning of the week (30/05, Figure 5). From top to bottom, the plots show of MOD-NAV differences in ITRF reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].




Figure 5: Difference MOD-GPSNAV, S/C C, 30/05. From top to bottom: X, Y and Z-axis.

## 3.2.2 Attitude observations

### 3.2.2.1 Swarm A

Nothing to report.

### 3.2.2.2 Swarm B

Nothing to report.

### 3.2.2.3 Swarm C

Nothing to report.

## 3.3 Magnetic Products

For the magnetic products, the weekly monitoring consists in:

- ASM instrument monitoring: quartz frequency (nominal range: [2.949E7 – 2.950E7] Hz) and ASM temperature (temperature range shall be: [-30;+50] °C, Rel. Variation shall not exceed: 0.1 °C/sec).
- VFM instrument monitoring: temperatures (Rel. Variation shall not exceed: 0.1 °C/sec).



- Visual inspection of daily time series of magnetic field intensity  $F$ ,  $B_{NEC}$  and  $B_{VFM}$ . Looking for gaps (or zero values in case of  $MAGx_{LR\_1B}$  products), out-of-threshold values (i.e. exceeding  $\pm 60000$  nT), and other strange features. Map plots of  $F$  and  $B_{NEC}$  for the whole week are then displayed.
- Monitoring of the ASM-VFM known anomaly: visual inspection of  $|B_{VFM}| - F$  taken from  $MAGx_{CA\_1B}$  products and recording of daily maximum variations and standard deviations. If  $\pm 1$  nT are exceeded on a given day, an alert is raised. Map plots of the residuals are shown along with weekly time series of the residuals with and without the "dB\_Sun" correction: in fact, at least a part of the discrepancies found in the measurements between ASM and VFM are modelled through a stray field (dB\_Sun) that is a function of the orientation of the VFM wrt Sun.
- Comparison of magnetic data ( $B_{NEC}$ ) with a model (Chaos5).
- Second derivative of vector field in VFM and NEC frame. Only measurement points within  $\pm 10^\circ$  latitude are considered, and values above 100 nT/s<sup>2</sup> are considered out of threshold.
- 5-min correlations between S/C A and S/C C  $B_{NEC}$  measurements.
- Differences between S/C A and C,  $B_{NEC}$  measurements. Values above 8000 nT are considered out of threshold.

In Table 3 are listed events that have to be reported.

**Table 3:** List of events related to magnetic products to be reported in the monitoring for 2016, Week 22: 30/05 - 05/06.

Observation ID	Description	Affected parameter	Sect. of Obs. Description	Sect. of Obs. Analysis
<b>SW-IDEAS-63</b>	OBS_ROUTINE: $MAGx_{CA\_1B}$ gaps	$MAGx_{CA\_1B}$	3.1	NA

### 3.3.1 VFM-ASM anomaly

**General observation:**

- S/C A – violation of:
  - VFM-ASM residuals threshold on 30/05 - 05/06;
  - mean value of residuals threshold on 30/05, 02/06 - 05/06;
  - standard deviation of residuals threshold on 02/06 - 05/06.
- S/C B – violation of:
  - VFM-ASM residuals threshold on 30/05, 03/06, 05/06;
  - standard deviation of residuals threshold on 31/05 - 05/06.

#### 3.3.1.1 ASM-VFM difference statistics

In Table 4, one can see the statistics of the differences between magnetic field absolute value measured by ASM and by VFM. In the second and third column are reported the maximum differences, maximum negative and maximum positive respectively. The maximum standard deviation is in the fourth column.

The ASM-VFM difference is defined as follow:




$$dF = |B_{VFM}| - F_{ASM}$$

Figure 6 and Figure 7 show the daily mean (circles) and standard deviation (crosses) of dF of the last month for Swarm A and Swarm B respectively.


**Table 4:** Swarm A and B, difference between absolute value of magnetic field measured by ASM and by VFM.

<b>Swarm A, 30/05 - 05/06, ASM-VFM difference</b>					
<b>Day</b>	<b>Max (nT)</b>	<b>Min (nT)</b>	<b>Standard deviation (nT)</b>	<b>Mean (nT)</b>	<b>Notes</b>
30/05	<b>1.05</b>	-0.58	0.35	<b>0.395</b>	
31/05	<b>1.07</b>	-0.76	0.31	0.286	Temp. test
01/06	<b>1.01</b>	-0.69	0.36	0.293	Temp. test
02/06	<b>1.19</b>	-0.78	<b>0.4</b>	<b>0.42</b>	
03/06	<b>1.21</b>	-0.89	<b>0.43</b>	<b>0.411</b>	
04/06	<b>1.11</b>	-0.89	<b>0.4</b>	<b>0.419</b>	
05/06	<b>1.17</b>	-0.86	<b>0.4</b>	<b>0.446</b>	
<b>Swarm B, 30/05 - 05/06, ASM-VFM difference</b>					
<b>Day</b>	<b>Max (nT)</b>	<b>Min (nT)</b>	<b>Standard deviation (nT)</b>	<b>Mean (nT)</b>	<b>Notes</b>
30/05	0.71	<b>-1.01</b>	0.38	-0.113	
31/05	0.8	-0.97	<b>0.42</b>	-0.109	
01/06	0.72	-0.96	<b>0.41</b>	-0.103	
02/06	0.77	-0.96	<b>0.41</b>	-0.088	
03/06	0.77	<b>-1.01</b>	<b>0.45</b>	-0.08	
04/06	0.73	-0.97	<b>0.43</b>	-0.081	
05/06	0.79	<b>-1.02</b>	<b>0.45</b>	-0.072	






**Figure 6:** Daily mean and standard deviation values of ASM-VFM residuals (defined as  $dF=|B_{vfm}|-F_{asm}$ ) for S/C A.



**Figure 7:** Daily mean and standard deviation values of ASM-VFM residuals (defined as  $dF=|B_{vfm}|-F_{asm}$ ) for S/C B.

### 3.3.1.2 Swarm A


The daily peak-to-peak difference around the week stays within [-0.89 - 1.21] nT. Below follow two plots of such differences for current week (Figure 8).



**Figure 8:** ASM-VFM residuals for S/C A, during monitoring period 30/05-05/06. In top figure are plotted: difference between  $|B_{VFM}|$  and  $F_{ASM}$  (without dB<sub>Sun</sub> correction) (blue colour), and the residuals with dB<sub>Sun</sub> corrections (red colour). In bottom figure residuals are presented on the world map.

### 3.3.1.3 Swarm B

The daily peak-to-peak difference around the week stays within [-1.02 - 0.8] nT. Below follow two plots of such differences for current week (Figure 9).



**Figure 9:** ASM-VFM residuals for S/C B, during monitoring period 30/05-05/06. In top figure are plotted: difference between |B<sub>VFM</sub>| and F<sub>ASM</sub> (without dB<sub>Sun</sub> correction) (blue colour), and the residuals with dB<sub>Sun</sub> corrections (red colour). In bottom figure residuals are presented on the world map.

### 3.3.1.4 Swarm C

No data because ASM is switched off.


### 3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC\_0)

For S/C A and B, the temperature and quartz frequency behaved as expected.


### 3.3.3 VFM Instrument parameters: VFM temperatures (MAG\_CA)

The VFM instrument parameters important for monitoring the instrument health are the VFM sensor temperatures: T<sub>CDC</sub>, T<sub>CSC</sub> and T<sub>EU</sub>.

For S/C A, B and C, for reported period, the temperatures behaved as expected, however due to the heater activation test the trend of the VFM temperature on Swarm A is not nominal, please see Figure 10 and Figure 11.



**Figure 10:** T\_CDC temperature of SC A with visible impact of the temperature test.




**Figure 11:** T\_CDC temperature of SC A with visible impact of the temperature test.




### 3.3.4 Magnetic time series visual inspection

#### 3.3.4.1 Swarm A

Map plots of magnetic field measurement for week 22 for S/C A can be seen in Figure 12 below.







**Figure 12:** S/C A, world map plots of the geomagnetic field and components measured during monitoring period 30/05-05/06. From top to bottom: F-magnetic field from ASM measurement, B<sub>NEC</sub> components (North, East, and Centre) of magnetic field from VFM measurement.



### 3.3.4.2 Swarm B

Map plots of magnetic field measurement for week 22 for S/C B can be seen in Figure 13 below.






**Figure 13:** S/C B, world map plots of the geomagnetic field and components measured during monitoring period 30/05-05/06. From top to bottom: F-magnetic field from ASM measurement,  $B_{NEC}$  components (North, East, and Centre) of magnetic field from VFM measurement.



### 3.3.4.3 Swarm C

Map plots of magnetic field measurement for week 22 for S/C C can be seen in Figure 14.




**Figure 14:** S/C C, world map plots of the geomagnetic field and components measured during monitoring period 30/05-05/06. From top to bottom:  $B_{NEC}$  components (North, East, and Centre) of magnetic field from VFM measurement.



### 3.3.5 S/C A and C magnetic correlation


In the plot below is shown the correlation in 5-minutes intervals of magnetic data between satellite A and C. B<sub>north</sub>, B<sub>east</sub>, and B<sub>center</sub> are the components of the magnetic field vector in NEC frame.




**Figure 15:** Correlation in magnetic data between satellite A and C for B<sub>north</sub>, B<sub>east</sub>, and B<sub>center</sub> components of B<sub>NEC</sub> are the components of the magnetic field vector in NEC frame

### 3.3.6 S/C A and C magnetic difference


The next three plots show the differences in magnetic data between satellite A and C. Threshold is set to 8 000 nT for each component.



**Figure 16:** Difference in B<sub>NEC</sub> North component between S/C A and S/C C.



**Figure 17:** Difference in B<sub>NEC</sub> East component between S/C A and S/C C.



**Figure 18:** Difference in B<sub>NEC</sub> Center component between S/C A and S/C C.

### 3.3.7 B<sub>NEC</sub> vs Chaos5 model residuals

The magnetic field measurement is compared to magnetic field estimated from the Chaos5 global geomagnetic field model (only Core and Crustal contributions). Currently in the monitoring routines the external contribution based on Dst index is not taken into account.




Figure 19, Figure 21 and Figure 23 show field residuals  $\Delta B = B_{NEC} - B_{Chaos}$  (all versus co-latitude in degrees), from top to bottom: 1)  $B_r$ , 2)  $B_\theta$  and 3)  $B_\phi$ .

As a general feature one can see the field residuals to be steady and usually below 50 nT at low and middle latitudes, up to  $|\pm 55| - |\pm 60|$  degrees; then the residual increases at high latitudes because the Chaos model does not take into account the contribution from the external field ([RD.17]).

Figure 20, Figure 22 and Figure 24 show, from top to bottom, the time series on 30/05 of: (1-2-3) residuals of  $B_{NEC} - B_{Chaos}$  by components, related to S/C A, B and C respectively.

The component most affected by residual spikes and variations is  $B_{\theta_{NEC}}$ , i.e. the component that shows the variations of the field wrt to co-latitude. At high latitudes, the order of magnitude of the variability is about  $\pm 200$  nT.

### 3.3.7.1 Swarm A



**Figure 19:** S/C A day 30/05  $B_{NEC} - B_{Chaos}$  vs colatitude.




Figure 20: S/C A day 30/05: time series of  $B_{NEC} - B_{Chaos}$  residuals.

### 3.3.7.2 Swarm B




Figure 21: S/C B day 30/05  $B_{NEC} - B_{Chaos}$  difference vs colatitude.




Figure 22: S/C B day 30/05 time series of  $B_{NEC} - B_{Chaos}$  residuals.

### 3.3.7.3 Swarm C




Figure 23: S/C C day 30/05  $B_{NEC} - B_{Chaos}$  difference vs colatitude.




Figure 24: S/C C day 30/05 time series of  $B_{NEC} - B_{Chaos}$  residuals.

### 3.3.8 Second derivative of $B_{NEC}$ and $B_{VFM}$

The second derivative of the vector magnetic field measurements in both VFM and NEC frame has been performed on 50Hz data (MAGx\_HR\_1B). In this analysis only measurement points within  $\pm 10^\circ$  latitude have been considered.








Figure 25, Figure 26 and Figure 27 show the daily standard deviation of the second derivative of  $B_{VFM}$  of the last month for S/C A, B, and C respectively. Second derivative of  $B_{NEC}$  is not shown due to artificial spikes introduced during quaternions interpolation from 1Hz to 50Hz.

**3.3.8.1 Swarm A**



**Figure 25:** Plot of standard deviation of B components in VFM frames.

**3.3.8.2 Swarm B**



**Figure 26:** Plot of standard deviation of B components in VFM frames.



### 3.3.8.3 Swarm C




Figure 27: Plot of standard deviation of B components in VFM frames.





#### **4. ON-DEMAND ANALYSIS**

Nothing to report.



End of Document