IDEAS+ Swarm Weekly Report For Year 2016, Week 36 (05/09 - 11/09)



IDEAS+-SER-OQC-REP-2071 Issue 1.0

Customer : ESRIN Document Ref : IDEAS+-SER-OOC-REP-2071

Contract No : 4000111304/14/I-AM Issue Date : 07 October 2016

WP No : 6110 Issue : 1.0



IDEAS+ Swarm Weekly Report 2016/36: 2016/09/05 - 2016/09/11

Abstract: This is the **Instrument Data Quality Evaluation and Analysis Service Plus** (IDEAS+)

Swarm Weekly report on Swarm products quality, covering the period from 05

September to 11 September 2016.

Author : Approval :

Igino Coco, Jan Miedzik and Enkelejda Qamili on behalf of

Swarm IDEAS+ Team

Marta De Laurentis
IDEAS+ Science and Ops.

Coordinator

Distribution: ESA/ESRIN EOP-GMQ

ESA/ESRIN EOP-GM Swarm MM

IDEAS+ Leadership Team IDEAS+ subcontractors

ESA/ESTEC Swarm PLSO ESA/ESOC Swarm FOS

Copyright © 2015 Telespazio VEGA UK Ltd

All rights reserved.

No part of this work may be disclosed to any third party translated reproduced copied or disseminated in any form or by any means except as defined in the contract or with the written permission of Telespazio Vega UK Ltd.

Telespazio VEGA UK Ltd

350 Capability Green, Luton, Bedfordshire, LU1 3LU, United Kingdom Tel: +44 (0) 1582 399 000 Fax: +44 (0) 1582 728 686 www.telespazio-vega.com

IDEAS+ Swarm Weekly Report For Year 2016, Week 36 (05/09 - 11/09)



TABLE OF CONTENTS


TAE	BLE OF CONTENTS		2
1.	INTRODUCTION		5
1.1		juration of monitored data:	
1.2			
2.	SUMMARY OF THE OBSERVA	ATIONS	7
2.1	General status of Swarm in	struments and Level 1B products quality	7
2.2	Plan for operational proces	ssor updates	7
2.3	Quality Working Group and	d Cal/Val Coordination	7
2.4	Summary of observations f	for 2016, Week 36 (05/09 - 11/09)	8
3.	ROUTINE QUALITY CONTRO	L	9
3.1	Gaps analysis		9
3.2	Orbit and Attitude Product	S	9
3	3.2.1 Position Statistics		9
3	3.2.2 Attitude observations		14
3.3	Magnetic Products		14
3	3.3.1 VFM-ASM anomaly		15
3	3.3.2 ASM Instrument parar	meters: quartz frequency and ASM temperature (ASMAVEC_0)	18
3	3.3.3 VFM Instrument parar	neters: VFM temperatures (MAG_CA)	18
3	3.3.4 Magnetic time series v	risual inspection	19
3	3.3.5 S/C A and C magnetic	correlation	24
3	3.3.6 S/C A and C magnetic	difference	24
3	3.3.7 B _{NEC} vs Chaos5 model	residuals	24
3	3.3.8 Second derivative of B	B _{NEC} and B _{VFM}	28
4	ON-DEMAND ANALYSIS		32





IDEAS+-SER-OQC-REP-2071
Issue 1.0

This page intentionally left blank.





AMENDMENT POLICY

This document shall be amended by releasing a new edition of the document in its entirety. The Amendment Record Sheet below records the history and issue status of this document.

AMENDMENT RECORD SHEET

ISSUE	DATE	REASON
1.0	07 Oct 2016	First issue



1. INTRODUCTION

This document refers to the activities carried out in the framework of the Sensor Performance, Products and Algorithms (SPPA) Office [RD.1], and as such, it reports on work related to:

- Algorithms and Processors Development, Maintenance and Evolution: these include all algorithm and software evolution and maintenance aspects for the different components, for both the Operational processors (OP) and Prototypes processors (PP) of L1 and L2
- Performance Assessment: these include all Quality Control activities (on-line and offline, systematic or on-demand), for the applicable product levels.
- System Calibration: these include the activities related to calibration, from sensor to system level. They also include aspects like cross calibration and handling of external calibration sources.
- Product validation: these include definition and maintenance of product validation plans.
- End-to-end Sensor Dataset Performance: these include activities related to the organisation and coordination of Quality Working Groups and all aspects of the Experimental platform. It also covers the product baseline, coordination and handling of external communities, and all aspects of ADF handling (both for the operational processors and for the prototypes).

This weekly report constitutes a work in progress throughout the mission lifetime, and new parts and complements will be added while the consolidation of knowledge on Swarm data and instruments will progress.

Section 2.1 always gives an overview of the general quality status of the mission instruments and products, while the main observations of the week are summarized in Section 2.1.

The document also includes information on data quality for the three Swarm spacecraft, inferred from automated HTML quality reports, which are produced on daily basis for each product. Please contact the IDEAS+ Swarm team if interested in accessing the reports via web or FTP (all details about interfaces and folder structure available on [RD.2]). Such quality reports represent the core of the Routine Quality Control (Chapter 3). A description of the implemented quality checks is given in [RD.3], and references therein.

Basing on specific findings of the routine quality control, or on-demand from other entities (i.e. Swarm PDGS, FOS, Mission Management, Post-Launch Support Office, Expert Support Laboratories, Quality Working Groups, and user community), anomalies can be triggered. Preliminary characterisations and investigations of such anomalies are given in Chapter 4.The anomalies documented in the Weekly Reports are tracked in the following way:

- 1. If triggered by ESA Eohelp or within the Service: IDEAS+ action and ticketing system (http://requests-sppa.serco.it/RT3/index.html).
- 2. If triggered by IDEAS+ Swarm team or other entities:
 - 2a. if the observation/analysis leads to an anomaly to be addressed to the processor provider (GMV): SPR on EO ARTS (https://arts.eo.esa.int), SWL1L2DB project;
 - 2b. if the observation/analysis does not lead to an anomaly or the investigation shall be escalated to other entities (PLSO/industry, ESL, and PDGS): Action tracked on EO ARTS, SW-IDEAS project, then addressed to the proper tracking system if needed (e.g. JIRA for ESLs, SW-CP-AR project on EO ARTS for PDGS).

Information on Level 1B Swarm products can be found in [RD.4].

1.1 **Current Operational configuration of monitored data:**

- Processors Version: L1BOP 3.17, L2-Cat2 1.15.p5.
- L0 input products baseline: 02

IDEAS+ Swarm Weekly Report For Year 2016, Week 36 (05/09 - 11/09)



IDEAS+-SER-OQC-REP-2071 Issue 1.0

- L1B baseline: MAGNET and PLASMA 04, ORBATT and ACCELE 03 (for definitions and description of the data baseline concept see https://earth.esa.int/web/quest/missions/esaoperational-eo-missions/swarm/data-access/product-baseline-definition)
- Level 2 Cat 2 baseline: EEF 01, IBI, FAC and TEC 02
- Input auxiliary files baseline: S/C A CCDB 0012 (13/09/2016), S/C B CCDB 0012 (16/06/2016), S/C C - CCDB 0012 (13/09/2016), ADF 0101
- MPPF-CVQ v.2.14.01

1.2 **Reference documents**

The following is a list of documents with a direct bearing on the content of this report. Where referenced in the text, these are identified as RD.n, where 'n' is the number in the list below:

- Sensor Performance, Products and Algorithms (SPPA), PGSI-GSOP-EOPG-TN-05-0025. [RD.1] Version 2.3.
- Swarm PDGS External DMC Interface Control Document, SW-ID-DS-GS-0001, Issue 3.2. [RD.2]
- Swarm MPPF-CVQ Monitoring Baseline Document, ST-ESA-SWARM-MBD-0001, Issue 1.7. [RD.3]
- Swarm Level 1B Product Definition, SW-RS-DSC-SY-0007, Issue 5.13. [RD.4]
- [RD.5] Swarm IDEAS Configuration Management Plan, IDEAS-SER-MGT-PLN-1081 v0.14.
- Swarm Quality Control Project Plan, IDEAS-SER-MGT-PLN-1071 [RD.6]
- SW L1BOP status 20141124 MoM [RD.7]
- [RD.8] Planned Updates for Level 1b, SW-PL-DTU-GS-008, Rev: 1dC.
- [RD.9] IDEAS+ Swarm Weekly Report: 25/08/2014 - 31/08/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20140825_20140831.pdf (ref. for SWL1L2DB-9)
- [RD.10] IDEAS+ Swarm Weekly Report: 29/09/2014 05/10/2014, IDEAS+-SER-OQC-REP-2071 SPPA SwarmWeeklyReport 20140929 20141005.pdf (ref. for SW-IDEAS-34)
- [RD.11] IDEAS+ Swarm Weekly Report: 06/10/2014 12/10/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20141006_20141012.pdf (ref. for SW-IDEAS-36)
- [RD.12] IDEAS+ Swarm Weekly Report: 20/10/2014 26/10/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20141020_20141026.pdf (ref. for SW-IDEAS-40, GPS sync loss)
- [RD.13] IDEAS+ Swarm Weekly Report: 15/09/2014 21/09/2014, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_20140915_20140921.pdf (ref. for SW-IDEAS-27)
- [RD.14] Swarm L1B 03.15 Validation Report, OSMV-OPMT-SRCO-RP-15-3385, Issue 1.3.
- [RD.15] IDEAS+ Swarm Weekly Report: 23/03/2015 29/03/2015, IDEAS+-SER-OQC-REP-2071_SPPA_SwarmWeeklyReport_201513_20150323_20150329.pdf.
- [RD.16] SWARM Weekly Operations Report #76, SW-RP-ESC-FS-6172
- [RD.17] Olsen, N., H. Luhr, C.C. Finlay, T.J. Sabaka, I. Michaelis, J. Rauberg and L. Tøffner-Clausen, The CHAOS-4 geomagnetic field model, Geophys. J. Int. 197, 815-827, 2014
- [RD.18] IDEAS+-SER-IPF-PLN-2272, Swarm Level 1B Operational Processor Verification Plan, IDEAS+-SER-IPF-PLN-2272_L1BOP_316_v1.5_final.pdf



2. SUMMARY OF THE OBSERVATIONS

2.1 General status of Swarm instruments and Level 1B products quality

Nothing to report

2.2 Plan for operational processor updates

L1BOP: In the next delivery of the L1BOP after the PDGS evolution activities (beginning of Nov. 2016) the following changes will be implemented into the MAGNET and ORBATT processors:

- Change the data resampling used for the interpolation of the ASM measurements to UTC second because the present used in L1B data is removing a significant part of the high frequency content of the ASM data.
- Modification of the F_error computation.
- More robust handling of the leap second occurrence, and modification in timestamp sorting, in order to make the processor complete the run even in case of VFM on-board anomalies, that could imply a packet frequency greater than 1 Hz.
- STR Inter-Boresight Angles correction model.
- The activity of PLASMA cross-verification has been restarted and hopefully a fully cross-verified PLASMA processor will be included in the next delivery.
- Changes in the CCDB with inclusion of several new parameters is foreseen for many of the implementations above.

For what concern the ORBATT processor, another change to be implemented is under discussion.

• An increase of the frequency of the STR LO product from 1 Hz to 2 Hz

L2 Cat-2: Any other delivery of the L2 Cat-2 OP is postponed after the PDGS evolution activity (Nov. 2016).

PDGS evolution related activity: Currently, the deployment in operations of L1BOP 3.18 and L2CAT2OP 1.16, including the update to the full CDF processing schema, is foreseen for mid-November 2016.

2.3 Quality Working Group and Cal/Val Coordination

The former "Task Force" deputed to study the ASM-VFM residuals, has been reshaped in a Magnetic Measurements Expert Group (MMEG). The first MMEG meeting was held in Warsaw (GMV premises) on **10-11/03/2016**. During this meeting the following decisions have been taken:

ASM-VFM Scalar Residual:

VFM calibration issue:

IDEAS+ Swarm Weekly Report For Year 2016, Week 36 (05/09 - 11/09)



IDEAS+-SER-OQC-REP-2071 Issue 1.0

- Finalize the covariance analysis between parameters in alpha/beta space
- Further analysis directly on non-orthogonalities in-flight and on ground test data and not the computed models.
- Scale factor s_{exp} shall be further investigated.
- Consolidate the analysis on dependency on sbeta and TEU.
- Further investigation on the correlation between ASM-VFM residuals and T_{EU}.
- Continue the investigation on ASM-VFM residuals using MAGx_HR_1B data.
- Consolidate test proposal to use Heater activation to generate T gradient on boom during Dawn/Dusk orbit.
- Continue the investigation on ASM-VFM residuals using the ASM-V data to constrain the disturbance field.
- Further investigation on the possible impact the plasma-induced perturbations have on the VFM/ASM instrument.

STR:

- IBA correction model: to check the impact this model have on the field modelling Euler Angles:
- Ionospheric/Magnetospheric currents have been found to be responsible for the apparent change of the Euler Angles. Static Euler angles are suggested to be used in L1B processing. Data Distribution:
 - L1B data produced with corrected quaternions. A TDS to be distributed to full community in preparation of the next DQW (Sept 2016)
 - Burst mode ASM data delivered by IPGP to be distributed for the day 19/01/2014
 - Generate RPRO MAGx_HR_1B data for the day 19/01/2014.
 - A complete data package containing all available data for the day 19/01/2014 will be distributed.

IRF is about to deliver the Sweep mode derived electron density and temperature and SC potential dataset, for the benefit of expert users (by end of September).

2.4 Summary of observations for 2016, Week 36 (05/09 - 11/09)

During the monitored week the following events have been found and investigated:

- ASM-VFM mean value SC A soft threshold violation.
- After introduction of new CCDB for SC A and SC C no discontinuities found.



3. ROUTINE QUALITY CONTROL

3.1 Gaps analysis

• **Several few seconds gaps in MAGx_CA_1B products** throughout the week. Some of them seem not to be associated to gaps in telemetry. Monitoring on-going.

3.2 Orbit and Attitude Products

In Table 1 are listed events that have to be reported.

Table 1: List of events related to attitude and orbit products to be reported in the monitoring for 2016, Week 36: 05/09 - 11/09.

Observation ID	Description	Affected parameter	Sect. of Obs. Description	Sect. of Obs. Analysis

The relevant parameters that have been monitored are:

- Position difference between calculated Medium Accuracy orbits (MODx_SC_1B) and on-board solution (GPSxNAV_0). Threshold values for such differences have not been assessed yet: we have just monitored the average values and maximum variations around the week. They are reported in tables in the sections below. In addition, some example plots are given from the HTML daily reports. For the time being we evaluated an anomaly should be raised if one (or more) of the following conditions occurs:
 - The **average difference** on a given day exceeds the position accuracy requirement for the mission (1.5 m),
 - The variability around the average is quite high: standard deviation threshold has been arbitrarily chosen to be twice the position accuracy requirement for the mission (2-sigma = 3 m).
 - o At least 4-5 spikes are observed on a given day, exceeding +/- 50 m.
- Visual inspection of Star Tracker characterisation flags (STRxATT_1B)
- Deviation of the quaternion norm from unity (deviation threshold = \pm 10⁻⁹)
- Visual inspection of Euler Angles derived from quaternions.

3.2.1 Position Statistics

In Table 2, one can see the statistics of the differences between MOD and on-board solution positions for S/C A, B and C respectively. In the third column the maximum differences (maximum negative and maximum positive) are reported. The maximum standard deviation is in the fourth column. Maxima, minima and standard deviations usually refer to the Z component that is often the most disturbed; in case another component is most affected, it will be specified in parentheses. Figure 1 shows a cumulative trend of the maximum daily standard deviation for the past 30 days of operations of the MOD-NAV difference, while Figure 2 shows the daily maximum difference, in absolute value, of the MOD-NAV difference, always for the past 30 days of operations.



IDEAS+-SER-OQC-REP-2071 Issue 1.0

Table 2: Swarm A, B and C, difference between MOD and on-board solution positions. If not specified maximum difference and maximum standard deviation refers to the Z-axis.

	Specifica maximus	Transcrete and	, maximam starr	uard deviation refers to the	L data.
		Swarm	A, 05/09 - 11/0	9, Position difference	
Day	Average difference (m)	Maximum difference (m)		Maximum standard deviation (m)	Notes
05/09	0.04	-7.5 5.3		1.2	
06/09	0.24	-6 (X)	8	1.31	
07/09	0.1	-8.3	6.9 (Y)	1.35	
08/09	0.15	-8.5	10.8	1.33	
09/09	0.11	-5.8	6.3	1.21	
10/09	0.06	-7.3	6.6 (Y)	1.11	
11/09	0.04	-6.8	5.5	1.03	
		Swarm	B, 05/09 - 11/0	9, Position difference	
Day	Average difference (m)	Maximum difference (m)		Maximum standard deviation (m)	Notes
05/09	0.04	-8.2	9	1.29	
06/09	0.13	-8.1	8	1.27	
07/09	0.13	-13.9	8.5	1.5	
08/09	0.06	-6.5	7.6	1.27	
09/09	0.12	-7.7	7	1.19	
10/09	0.12	-5.7	5.4	1.11	
11/09	0.14	-8.5	6.2	1	
		Swarm	C, 05/09 - 11/0	9, Position difference	
Day	Average difference (m)	Maximum difference (m)		Maximum standard deviation (m)	Notes
05/09	0.11	-7	12.3	1.18	
06/09	0.2	-7.4	6.7	1.36	
07/09	0.11	-6.3 8.3 (Y)		1.31	
08/09	0.14	-6.3 (Y) 6.9 (X)		1.26	
09/09	0.11	-5.6	6	1.21	
10/09	0.08	-6 (X)	5.8	1.1	
11/09	11/09 0.08 -6.8 4.9		1.05		






Figure 1: Plot of the standard deviation of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.




Figure 2: Plot of the maximum difference of the absolute value of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.



3.2.1.1 Swarm A

Below is presented plot of MOD-NAV differences for S/C A, taken at the beginning of the week (05/09, Figure 3). From top to bottom, the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].




Figure 3: Difference MOD-GPSNAV, S/C A, 05/09. From top to bottom: X, Y and Z-axis



3.2.1.2 Swarm B

Below is presented plot of MOD-NAV differences for S/C B, taken at the beginning of the week (05/09, Figure 4). From top to bottom, the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].




Figure 4: Difference MOD-GPSNAV, S/C B, 05/09. From top to bottom: X, Y and Z-axis



3.2.1.3 Swarm C

Below is presented plot of MOD-NAV differences for S/C C, taken at the beginning of the week (05/09, Figure 5). From top to bottom, the plots show of MOD-NAV differences in ITFR reference frame: on X, Y and Z-axis respectively. The difference between both solutions is given in [m].




Figure 5: Difference MOD-GPSNAV, S/C C, 05/09. From top to bottom: X, Y and Z-axis

3.2.2 Attitude observations

3.2.2.1 Swarm A

Nothing to report.

3.2.2.2 Swarm B

Nothing to report.

3.2.2.3 Swarm C

Nothing to report.

3.3 Magnetic Products

For the magnetic products, the weekly monitoring consists in:

- ASM instrument monitoring: quartz frequency (nominal range: [2.949E7 2.950E7] Hz) and ASM temperature (temperature range shall be: [-30;+50] °C, Rel. Variation shall not exceed: 0.1 °C/sec).
- VFM instrument monitoring: temperatures (Rel. Variation shall not exceed: 0.1 °C/sec).



IDEAS+-SER-OQC-REP-2071 Issue 1.0

- Visual inspection of daily time series of magnetic field intensity F, B_{NEC} and B_{VFM} . Looking for gaps (or zero values in case of MAGx_LR_1B products), out-of-threshold values (i.e. exceeding +/- 60000 nT), and other strange features. Map plots of F and B_{NEC} for the whole week are then displayed.
- Monitoring of the ASM-VFM known anomaly: visual inspection of |B_{VFM}| F taken from MAGx_CA_1B products and recording of daily maximum variations and standard deviations. If +/- 1 nT are exceed on a given day, an alert is raised. Map plots of the residuals are shown along with weekly time series of the residuals with and without the "dB_Sun" correction: in fact, at least a part of the discrepancies found in the measurements between ASM and VFM are modelled through a stray field (dB_Sun) that is a function of the orientation of the VFM wrt Sun.
- Comparison of magnetic data (B_{NEC}) with a model (Chaos5).
- Second derivative of vector field in VFM and NEC frame. Only measurement points within ±10° latitude are considered, and values above 100 nT/s2 are considered out of threshold.
- 5-min correlations between S/C A and S/C C B_{NEC} measurements.
- Differences between S/C A and C, BNEC measurements. Values above 8000 nT are considered out of threshold.

In Table 3 are listed events that have to be reported.

Table 3: List of events related to magnetic products to be reported in the monitoring for 2016, Week 36: 05/09 - 11/09.

Observation ID	Description	Affected parameter	Sect. of Obs. Description	Sect. of Obs. Analysis
SW-IDEAS-63	OBS_ROUTINE: MAGx_CA_1B gaps	MAGx_CA_1B	3.1	NA

3.3.1 VFM-ASM anomaly

General observation:

- S/C A violation of:
 - VFM-ASM residuals threshold on 05/09, 08/09, 09/09;
 - o mean value of residuals threshold on 05/09 11/09.

3.3.1.1 ASM-VFM difference statistics

In Table 4, one can see the statistics of the differences between magnetic field absolute value measured by ASM and by VFM. In the second and third column are reported the maximum differences, maximum negative and maximum positive respectively. The maximum standard deviation is in the fourth column.

The ASM-VFM difference is defined as follow:

 $dF = |B_{VFM}| - F_{ASM}$

Figure 6 and Figure 7 show the daily mean (circles) and standard deviation (crosses) of dF of the last month for Swarm A and Swarm B respectively.



Table 4: Swarm A and B, difference between absolute value of magnetic field measured by ASM and by VFM.

	and by vrivi.				
		Swarm A, (05/09 - 11/09, ASM-VFM c	lifference	
Day	Max (nT)	Min (nT)	Standard deviation (nT)	Mean (nT)	Notes
05/09	1.09	-0.68	0.34	0.395	
06/09	0.99	-0.71	0.31	0.344	
07/09	0.92	-0.7	0.29	0.328	
08/09	1.07	-0.56	0.27	0.337	
09/09	1.14	-0.56	0.25	0.327	
10/09	0.98	-0.45	0.21	0.326	Updated CCDB
11/09	0.88	-0.42	0.19	0.311	Updated CCDB
		Swarm B, (05/09 - 11/09, ASM-VFM c	lifference	
Day	Max (nT)	Min (nT)	Standard deviation (nT)	Mean (nT)	Notes
05/09	0.82	-0.65	0.31	0.105	
06/09	0.68	-0.82	0.29	0.078	
07/09	0.64	-0.84	0.31	0.058	
08/09	0.68	-0.88	0.31	0.052	
09/09	0.68	-0.85	0.31	0.043	
10/09	0.64	-0.69	0.27	0.021	
11/09	0.74	-0.62	0.28	0.007	




Figure 6: Daily mean and standard deviation values of ASM-VFM residuals (defined as $dF = |B_{VFM}| - F_{ASM}$) for S/C A.






Figure 7: Daily mean and standard deviation values of ASM-VFM residuals (defined as $dF = |B_{VFM}| - F_{ASM}$) for S/C B.

3.3.1.2 Swarm A

The daily peak-to-peak difference around the week stays within [-0.71 - 1.14] nT. Below follow two plots of such differences for current week (Figure 8).




Figure 8: ASM-VFM residuals for S/C A, during monitoring period 05/09-11/09. In top figure are plotted: difference between |B_VFM| and F_ASM (without dB_Sun correction) (blue colour), and the residuals with dB_Sun corrections (red colour). In bottom figure residuals are presented on the world map.



3.3.1.3 Swarm B

The daily peak-to-peak difference around the week stays within [-0.88 - 0.82] nT. Below follow two plots of such differences for current week (Figure 9).




Figure 9: ASM-VFM residuals for S/C B, during monitoring period 05/09-11/09. In top figure are plotted: difference between |B_VFM| and F_ASM (without dB_Sun correction) (blue colour), and the residuals with dB_Sun corrections (red colour). In bottom figure residuals are presented on the world map.

3.3.1.4 Swarm C

No data because ASM is switched off.

3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC_0)

For S/C A and B, the temperature and quartz frequency behaved as expected.

3.3.3 VFM Instrument parameters: VFM temperatures (MAG_CA)

The VFM instrument parameters important for monitoring the instrument health are the VFM sensor temperatures: T_CDC, T_CSC and T_EU.


For S/C A, B and C, for reported period, the temperatures behaved as expected.




3.3.4 Magnetic time series visual inspection


3.3.4.1 Swarm A

Map plots of magnetic field measurement for week 36 for S/C A can be seen in Figure 10 below.














Figure 10: S/C A, world map plots of the geomagnetic field and components measured during monitoring period 05/09-11/09. From top to bottom: F-magnetic field from ASM measurement, B_{NEC} components (North, East, and Centre) of magnetic field from VFM measurement.




3.3.4.2 Swarm B

Map plots of magnetic field measurement for week 36 for S/C B can be seen in Figure 11 below.














Figure 11: S/C B, world map plots of the geomagnetic field and components measured during monitoring period 05/09-11/09. From top to bottom: F-magnetic field from ASM measurement, B_{NEC} components (North, East, and Centre) of magnetic field from VFM measurement.



3.3.4.3 Swarm C

Map plots of magnetic field measurement for week 36 for S/C C can be seen in Figure 12.








Figure 12: S/C C, world map plots of the geomagnetic field and components measured during monitoring period 05/09-11/09. From top to bottom: B_{NEC} components (North, East, and Centre) of magnetic field from VFM measurement.



3.3.5 S/C A and C magnetic correlation

In the plot below is shown the correlation in 5-minutes intervals of magnetic data between satellite A and C. B_north, B_east, and B_center are the components of the magnetic field vector in NEC frame.




Figure 13: Correlation in magnetic data between satellite A and C for B_north, B_east, and B_center components of B_{NEC} are the components of the magnetic field vector in NEC frame

3.3.6 S/C A and C magnetic difference

The next three plots show the differences in magnetic data between satellite A and C. Threshold is set to 8 000 nT for each component.




Figure 14: Difference in B_NEC North component between S/C A and S/C C.




Figure 15: Difference in B_NEC East component between S/C A and S/C C.




Figure 16: Difference in B_NEC Center component between S/C A and S/C C.

3.3.7 B_{NEC} vs Chaos5 model residuals

The magnetic field measurement is compared to magnetic field estimated from the Chaos5 global geomagnetic field model (only Core and Crustal contributions). Currently in the monitoring routines the external contribution based on Dst index is not taken into account.



Figure 17, Figure 19 and Figure 21 show field residuals $dB=B_{NEC}$ - B_{Chaos} (all versus co-latitude in degrees), from top to bottom: 1) Br, 2) B θ and 3) $B\phi$.

As a general feature one can see the field residuals to be steady and usually below 50 nT at low and middle latitudes, up to |55| - |60| degrees; then the residual increases at high latitudes because the Chaos model does not take into account the contribution from the external field ([RD.17]).

Figure 18, Figure 20 and Figure 22 show, from top to bottom, the time series on 05/09 of: (1-2-3) residuals of B_{NEC}-B_{Chaos} by components, related to S/C A, B and C respectively.

The component most affected by residual spikes and variations is $B\theta_{NEC}$, i.e. the component that shows the variations of the field wrt to co-latitude. At high latitudes, the order of magnitude of the variability is about +/- 200 nT.

3.3.7.1 Swarm A




Figure 17: S/C A day 05/09 B_{NEC} - B_{Chaos} vs colatitude.






Figure 18: S/C A day 05/09: time series of $B_{NEC} - B_{Chaos}$ residuals.

3.3.7.2 Swarm B




Figure 19: S/C B day 05/09 B_{NEC} - B_{Chaos} difference vs colatitude.






Figure 20: S/C B day 05/09 time series of B_{NEC} – B_{Chaos} residuals.

3.3.7.3 Swarm C




Figure 21: S/C C day $05/09\ B_{NEC}$ - B_{Chaos} difference vs colatitude.






Figure 22: S/C C day 05/09 time series of B_{NEC} – B_{Chaos} residuals.

3.3.8 Second derivative of B_{NEC} and B_{VFM}

The second derivative of the vector magnetic field measurements in both VFM and NEC frame has been performed on 50Hz data (MAGx_HR_1B). In this analysis only measurement points within $\pm 10^{\circ}$ latitude have been considered. Figure 23, Figure 24 and Figure 25 show the daily standard deviation of the second derivative of B_{NEC} and B_{VFM} of the last month for S/C A, B, and C respectively. The spikes (2^{nd} derivative value above 100) is in all cases caused by artificial spike introduced during quaternions interpolation from 1Hz to 50Hz (investigated).



3.3.8.1 Swarm A




Figure 23: Plot of standard deviation of the second derivative of B components in VFM and NEC frames.



Swarm B




Figure 24: Plot of standard deviation of the second derivative of B components in VFM and NEC frames.



3.3.8.2 Swarm C





Figure 25: Plot of standard deviation of the second derivative of B components in VFM and NEC frames.





4. ON-DEMAND ANALYSIS

Nothing to report.

IDEAS+ Swarm Weekly Report For Year 2016, Week 36 (05/09 - 11/09)



End of Document