


---

# Swarm DISC Weekly Report 2022/47: 2022/11/21 - 2022/11/27

---



**Abstract** : This is the **Swarm Data Innovation and Science Cluster** (Swarm DISC) Weekly report on Swarm products quality, covering the period from 21 November to 27 November 2022.

**Doc. No** : SW-RP-SER-GS-010

<b>Author</b> :	<b>Approval</b> :
Đorđe Stevanović, Nicola Comparetti and Maria E Mazzocato on behalf of Swarm DISC Team	Jerome Bouffard ESA – EOP/GMQ

<b>Distribution</b> :	
ESA/ESRIN EOP-GMQ ESA/ESRIN EOP-GM Swarm MM Swarm DISC Management Team Swarm DISC subcontractors	ESA/ESTEC Swarm PLSO ESA/ESOC Swarm FOS



## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 2 of 22

# TABLE OF CONTENTS

<b>TABLE OF CONTENTS .....</b>	<b>2</b>
<b>1. INTRODUCTION .....</b>	<b>4</b>
1.1 Annual Swarm Data Quality Workshop .....	4
1.2 Current Operational configuration of monitored data: .....	5
1.3 Recent evolutions: .....	5
1.4 Reference documents .....	5
<b>2. SUMMARY OF THE OBSERVATIONS .....</b>	<b>7</b>
2.1 Changes in the general status of Swarm instruments and Level 1B products quality .....	7
2.2 Relevant observations of the week 47 (21/11 - 27/11) .....	7
<b>3. ROUTINE QUALITY CONTROL .....</b>	<b>8</b>
3.1 Gaps analysis .....	8
3.2 Orbit and Attitude Products .....	8
3.2.1 Position Statistics .....	8
3.2.2 Attitude observations .....	9
3.2.2.1 Swarm A .....	9
3.2.2.2 Swarm B .....	9
3.2.2.3 Swarm C .....	9
3.3 Magnetic Products .....	10
3.3.1 VFM-ASM anomaly .....	10
3.3.1.1 ASM-VFM difference statistics .....	10
3.3.1.2 Swarm A .....	11
3.3.1.3 Swarm B .....	12
3.3.1.4 Swarm C .....	12
3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC_0) .....	12
3.3.3 VFM Instrument parameters: VFM temperatures (MAG_CA) .....	12
3.3.4 Magnetic time series visual inspection .....	13
3.3.4.1 Swarm A .....	13
3.3.4.2 Swarm B .....	14
3.3.4.3 Swarm C .....	15
3.3.5 $B_{NEC}$ vs Chaos7 model residuals .....	15
3.3.5.1 Swarm A .....	16
3.3.5.2 Swarm B .....	16
3.3.5.3 Swarm C .....	17
3.4 Plasma Products .....	17
3.4.1 Plasma time series visual inspection .....	18
3.4.1.1 Swarm A .....	18
3.4.1.2 Swarm B .....	18
3.4.1.3 Swarm C .....	19
3.4.2 Plasma products latitudinal variations .....	19
3.4.2.1 Swarm A .....	19
3.4.2.2 Swarm B .....	20
3.4.2.3 Swarm C .....	20
<b>4. SPECIAL INVESTIGATIONS .....</b>	<b>21</b>



## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 3 of 22

### **AMENDMENT POLICY**

This document shall be amended by releasing a new edition of the document in its entirety. The Amendment Record Sheet below records the history and issue status of this document.

### **AMENDMENT RECORD SHEET**

<b>ISSUE</b>	<b>DATE</b>	<b>REASON</b>
1.0	16 Jan 2023	First issue

## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 4 of 22

# 1. Introduction

This document refers to the activities carried out in the framework of the ESA Sensor Performance, Products and Algorithms (SPPA) Office [RD. 01].

Chapter 1 gives an overview on the outcomes from the annual Swarm Data Quality Workshop and reports the information on the current operational configuration and its future improvements. It also contains the list of used reference documents.

In Chapter 2, the Section 2.1 gives an overview of the general quality status of the Swarm mission instruments and products, while the main observations of the week are summarized in the Section 2.2.

The document also includes information on data quality for the three Swarm spacecraft, inferred from automated HTML quality reports, which are produced on daily basis for each product. If interested in accessing the reports via web or FTP, please contact the Swarm DISC team at the following email address: <swarm@eo-sppa.org>. Such data quality reports represent the main component of the Routine Quality Control performed by ESA SPPA (Chapter 3). A description of the implemented quality checks is given in [RD. 02], and references therein.

Based on specific findings of the routine quality control, or requests from other entities (i.e. Swarm Payload Data Ground Segment (PDGS), Flight Operation Segment (FOS), Mission Management, Post-Launch Support Office (PLSO), Expert Support Laboratories (ESL), Quality Working Groups (QWG), and user community), investigations on anomalies can be triggered. Preliminary characterisations on such anomalies are given in Chapter 4.

Information on Swarm Level 1B products can be found in [RD. 03].

**This weekly report is based on QC methods and diagnostics that tend to be continuously evolved and improved throughout the mission lifetime, reporting on the data quality, product evolutions, and status of the instruments on weekly basis.**

## 1.1 Annual Swarm Data Quality Workshop

The 12th Swarm Data Quality Workshop was hosted by the Swedish Institute of Space Physics (IRFU) in Uppsala, Sweden, from 10 to 14 October 2022. Thanks to the participation of more than 150 scientists (in presence and remotely) from different institutions in Europe, America and Asia, the event was a great success.

The DQW#12, structured in 10 thematic sessions including talks and dedicated time slots for discussions, was instrumental in addressing the processing and use of Swarm data and defining a road map for future activities.

The main topics addressed during the workshop were related to:

- collecting feedback about new Swarm dataset generated with an improved version of L1B and L2-Cat2 operational processors
- identifying possible new Swarm data products and services, with particular attention to Space Weather applications
- promoting synergies with other satellite missions
- discussing on future Swarm-based science applications and orbital constellation evolution

More information on the Swarm DQW#12 outcomes, summary and recommendations will be published soon.

## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 5 of 22

### 1.2 Current Operational configuration of monitored data:

Processor	
Name	Version
L1BOP	v3.24p4
L2-Cat2	v01.20
Products	
Name	Baseline
L0 inputs	02
L1B MAGNET and PLASMA	06
L1B ORBATT and ACCELE	05
L2-Cat2 EEF	02
L2-Cat2 IBI, FAC and TEC	04
Others	
Input auxiliary files	S/C A, CCDB 0028 (27/09/2022) S/C B, CCDB 0028 (27/09/2022) S/C C, CCDB 0029 (27/09/2022)
MPPF-CVQ	v03.13 (18/03/2022)

### 1.3 Recent evolutions:

An improved version for both L1B and L2 (FAC, IBI, and TEC) Operational Processors was transferred into operation on 28/09/2022.

The main improvements introduced in L1B data processing chain are:

- Improved thruster activation information
- New Flags\_q parameter
- Removed duplicated timestamp in GPS RINEX files
- Improved thermal model for Inter Boresight Angle (IBA) variation
- Improved dB\_Sun and new dF\_Sun stray fields
- Improved F\_ASM data processing during Burst Mode sessions
- Production of MAGNET data when no VFM data are available
- Usage of POD as input for MAGNET
- New LP calibration parameter
- Improved Flags for Plasma products

Please note that the L1B data improvements have a negligible impact on L2 IBI, FAC and TEC products. Moreover, the porting to an updated operational system was performed for both L1B and L2 Operational processors, without any impact on Swarm L1B and L2 data content. For more information please refer to the Technical Note [RD .05].

### 1.4 Reference documents

The following is a list of documents with a direct bearing on the content of this report. Where referenced in the text, these are identified as RD.n, where 'n' is the number in the list below:

[RD. 01] Sensor Performance, Products and Algorithms (SPPA), PGSI-GSOP-EOPG-TN-05-0025. Version 2.3.

[RD. 02] Swarm MPPF-CVQ Monitoring Baseline Document, ST-ESA-SWARM-MBD-0001, Issue 1.7.



## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 6 of 22

- 
- [RD. 03] Swarm Level 1B Product Definition, SW-RS-DSC-SY-0007, Issue 5.23.
  - [RD. 04] Olsen, N., H. Luhr, C.C. Finlay, T.J. Sabaka, I. Michaelis, J. Rauberg and L. Tøffner-Clausen, The CHAOS-4 geomagnetic field model, Geophys. J. Int. 197, 815–827, 2014
  - [RD. 05] <https://earth.esa.int/documents/10174/1514862/Swarm-L1B-and-L2-operational-processors.pdf>



## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 7 of 22

## 2. Summary of the observations

### 2.1 Changes in the general status of Swarm instruments and Level 1B products quality

- The Absolute Scalar Magnetometer (ASM) on board Swarm Bravo is commanded in burst mode from 18/11/2022 at 00:00 UT to 25/11/2022 at 00:00 UT.
- The Absolute Scalar Magnetometer (ASM) on board Swarm Alpha is commanded in burst mode from 21/11/2022 at 00:00 UT to 28/11/2022 at 00:00 UT.
- Due to an issue detected in the Swarm L1B data processing chain, several data gaps are present in the magnetic field vector components derived from Swarm Alpha VFM instrument starting from 20 November 2022 at 23:08 UTC.

### 2.2 Relevant observations of the week 47 (21/11 - 27/11)

During the monitored week the following no events have been found.

## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 8 of 22

### 3. Routine Quality control

#### 3.1 Gaps analysis

- 20th and 21st November:

F\_ASM data stored in MAGx\_LR\_1B for Swarm Alpha is set to zero from 20/11/2022 at 23:59:30 to 21/11/2022 at 00:00:01 UT. A data gap in MAGx\_CA\_1B production is present in the same time frame

- 24th and 25th November:

F\_ASM data stored in MAGx\_LR\_1B for Swarm Bravo is set to zero from 24/11/2022 at 23:59:28 to 25/11/2022 at 00:00:24 UT. A data gap in MAGx\_CA\_1B production is present in the same time frame.

- 27th and 28th November:

F\_ASM data stored in MAGx\_LR\_1B for Swarm Alpha is set to zero from 27/11/2022 at 23:59:28 to 28/11/2022 at 00:00:24 UT. A data gap in MAGx\_CA\_1B production is present in the same time frame.

#### 3.2 Orbit and Attitude Products

The relevant parameters that have been monitored are:

- Position difference between calculated Medium Accuracy orbits (MODx\_SC\_1B) and on-board solution (GPSxNAV\_0). Threshold values for such differences have not been assessed yet: we have just monitored the average values and maximum variations within the week. For the time being we evaluated an anomaly should be raised if one (or more) of the following conditions occurs:
  - The **average difference** on a given day exceeds the position accuracy requirement for the mission (1.5 m),
  - The variability around the average is quite high: **standard deviation** threshold has been arbitrarily chosen to be twice the position accuracy requirement for the mission (2-sigma = 3 m).
  - At least 4-5 spikes are observed on a given day, exceeding +/- 50 m.
- Visual inspection of Star Tracker characterisation flags (STRxATT\_1B)
- Deviation of the quaternion norm from unity (deviation threshold = +/- 10<sup>-9</sup>)
- Visual inspection of Euler Angles derived from quaternions.

##### 3.2.1 Position Statistics

In Figure 3-1 and Figure 3-2 one can see the statistics of the differences between MOD and on-board solution positions for S/C A, B and C respectively. Figure 3-1 shows a cumulative trend of the






Swarm DISC Weekly Report


Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 9 of 22

maximum daily standard deviation for the past 30 days of operations of the MOD-NAV difference, while Figure 3-2 shows the daily maximum difference, in absolute value, of the MOD-NAV difference, always for the past 30 days of operations.



**Figure 3-1:** Plot of the standard deviation of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.



**Figure 3-2:** Plot of the maximum difference of the absolute value of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.

### 3.2.2 Attitude observations

#### 3.2.2.1 Swarm A

Nominal. Nothing to report.

#### 3.2.2.2 Swarm B

Nominal. Nothing to report.

#### 3.2.2.3 Swarm C

- 26th November:  
Swarm-C STR data gap from 12:47:30 to 12:47:46

## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 10 of 22

### 3.3 Magnetic Products

For the magnetic products, the weekly monitoring consists of:

- ASM instrument monitoring: quartz frequency (nominal range: [2.949E7 – 2.950E7] Hz) and ASM temperature (temperature range shall be: [-30;+50] °C, Rel. Variation shall not exceed: 0.1 °C/sec).
- VFM instrument monitoring: temperatures (Rel. Variation shall not exceed: 0.1 °C/sec).
- Visual inspection of daily time series of magnetic field intensity  $F$ ,  $B_{NEC}$  and  $B_{VFM}$ . Looking for gaps (or zero values in case of  $MAGx_{LR\_1B}$  products), out-of-threshold values (i.e. exceeding +/- 60000 nT), and other strange features. Map plots of  $F$  and  $B_{NEC}$  for the whole week are then displayed.
- Monitoring of the ASM-VFM known anomaly: visual inspection of  $|B_{VFM}| - F$  taken from  $MAGx_{CA\_1B}$  products and recording of daily maximum variations and standard deviations. If +/- 1 nT are exceeded on a given day, an alert is raised. Map plots of the residuals are shown along with weekly time series of the residuals with and without the "dB\_Sun" correction: in fact, at least a part of the discrepancies found in the measurements between ASM and VFM are modelled through a stray field (dB\_Sun) that is a function of the orientation of the VFM wrt Sun.
- Comparison of magnetic data ( $B_{NEC}$ ) with a model (Chaos7).

#### 3.3.1 VFM-ASM anomaly

- S/C A – violation of:
  - VFM-ASM residuals threshold on 21/11, 22/11, 23/11, 24/11, 25/11, 26/11, 27/11;
  - standard deviation of residuals threshold on 21/11, 22/11, 23/11, 24/11, 25/11, 26/11, 27/11.
- S/C B – no violation of VFM-ASM residual thresholds.

##### 3.3.1.1 ASM-VFM difference statistics

The ASM-VFM difference is defined as follow:

$$dF = |B_{VFM}| - F_{ASM}$$

Figure 3-3 and Figure 3-4 show the daily mean (circles) and standard deviation (crosses) of  $dF$  of the last month for Swarm A and Swarm B respectively.



Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 11 of 22




Figure 3-3: Daily mean and standard deviation values of ASM-VFM residuals (defined as  $dF = |B_{VFM}| - F_{ASM}$ ) for S/C A.




Figure 3-4: Daily mean and standard deviation values of ASM-VFM residuals (defined as  $dF = |B_{VFM}| - F_{ASM}$ ) for S/C B.

3.3.1.2 Swarm A

The daily peak-to-peak difference around the week stays within  $[-27.61 - 25.45]$  nT. Below follow two plots of such differences for current week (Figure 3-5).




Figure 3-5: ASM-VFM residuals for S/C A, during monitoring period 21/11-27/11. In top figure are plotted: difference between  $|B_{VFM}|$  and  $F_{ASM}$  (without  $dB_{Sun}$  correction) (blue colour), and the residuals with  $dB_{Sun}$  corrections (red colour). In bottom figure residuals are presented on the world map.


## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 12 of 22

### 3.3.1.3 Swarm B

The daily peak-to-peak difference around the week stays within  $[-0.41 - 0.6]$  nT. Below follow two plots of such differences for current week (Figure 3-6).



**Figure 3-6:** ASM-VFM residuals for S/C B, during monitoring period 21/11-27/11. In top figure are plotted: difference between  $|B_{VFM}|$  and  $F_{ASM}$  (without  $dB_{Sun}$  correction) (blue colour), and the residuals with  $dB_{Sun}$  corrections (red colour). In bottom figure residuals are presented on the world map.

### 3.3.1.4 Swarm C

No data because ASM is switched off.

## 3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC\_0)

For S/C A and B, the temperature and quartz frequency behaved as expected.

## 3.3.3 VFM Instrument parameters: VFM temperatures (MAG\_CA)


The VFM instrument parameters important for monitoring the instrument health are the VFM sensor temperatures:  $T_{CDC}$ ,  $T_{CSC}$  and  $T_{EU}$ .

For S/C A, B and C, for reported period, the temperatures behaved as expected.

### 3.3.4 Magnetic time series visual inspection

#### 3.3.4.1 Swarm A

Map plots of magnetic field measurement for week 47 for S/C A can be seen in Figure 3-7 below.



**Figure 3-7:** S/C A, world map plots of the geomagnetic field and components measured during monitoring period 21/11-27/11. From top to bottom: F-magnetic field from ASM measurement,  $B_{NEC}$  components (North, East, and Centre) of magnetic field from VFM measurement.


Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 14 of 22

3.3.4.2 Swarm B

Map plots of magnetic field measurement for week 47 for S/C B can be seen in Figure 3-8 below.



**Figure 3-8:** S/C B, world map plots of the geomagnetic field and components measured during monitoring period 21/11-27/11. From top to bottom: F-magnetic field from ASM measurement,  $B_{NEC}$  components (North, East, and Centre) of magnetic field from VFM measurement.


## Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 15 of 22

### 3.3.4.3 Swarm C

Map plots of magnetic field measurement for week 47 for S/C C can be seen in Figure 3-9.



**Figure 3-9:** S/C C, world map plots of the geomagnetic field and components measured during monitoring period 21/11-27/11. From top to bottom:  $B_{NEC}$  components (North, East, and Centre) of magnetic field from VFM measurement.

### 3.3.5 $B_{NEC}$ vs Chaos7 model residuals

The magnetic field measurement is compared to magnetic field estimated from the Chaos7 global geomagnetic field model (only Core and Crustal contributions). Currently in the monitoring routines the external contribution based on Dst index is not taken into account.

Left side of Figure 3-10, Figure 3-11 and Figure 3-12 show field residuals  $\Delta B = B_{NEC} - B_{Chaos}$  (all versus co-latitude in degrees), from top to bottom: 1)  $B_r$ , 2)  $B_\theta$  and 3)  $B_\phi$ .

As a general feature one can see the field residuals to be steady and usually below 50 nT at low and middle latitudes, up to  $|55| - |60|$  degrees; then the residual increases at high latitudes because the Chaos model does not take into account the contribution from the external field ([RD. 04]).

Right side of Figure 3-10, Figure 3-11 and Figure 3-12 show, from top to bottom, the time series on first day of the week of: (1-2-3) residuals of  $B_{NEC} - B_{Chaos}$  by components, related to S/C A, B and C respectively.

The component most affected by residual spikes and variations is  $B_{\theta_{NEC}}$ , i.e. the component that shows the variations of the field wrt to co-latitude. At high latitudes, the order of magnitude of the variability is about  $\pm 200$  nT.



Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 16 of 22

3.3.5.1 Swarm A




Figure 3-10: S/C A day 21/11: time series of  $B_{NEC} - B_{Chaos}$  residuals (right) and  $B_{NEC} - B_{Chaos}$  vs colatitude (left).

3.3.5.2 Swarm B




Figure 3-11: S/C B day 21/11: time series of  $B_{NEC} - B_{Chaos}$  residuals (right) and  $B_{NEC} - B_{Chaos}$  vs colatitude (left).





Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 17 of 22

3.3.5.3 Swarm C




Figure 3-12: S/C C day 21/11: time series of  $B_{NEC} - B_{Chaos}$  residuals (right) and  $B_{NEC} - B_{Chaos}$  vs colatitude (left)

3.4 Plasma Products

The monitored plasma products are the electron density (Ne) and electron temperature (Te) measured by the EFI-LP instruments. The monitoring of the data is done on different temporal basis (daily, weekly, monthly, yearly) in order to have a comprehensive view on the data quality. Here we report only two examples of the performed data monitoring, which are the most representative of the data quality.

Figures from Figure 3-13 to Figure 3-15 show the weekly profiles of the electron density and temperature as a function of time for the last week of operations. Data have been down sampled from 0.5s to 2min in order to have a clearer representation (grey lines). Also, the 20 minutes moving window average is shown in the figures (black points). From these figures, it is possible to see if there are measurements with large discrepancies from the average behaviour, and their time location. Information on the local magnetic time is reported in the captions.

Figures from Figure 3-16 to Figure 3-18 show the variations of the electron density and temperature as a function of the latitudes in quasi-dipole (QD) coordinate system, during the last week of operation. These analyses are useful to study the dependence of the variables on the QD magnetic coordinate system.

These analyses are shown for the ascending (upper panels) and descending (bottom panels) phase orbits, separately.

It is visible that sometimes the electron temperature reaches very high values, exceeding ten thousand Kelvin, particularly at high latitudes. The nature of this feature is currently under investigation.




Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 18 of 22


### 3.4.1 Plasma time series visual inspection

#### 3.4.1.1 Swarm A



**Figure 3-13:** The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 7 a.m. for ascending phase and 7 p.m. for descending phase.

#### 3.4.1.2 Swarm B



**Figure 3-14:** The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 9 a.m. for descending phase and 9 p.m. for ascending phase.




Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 19 of 22


3.4.1.3 Swarm C



**Figure 3-15:** The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 7 a.m. for ascending phase and 7 p.m. for descending phase.

3.4.2 Plasma products latitudinal variations

3.4.2.1 Swarm A



**Figure 3-16:** The panels shown the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.




Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1


Page 20 of 22

3.4.2.2 Swarm B



**Figure 3-17:** The panels shown the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.

3.4.2.3 Swarm C



**Figure 3-18:** The panels shown the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.



Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 21 of 22

---

## 4. Special Investigations

Nothing to report.



**Swarm DISC Weekly Report**

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 22 of 22

---

End of Document