
Swarm DISC Weekly Report 2023/42: 2023/10/16 - 2023/10/22



Abstract : This is the **Swarm Data Innovation and Science Cluster** (Swarm DISC) Weekly report on Swarm products quality, covering the period from 16 October to 22 October 2023.

Doc. No : SW-RP-SER-GS-010

Author :	Approval :
Đorđe Stevanović, Nicola Comparetti and Maria E Mazzocato on behalf of Swarm DISC Team	Enkelejda Qamili ESA – EOP/GMQ

Distribution :	
ESA/ESRIN EOP-GMQ ESA/ESRIN EOP-GM Swarm MM Swarm DISC Management Team Swarm DISC subcontractors	ESA/ESTEC Swarm PLSO ESA/ESOC Swarm FOS

TABLE OF CONTENTS

TABLE OF CONTENTS	2
1. INTRODUCTION	4
1.1 Annual Swarm Data Quality Workshop	4
1.2 Current Operational configuration of monitored data:	5
1.3 Recent evolutions:	5
1.4 Reference documents	6
2. SUMMARY OF THE OBSERVATIONS	7
2.1 Changes in the general status of Swarm instruments and Level 1B products quality	7
2.2 Relevant observations of the week 42 (16/10 - 22/10)	7
3. ROUTINE QUALITY CONTROL	8
3.1 Gaps analysis	8
3.2 Orbit and Attitude Products	8
3.2.1 Position Statistics	8
3.2.2 Attitude observations	9
3.2.2.1 Swarm A	9
3.2.2.2 Swarm B	9
3.2.2.3 Swarm C	9
3.3 Magnetic Products	9
3.3.1 VFM-ASM anomaly	10
3.3.1.1 ASM-VFM difference statistics	10
3.3.1.2 Swarm A	11
3.3.1.3 Swarm B	12
3.3.1.4 Swarm C	12
3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC_0)	12
3.3.3 VFM Instrument parameters: VFM temperatures (MAG_CA)	12
3.3.4 Magnetic time series visual inspection	13
3.3.4.1 Swarm A	13
3.3.4.2 Swarm B	14
3.3.4.3 Swarm C	15
3.3.5 B_{NEC} vs Chaos7 model residuals	15
3.3.5.1 Swarm A	16
3.3.5.2 Swarm B	16
3.3.5.3 Swarm C	17
3.4 Plasma Products	17
3.4.1 Plasma time series visual inspection	18
3.4.1.1 Swarm A	18
3.4.1.2 Swarm B	18
3.4.1.3 Swarm C	19
3.4.2 Plasma products latitudinal variations	19
3.4.2.1 Swarm A	19
3.4.2.2 Swarm B	20
3.4.2.3 Swarm C	20
4. SPECIAL INVESTIGATIONS	21

AMENDMENT POLICY

This document shall be amended by releasing a new edition of the document in its entirety. The Amendment Record Sheet below records the history and issue status of this document.

AMENDMENT RECORD SHEET

ISSUE	DATE	REASON
1.0	07 Dec 2023	First issue

1. Introduction

This document refers to the activities carried out in the framework of the ESA Sensor Performance, Products and Algorithms (SPPA) Office [RD. 01].

Chapter 1 gives an overview on the outcomes from the annual Swarm Data Quality Workshop and reports the information on the current operational configuration and its future improvements. It also contains the list of used reference documents.

In Chapter 2, the Section 2.1 gives an overview of the general quality status of the Swarm mission instruments and products, while the main observations of the week are summarized in the Section 2.2.

The document also includes information on data quality for the three Swarm spacecraft, inferred from automated HTML quality reports, which are produced on daily basis for each product. If interested in accessing the reports via web or FTP, please contact the Swarm DISC team at the following email address: <swarm@eo-sppa.org>. Such data quality reports represent the main component of the Routine Quality Control performed by ESA SPPA (Chapter 3). A description of the implemented quality checks is given in [RD. 02], and references therein.

Based on specific findings of the routine quality control, or requests from other entities (i.e. Swarm Payload Data Ground Segment (PDGS), Flight Operation Segment (FOS), Mission Management, Post-Launch Support Office (PLSO), Expert Support Laboratories (ESL), Quality Working Groups (QWG), and user community), investigations on anomalies can be triggered. Preliminary characterisations on such anomalies are given in Chapter 4.

Information on Swarm Level 1B products can be found in [RD. 03].

This weekly report is based on QC methods and diagnostics that tend to be continuously evolved and improved throughout the mission lifetime, reporting on the data quality, product evolutions, and status of the instruments on weekly basis.

1.1 Annual Swarm Data Quality Workshop

The 13th Swarm Data Quality Workshop was held on 10-11-12 October 2023, hosted by the European Space Agency facility in Italy (ESA -ESRIN), located in Frascati. Thanks to the participation of more than 120 scientists and instruments' experts (in presence and remotely) from different institutions in Europe, America and Asia, the event has been a great success.

The DQW#13 has been a 3 days' workshop dedicated to Swarm Data Calibration and Validation topics.

All the topics related to Swarm-based science and applications, which used to be included in the full week DQW event, will find the due space and audience during the next Swarm 10th Anniversary and Science Meeting that will be held from 8 to 12 April 2024 (<https://www.swarm-anniversary-and-science.org/>), organized by ESA (with the support of Swarm Data Quality team) and hosted by DTU Space (Copenhagen, Denmark).

The DQW#13, structured in 7 thematic sessions including talks and dedicated time slots for discussions, was instrumental in addressing the processing and use of Swarm data and defining a road map for future activities.

The main topics addressed during the workshop were related to:

collecting feedback about Swarm L1B and L2-Cat2 data quality, targeting new processing algorithms and correction improvements;

Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 5 of 22

Collecting innovative ideas for future Swarm-based activities, emerging applications, products, and services;
 promoting synergies with other satellite missions;
 discussing on future orbital constellation evolution.

More information on the Swarm DQW#13 outcomes, summary and recommendations will be published soon at <https://swarmdisc.org/>.

1.2 Current Operational configuration of monitored data:

Processor	
Name	Version
L1BOP	v3.24p4
L2-Cat2	v01.20
Products	
Name	Baseline
L0 inputs	02
L1B MAGNET and PLASMA	06
L1B ORBATT and ACCELE	05
L2-Cat2 EEF	02
L2-Cat2 IBI, FAC and TEC	04
Others	
Input auxiliary files	S/C A, CCDB 0029 (28/08/2023) S/C B, CCDB 0030 (28/08/2023) S/C C, CCDB 0030 (28/08/2023)
MPPF-CVQ	v03.13 (18/03/2022)

1.3 Recent evolutions:

On 24/06/2023 an improved version (v3.24p5) of L1B Operational Processors was transferred into operation in order to fix two minor issues detected on the clock offset information stored inside the Position dataset of MODx_SC_1B operational product.

This patch contained an updated library that affected the computation of the quaternions describing the transformation between the STR and the NEC reference frames, causing a degradation of attitude and magnetic measurements with errors close to 10 arcminutes in attitude and up to 150 nT in B_NEC. In order to restore the nominal data quality, a rollback from L1BOP 3.24p5 to 3.24p4 has been performed on 25 August 2023.

In addition, a further degradation of attitude (errors up to 8 arcseconds) and magnet (errors up to 3 nT in B_NEC) data has been identified in the data covering from October/December 2018 onward, when the STR were commanded to operate at 2 Hz sampling rate. Such degradation was introduced during the last full reprocessing when ORBATT and MAGNET data have been processed with the STR_q_CHU parameters of AUXxSW1_C_ CCDB file corresponding to 1 Hz sampling rate.

This issue was fixed with an updated set of AUXxSW1_C_ CCDB files deployed in operation on 25 August 2023.

Together with the rollback to 3.24p4, this action restored the nominal attitude and magnetic data quality from 25 August 2023 onwards.

Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 6 of 22

To recover the entire ORBATT and MAGNET quality a reprocessing campaign of the following time periods will be performed soon:

- Swarm Alpha: from 12 December 2018 to 24 August 2023
- Swarm Bravo: from 17 October 2018 to 24 August 2023
- Swarm Charlie: from 18 December 2018 to 24 August 2023

1.4 Reference documents

The following is a list of documents with a direct bearing on the content of this report. Where referenced in the text, these are identified as RD.n, where 'n' is the number in the list below:

- [RD. 01] Sensor Performance, Products and Algorithms (SPPA), PGSI-GSOP-EOPG-TN-05-0025. Version 2.3.
- [RD. 02] Swarm MPPF-CVQ Monitoring Baseline Document, ST-ESA-SWARM-MBD-0001, Issue 1.7.
- [RD. 03] Swarm Level 1B Product Definition, SW-RS-DSC-SY-0007, Issue 5.23.
- [RD. 04] Olsen, N., H. Luhr, C.C. Finlay, T.J. Sabaka, I. Michaelis, J. Rauberg and L. Tøffner-Clausen, The CHAOS-4 geomagnetic field model, *Geophys. J. Int.* 197, 815–827, 2014
- [RD. 05] <https://earth.esa.int/documents/10174/1514862/Swarm-L1B-and-L2-operational-processors.pdf>

2. Summary of the observations

2.1 Changes in the general status of Swarm instruments and Level 1B products quality

- The Absolute Scalar Magnetometer (ASM) on board Swarm Bravo is commanded in burst mode from 21/10/2023 at 00:00 UTC to 28/10/2023 at 00:00 UTC.
- The fourth and last manoeuvre of the orbit raise campaign was executed on Swarm Bravo on 19/10/2023 starting at 02:40 until 05:49 UTC . The thrusting phase lasted for two consecutive revolutions. It is the last of four manoeuvres to increase the orbital altitude of Swarm-B with respect to the lower pair. The expected final delta mean altitude with respect to Swarm-A/C is ~18 km at the end of the campaign. Consequently, the quality of magnetic data was affected during the manoeuvre (as showed in the Figure 3-6)

2.2 Relevant observations of the week 42 (16/10 - 22/10)

During the monitored week no events have been found.

3. Routine Quality control

3.1 Gaps analysis

- 20th -21st October:

F_ASM data stored in MAGx_LR_1B for Swarm Bravo is set to zero from 20/10/2023 at 23:59:29 to 21/10/2023 at 00:00:06 UTC. A data gap in MAGx_CA_1B production is present in the same time frame. This gap is related to the switch of the ASM to Burst/Vector Mode (see section 2.1).

3.2 Orbit and Attitude Products

The relevant parameters that have been monitored are:

- Position difference between calculated Medium Accuracy orbits (**MODx_SC_1B**) and on-board solution (**GPSxNAV_0**). Threshold values for such differences have not been assessed yet: we have just monitored the average values and maximum variations within the week. For the time being we evaluated an anomaly should be raised if one (or more) of the following conditions occurs:
 - The **average difference** on a given day exceeds the position accuracy requirement for the mission (1.5 m),
 - The variability around the average is quite high: **standard deviation** threshold has been arbitrarily chosen to be twice the position accuracy requirement for the mission (2-sigma = 3 m).
 - At least 4-5 spikes are observed on a given day, exceeding +/- 50 m.
- Visual inspection of Star Tracker characterisation flags (**STRxATT_1B**)
- Deviation of the quaternion norm from unity (deviation threshold = +/- 10⁻⁹)
- Visual inspection of Euler Angles derived from quaternions.

3.2.1 Position Statistics

In Figure 3-1 and Figure 3-2 one can see the statistics of the differences between MOD and on-board solution positions for S/C A, B and C respectively. Figure 3-1 shows a cumulative trend of the maximum daily standard deviation for the past 30 days of operations of the MOD-NAV difference, while Figure 3-2 shows the daily maximum difference, in absolute value, of the MOD-NAV difference, always for the past 30 days of operations.




Figure 3-1: Plot of the standard deviation of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.




Figure 3-2: Plot of the maximum difference of the absolute value of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.

3.2.2 Attitude observations

3.2.2.1 Swarm A

Nominal. Nothing to report.

3.2.2.2 Swarm B

Nominal. Nothing to report.

3.2.2.3 Swarm C

Nominal. Nothing to report.

3.3 Magnetic Products

For the magnetic products, the weekly monitoring consists of:

Swarm DISC Weekly Report

- ASM instrument monitoring: quartz frequency (nominal range: [2.949E7 – 2.950E7] Hz) and ASM temperature (temperature range shall be: [-30;+50] °C, Rel. Variation shall not exceed: 0.1 °C/sec).
- VFM instrument monitoring: temperatures (Rel. Variation shall not exceed: 0.1 °C/sec).
- Visual inspection of daily time series of magnetic field intensity F , B_{NEC} and B_{VFM} . Looking for gaps (or zero values in case of MAG_{LR_1B} products), out-of-threshold values (i.e. exceeding +/- 60000 nT), and other strange features. Map plots of F and B_{NEC} for the whole week are then displayed.
- Monitoring of the ASM-VFM known anomaly: visual inspection of $|B_{VFM}| - F$ taken from MAG_{CA_1B} products and recording of daily maximum variations and standard deviations. If +/- 1 nT are exceeded on a given day, an alert is raised. Map plots of the residuals are shown along with weekly time series of the residuals with and without the "dB_Sun" correction: in fact, at least a part of the discrepancies found in the measurements between ASM and VFM are modelled through a stray field (dB_Sun) that is a function of the orientation of the VFM wrt Sun.
- Comparison of magnetic data (B_{NEC}) with a model (Chaos7).

3.3.1 VFM-ASM anomaly

- S/C A – no violation of residuals threshold.
- S/C B – violation of:
 - o VFM-ASM residuals threshold on 19/10, 22/10;
 - o mean value of residuals threshold on 21/10, 22/10.

3.3.1.1 ASM-VFM difference statistics

The ASM-VFM difference is defined as follow:

$$dF = |B_{VFM}| - F_{ASM}$$

Figure 3-3 and Figure 3-4 show the daily mean (circles) and standard deviation (crosses) of dF of the last month for Swarm A and Swarm B respectively.




Figure 3-3: Daily mean and standard deviation values of ASM-VFM residuals (defined as $dF = |B_{VFM}| - F_{ASM}$) for S/C A.




Figure 3-4: Daily mean and standard deviation values of ASM-VFM residuals (defined as $dF=|B_{VFM}|-F_{ASM}$) for S/C B.

3.3.1.2 Swarm A

The daily peak-to-peak difference around the week stays within $[-0.99 - 0.66]$ nT. Below follow two plots of such differences for current week (Figure 3-5).




Figure 3-5: ASM-VFM residuals for S/C A, during monitoring period 16/10-22/10. In top figure are plotted: difference between $|B_{VFM}|$ and F_{ASM} (without dB_{Sun} correction) (blue colour), and the residuals with dB_{Sun} corrections (red colour). In bottom figure residuals are presented on the world map.

3.3.1.3 Swarm B

The daily peak-to-peak difference around the week stays within [-1.04 - 1.51] nT. Below follow two plots of such differences for current week (Figure 3-6).




Figure 3-6: ASM-VFM residuals for S/C B, during monitoring period 16/10-22/10. In top figure are plotted: difference between $|B_{VFM}|$ and F_{ASM} (without dB_{Sun} correction) (blue colour), and the residuals with dB_{Sun} corrections (red colour). In bottom figure residuals are presented on the world map.

3.3.1.4 Swarm C

No data because ASM is switched off.

3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC_0)

For S/C A and B, the temperature and quartz frequency behaved as expected.

3.3.3 VFM Instrument parameters: VFM temperatures (MAG_CA)

The VFM instrument parameters important for monitoring the instrument health are the VFM sensor temperatures: T_{CDC} , T_{CSC} and T_{EU} .

For S/C A, B and C, for reported period, the temperatures behaved as expected.

3.3.4 Magnetic time series visual inspection

3.3.4.1 Swarm A

Map plots of magnetic field measurement for week 42 for S/C A can be seen in Figure 3-7 below.




Figure 3-7: S/C A, world map plots of the geomagnetic field and components measured during monitoring period 16/10-22/10. From top to bottom: F-magnetic field from ASM measurement, B_{NEC} components (North, East, and Centre) of magnetic field from VFM measurement.

3.3.4.2 Swarm B

Map plots of magnetic field measurement for week 42 for S/C B can be seen in Figure 3-8 below.




Figure 3-8: S/C B, world map plots of the geomagnetic field and components measured during monitoring period 16/10-22/10. From top to bottom: F-magnetic field from ASM measurement, B_{NEC} components (North, East, and Centre) of magnetic field from VFM measurement.

3.3.4.3 Swarm C

Map plots of magnetic field measurement for week 42 for S/C C can be seen in Figure 3-9.




Figure 3-9: S/C C, world map plots of the geomagnetic field and components measured during monitoring period 16/10-22/10. From top to bottom: B_{NEC} components (North, East, and Centre) of magnetic field from VFM measurement.

3.3.5 B_{NEC} vs Chaos7 model residuals

The magnetic field measurement is compared to magnetic field estimated from the Chaos7 global geomagnetic field model (only Core and Crustal contributions). Currently in the monitoring routines the external contribution based on Dst index is not taken into account.

Left side of Figure 3-10, Figure 3-11 and Figure 3-12 show field residuals $\Delta B = B_{NEC} - B_{Chaos7}$ (all versus co-latitude in degrees), from top to bottom: 1) B_r , 2) B_θ and 3) B_ϕ .

As a general feature one can see the field residuals to be steady and usually below 50 nT at low and middle latitudes, up to $|\pm 55| - |\pm 60|$ degrees; then the residual increases at high latitudes because the Chaos model does not take into account the contribution from the external field ([RD. 04]).

Right side of Figure 3-10, Figure 3-11 and Figure 3-12 show, from top to bottom, the time series on first day of the week of: (1-2-3) residuals of $B_{NEC} - B_{Chaos}$ by components, related to S/C A, B and C respectively.

The component most affected by residual spikes and variations is $B_{\theta_{NEC}}$, i.e. the component that shows the variations of the field wrt to co-latitude. At high latitudes, the order of magnitude of the variability is about +/- 200 nT.

3.3.5.1 Swarm A




Figure 3-10: S/C A day 16/10: time series of $B_{NEC} - B_{Chaos}$ residuals (right) and $B_{NEC} - B_{Chaos}$ vs colatitude (left).

3.3.5.2 Swarm B




Figure 3-11: S/C B day 16/10: time series of $B_{NEC} - B_{Chaos}$ residuals (right) and $B_{NEC} - B_{Chaos}$ vs colatitude (left).

3.3.5.3 Swarm C




Figure 3-12: S/C C day 16/10: time series of $B_{NEC} - B_{Chaos}$ residuals (right) and $B_{NEC} - B_{Chaos}$ vs colatitude (left)

3.4 Plasma Products

The monitored plasma products are the electron density (Ne) and electron temperature (Te) measured by the EFI-LP instruments. The monitoring of the data is done on different temporal basis (daily, weekly, monthly, yearly) in order to have a comprehensive view on the data quality. Here we report only two examples of the performed data monitoring, which are the most representative of the data quality.

Figures from Figure 3-13 to Figure 3-15 show the weekly profiles of the electron density and temperature as a function of time for the last week of operations. Data have been down sampled from 0.5s to 2min in order to have a clearer representation (grey lines). Also, the 20 minutes moving window average is shown in the figures (black points). From these figures, it is possible to see if there are measurements with large discrepancies from the average behaviour, and their time location. Information on the local magnetic time is reported in the captions.

Figures from Figure 3-16 to Figure 3-18 show the variations of the electron density and temperature as a function of the latitudes in quasi-dipole (QD) coordinate system, during the last week of operation. These analyses are useful to study the dependence of the variables on the QD magnetic coordinate system.

These analyses are shown for the ascending (upper panels) and descending (bottom panels) phase orbits, separately.

It is visible that sometimes the electron temperature reaches very high values, exceeding ten thousand Kelvin, particularly at high latitudes. The nature of this feature is currently under investigation.

3.4.1 Plasma time series visual inspection

3.4.1.1 Swarm A




Figure 3-13: The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 1:30 a.m. for ascending phase and 1:30 p.m. for descending phase.

3.4.1.2 Swarm B




Figure 3-14: The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 4:30 a.m. for descending phase and 4:30 p.m. for ascending phase.

Swarm DISC Weekly Report

Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 19 of 22

3.4.1.3 Swarm C




Figure 3-15: The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 1:30 a.m. for ascending phase and 1:30 p.m. for descending phase.

3.4.2 Plasma products latitudinal variations

3.4.2.1 Swarm A




Figure 3-16: The panels show the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.

3.4.2.2 Swarm B




Figure 3-17: The panels shown the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.

3.4.2.3 Swarm C




Figure 3-18: The panels shown the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.

4. Special Investigations

Nothing to report.

End of Document