






# Swarm DISC Weekly Report 2024/13: 2024/03/25 - 2024/03/31



Abstract This is the Swarm Data Innovation and Science Cluster (Swarm DISC) Weekly report on

Swarm products quality, covering the period from 25 March to 31 March 2024.

Doc. No SW-RP-SER-GS-010

Author Approval

> Enkelejda Qamili Đorđe Stevanović, Nicola Comparetti and Maria E. ESA - EOP/GMQ

Mazzocato on behalf of Swarm **DISC Team** 

Distribution **ESA/ESRIN EOP-GMQ** 

**ESA/ESTEC Swarm PLSO** ESA/ESRIN EOP-GM Swarm MM ESA/ESOC Swarm FOS

Swarm DISC Management Team Swarm DISC subcontractors

© Serco S.p.a., Italy, 2024. Proprietary and intellectual rights of Serco S.p.a., Italy are involved in the subject-matter of this material and all manufacturing reproduction, use, disclosure, and sales rights pertaining to such subject-matter are expressly reserved. This material is submitted for a specific purpose as agreed in writing, and the recipient by accepting this material agrees that this material will not be used, copied, or reproduced in whole or in part nor its contents (or any part the reof) revealed in any manner or to any third party, except own staff, to meet the purpose for which it was submitted and subject to the terms of the written agreement.







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 2 of 23

## **TABLE OF CONTENTS**

TA	TABLE OF CONTENTS				
1.	INTRODUCTION	4			
1.1	1 Annual Swarm Data Quality Workshop				
1.2	.2 Current Operational configuration of monitored data:				
1.3					
1.4	Reference documents	6			
2.	SUMMARY OF THE OBSERVATIONS	7			
2.1					
2.2					
3.	ROUTINE QUALITY CONTROL				
3.1 3.2					
5.2	3.2.1 Position Statistics				
	3.2.2 Attitude observations				
	3.2.2.1 Swarm A				
	3.2.2.2 Swarm B				
	3.2.2.3 Swarm C				
3.3					
5.5	3.3.1 VFM-ASM a nomaly				
	3.3.1.1 ASM-VFM difference statistics				
	3.3.1.2 Swarm A				
	3.3.1.3 Swarm B				
	3.3.1.4 Swarm C				
	3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC_0)	12			
	3.3.3 VFM Instrument parameters: VFM temperatures (MAG_CA)				
	3.3.4 Magnetic time series visual inspection				
	3.3.4.1 Swarm A	13			
	3.3.4.2 Swarm B	14			
	3.3.4.3 Swarm C				
	3.3.5 BNEC vs Chaos7 model residuals				
	3.3.5.1 Swarm A				
	3.3.5.2 Swarm B				
	3.3.5.3 Swarm C				
3.4					
	3.4.1 Plasma time series visual inspection				
	3.4.1.1 Swarm A				
	3.4.1.2 Swarm B				
	3.4.1.3 Swarm C				
	3.4.2 Plasma products latitudinal variations				
	3.4.2.1 Swarm A				
	3.4.2.2 Swarm B				
	3.4.2.3 Swarm C	20			
4	SDECIAL INVESTIGATIONS	21			







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 3 of 23

## **AMENDMENT POLICY**

This document shall be amended by releasing a new edition of the document in its entirety. The Amendment Record Sheet below records the history and issue status of this document.

#### **AMENDMENT RECORD SHEET**

ISSUE	DATE	REASON
1.0	14 Jun 2024	First issue







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 4 of 23

## 1. Introduction

This document refers to the activities carried out in the framework of the ESA Sensor Performance, Products and Algorithms (SPPA) Office [RD. 01].

Chapter 1 gives an overview on the outcomes from the annual Swarm Data Quality Workshop and reports the information on the current operational configuration and its future improvements. It also contains the list of used reference documents.

In Chapter 2, the Section 2.1 gives an overview of the general quality status of the Swarm mission instruments and products, while the main observations of the week are summarized in the Section 2.2.

The document also includes information on data quality for the three Swarm spacecraft, inferred from automated HTML quality reports, which are produced on daily basis for each product. If interested in accessing the reports via web or FTP, please contact the Swarm DISC team at the following email address: <swarm@eo-sppa.org>. Such data quality reports represent the main component of the Routine Quality Control performed by ESA SPPA (Chapter 3). A description of the implemented quality checks is given in [RD. 02], and references therein.

Based on specific findings of the routine quality control, or requests from other entities (i.e. Swam Payload Data Ground Segment (PDGS), Flight Operation Segment (FOS), Mission Management, Post-Launch Support Office (PLSO), Expert Support Laboratories (ESL), Quality Working Groups (QWG), and user community), investigations on anomalies can be triggered. Preliminary characterisations on such anomalies are given in Chapter 4.

Information on Swarm Level 1B products can be found in [RD. 03].

This weekly report is based on QC methods and diagnostics that tend to be continuously evolved and improved throughout the mission lifetime, reporting on the data quality, product evolutions, and status of the instruments on weekly basis.

## 1.1 Annual Swarm Data Quality Workshop

The 13th Swarm Data Quality Workshop was held on 10-11-12 October 2023, hosted by the European Space Agency facility in Italy (ESA -ESRIN), located in Frascati. Thanks to the participation of more than 120 scientists and instruments' experts (in presence and remotely) from different institutions in Europe, America and Asia, the event has been a great success.

The DQW#13 has been a 3 days' workshop dedicated to Swarm Data Calibration and Validation topics.

All the topics related to Swarm-based science and applications, which used to be included in the full week DQW event, will find the due space and audience during the next Swarm 10th Anniversary and Science Meeting that will be held from 8 to 12 April 2024 (<a href="https://www.swarm-anniversary-and-science.org">https://www.swarm-anniversary-and-science.org</a>), organized by ESA (with the support of Swarm Data Quality team) and hosted by DTU Space (Copenhagen, Denmark).

The DQW#13, structured in 7 thematic sessions including talks and dedicated time slots for discussions, was instrumental in addressing the processing and use of Swarm data and defining a road map for future activities.

The main topics addressed during the workshop were related to:

- collecting feedback about Swarm L1B and L2-Cat2 data quality, targeting new processing algorithms and correction improvements,
- collecting innovative ideas for future Swarm-based activities, emerging applications, products, and services,







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 5 of 23

- promoting synergies with other satellite missions,
- discussing on future orbital constellation evolution.

More information on the Swarm DQW#13 outcomes, summary and recommendations will be published soon at <a href="https://swarmdisc.org/">https://swarmdisc.org/</a>.

## 1.2 **Current Operational configuration of monitored data:**

Processor			
Name	Version		
L1BOP	v3.24p4		
L2-Cat2	v01.20		
Products			
Name	Baseline		
L0 inputs	02		
L1B MAGNET and PLASMA	06		
L1B ORBATT and ACCELE	05		
L2-Cat2 EEF	02		
L2-Cat2 IBI, FAC and TEC	04		
Others			
Input auxiliary files	S/C A, CCDB 0029 (28/08/2023)		
	S/C B, CCDB 0030 (28/08/2023)		
	S/C C, CCDB 0030 (28/08/2023)		
MPPF-CVQ	v03.13 (18/03/2022)		

#### 1.3 Recent evolutions:

On 24/06/2023 an improved version (v3.24p5) of L1B Operational Processors was transferred into operation to fix two minor issues detected on the clock offset information stored inside the Position dataset of MODx\_SC\_1B operational product.

This patch contained an updated library that affected the computation of the quaternions describing the transformation between the STR and the NEC reference frames, causing a degradation of attitude and magnetic measurements with errors close to 10 arcminutes in attitude and up to 150 nT in B\_NEC. In order to restore the nominal data quality, a rollback from L1BOP 3.24p5 to 3.24p4 has been performed on 25 August 2023.

In addition, a further degradation of attitude (errors up to 8 arcseconds) and magnet (errors up to 3 nT in B\_NEC) data has been identified in the data covering from October/December 2018 onward, when the STR were commanded to operate at 2 Hz sampling rate. Such degradation was introduced during the last full reprocessing when ORBATT and MAGNET data were processed with the STR\_q\_CHU parameters of the AUXxSW1\_C\_ CCDB file corresponding to a 1 Hz sampling rate.

This issue was fixed with an updated set of AUXxSW1\_C\_ CCDB files deployed in operation on 25 August 2023.

Together with the rollback to 3.24p4, this action restored the nominal attitude and magnetic data quality from 25 August 2023 onwards.







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 6 of 23

#### 1.4 Reference documents

The following is a list of documents with a direct bearing on the content of this report. Where referenced in the text, these are identified as RD.n, where 'n' is the number in the list below:

- [RD. 01] Sensor Performance, Products and Algorithms (SPPA), PGSI-GSOP-EOPG-TN-05-0025. Version 2.3.
- [RD. 02] Swarm MPPF-CVQ Monitoring Baseline Document, ST-ESA-SWARM-MBD-0001, Issue 1.7.
- [RD. 03] Swarm Level 1B Product Definition, SW-RS-DSC-SY-0007, Issue 5.23.
- [RD. 04] Olsen, N., H. Luhr, C.C. Finlay, T.J. Sabaka, I. Michaelis, J. Rauberg and L. Tøffner-Clausen, The CHAOS-4 geomagnetic field model, Geophys. J. Int. 197, 815–827, 2014
- [RD. 05] <a href="https://earth.esa.int/documents/10174/1514862/Swarm-L1B-and-L2-operational-processors.pdf">https://earth.esa.int/documents/10174/1514862/Swarm-L1B-and-L2-operational-processors.pdf</a>







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 7 of 23

## 2. Summary of the observations

# 2.1 Changes in the general status of Swarm instruments and Level 1B products quality

- The Absolute Scalar Magnetometer (ASM) on board Swarm Alpha is commanded in burst mode from 18/03/2024 at 00:00 UTC to 25/03/2024 at 00:00 UTC.
- The Absolute Scalar Magnetometer (ASM) on board Swarm Bravo is commanded in burst mode from 27/03/2024 at 00:00 UTC to 03/04/2024 at 00:00 UTC.
- On 30/03/2024 there was an anomaly on the Swarm-B telemetry files causing small data gaps (see Section 3.1).

## 2.2 Relevant observations of the week 13 (25/03 - 31/03)

During the monitored week no events have been found.







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 8 of 23

## 3. Routine Quality control

## 3.1 Gaps analysis

#### 25th March:

F\_ASM data stored in MAGx\_LR\_1B for Swarm Alpha is set to zero from 24/03/2024 at 23:59:28 to 25/03/2024 at 00:00:24 UTC. A data gap in MAGx\_CA\_1B production is present in the same time frame. This gap is related to the switch of the ASM to Burst/Vector Mo de (see section 2.1).

#### • 26-27th March:

F\_ASM data stored in MAGx\_LR\_1B for Swarm Bravo is set to zero from 26/03/2024 at 23:59:29 to 27/03/2024 at 00:00:06 UTC. A data gap in MAGx\_CA\_1B production is present in the same time frame. This gap is related to the switch of the ASM to Burst/Vector Mode (see section 2.1).

#### • 30th March:

Swarm-B two small (<10s) data gaps detected in telemetry files between 03:44 and 03:54 UTC. All instruments were affected.

## 3.2 **Orbit and Attitude Products**

The relevant parameters that have been monitored are:

- Position difference between calculated Medium Accuracy orbits (MODx\_SC\_1B) and on-board solution (GPSxNAV\_0). Threshold values for such differences have not been assessed yet: we have just monitored the average values and maximum variations within the week. For the time being we evaluated an anomaly should be raised if one (or more) of the following conditions occurs:
  - The average difference on a given day exceeds the position accuracy requirement for the mission (1.5 m),
  - The variability around the average is quite high: standard deviation threshold has been arbitrarily chosen to be twice the position accuracy requirement for the mission (2-sigma = 3 m).
  - At least 4-5 spikes are observed on a given day, exceeding +/- 50 m.
- Visual inspection of Star Tracker characterisation flags (STRxATT\_1B)
- Deviation of the quaternion norm from unity (deviation threshold =  $+/-10^{-9}$ )
- Visual inspection of Euler Angles derived from quaternions.









Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 9 of 23

## 3.2.1 **Position Statistics**

In Figure 3-1 and Figure 3-2 one can see the statistics of the differences between MOD and on-board solution positions for S/C A, B and C respectively. Figure 3-1 shows a cumulative trend of the maximum daily standard deviation for the past 30 days of operations of the MOD-NAV difference, while Figure 3-2 shows the daily maximum difference, in absolute value, of the MOD-NAV difference, always for the past 30 days of operations.



**Figure 3-1:** Plot of the standard deviation of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.



**Figure 3-2:** Plot of the maximum difference of the absolute value of the difference between MOD and NAV solutions for all satellites. Plot covers last month of operation.

#### 3.2.2 Attitude observations

## 3.2.2.1 **Swarm A**

Nominal. Nothing to report.

#### 3.2.2.2 **Swarm B**

Nominal. Nothing to report.







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 10 of 23

#### 3.2.2.3 **Swarm C**

Nominal. Nothing to report.

## 3.3 Magnetic Products

For the magnetic products, the weekly monitoring consists of:

- ASM instrument monitoring: quartz frequency (nominal range: [2.949E7 2.950E7] Hz) and ASM temperature (temperature range shall be: [-30;+50] °C, Rel. Variation shall not exceed: 0.1 °C/sec).
- VFM instrument monitoring: temperatures (Rel. Variation shall not exceed: 0.1 °C/sec).
- Visual inspection of daily time series of magnetic field intensity F, BNEC and BVFM. Looking for gaps (or zero values in case of MAGx\_LR\_1B products), out-of-threshold values (i.e. exceeding +/- 60000 nT), and other strange features. Map plots of F and BNEC for the whole week are then displayed.
- Monitoring of the ASM-VFM known anomaly: visual inspection of |BVFM| F taken from MAGX\_CA\_1B products and recording of daily maximum variations and standard deviations. If +/- 1 nT are exceed on a given day, an alert is raised. Map plots of the residuals are shown along with weekly time series of the residuals with and without the "dB\_Sun" correction: in fact, at least a part of the discrepancies found in the measurements between ASM and VFM are modelled through a stray field (dB\_Sun) that is a function of the orientation of the VFM wrt Sun.
- Comparison of magnetic data (BNEC) with a model (Chaos 7).

#### 3.3.1 **VFM-ASM anomaly**

- S/C A no violation of residuals threshold.
- S/C B violation of:
  - o VFM-ASM residuals threshold on 25/03, 26/03, 27/03, 28/03, 29/03, 30/03, 31/03;
  - o mean value of residuals threshold on 25/03, 26/03, 27/03, 28/03, 29/03, 30/03, 31/03.

#### 3.3.1.1 **ASM-VFM difference statistics**

The ASM-VFM difference is defined as follow:

dF = |BVFM| - FASM

Figure 3-3 and Figure 3-4 show the daily mean (circles) and standard deviation (crosses) of dF of the last month for Swarm A and Swarm B respectively.







Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 11 of 23




Figure 3-3: Daily mean and standard deviation values of ASM-VFM residuals (defined as dF=|BVFM|-FASM) for S/C A.





Figure 3-4: Daily mean and standard deviation values of ASM-VFM residuals (defined as dF=|BVFM|-FASM) for S/C B.

#### 3.3.1.2 **Swarm A**

The daily peak-to-peak difference around the week stays within [-0.6 - 0.79] nT. Below follow two plots of such differences for current week (Figure 3-5).



**Figure 3-5:** ASM-VFM residuals for S/C A, during monitoring period 25/03-31/03. In top figure are plotted: difference between |B\_VFM| and F\_ASM (without dB\_Sun correction) (blue colour), and the residuals with dB\_Sun corrections (red colour). In bottom figure residuals are presented on the world map.








Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 12 of 23

#### 3.3.1.3 **Swarm B**

The daily peak-to-peak difference around the week stays within [-0.22 - 1.14] nT. Below follow two plots of such differences for current week (Figure 3-6).



**Figure 3-6:** ASM-VFM residuals for S/C B, during monitoring period 25/03-31/03. In top figure are plotted: difference between |B\_VFM| and F\_ASM (without dB\_Sun correction) (blue colour), and the residuals with dB\_Sun corrections (red colour). In bottom figure residuals are presented on the world map.

#### 3.3.1.4 **Swarm C**

No data because ASM is switched off.

## 3.3.2 ASM Instrument parameters: quartz frequency and ASM temperature (ASMAVEC\_0)

For S/C A and B, the temperature and quartz frequency behaved as expected.

## 3.3.3 **VFM Instrument parameters: VFM temperatures (MAG\_CA)**

The VFM instrument parameters important for monitoring the instrument health are the VFM sensor temperatures: T\_CDC, T\_CSC and T\_EU.

For S/C A, B and C, for reported period, the temperatures behaved as expected.









Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 13 of 23


## 3.3.4 Magnetic time series visual inspection


#### 3.3.4.1 **Swarm A**

Map plots of magnetic field measurement for week 13 for S/C A can be seen in Figure 3-7 below.









**Figure 3-7:** S/C A, world map plots of the geomagnetic field and components measured during monitoring period 25/03-31/03. From top to bottom: F-magnetic field from ASM measurement, BNEC components (North, East, and Centre) of magnetic field from VFM measurement.










Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 14 of 23


#### 3.3.4.2 **Swarm B**

Map plots of magnetic field measurement for week 13 for S/C B can be seen in Figure 3-8 below.









**Figure 3-8:** S/C B, world map plots of the geomagnetic field and components measured during monitoring period 25/03-31/03. From top to bottom: F-magnetic field from ASM measurement, BNEC components (North, East, and Centre) of magnetic field from VFM measurement.










Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 15 of 23

#### 3.3.4.3 **Swarm C**

Map plots of magnetic field measurement for week 13 for S/C C can be seen in Figure 3-9.







**Figure 3-9:** S/C C, world map plots of the geomagnetic field and components measured during monitoring period 25/03-31/03. From top to bottom: BNEC components (North, East, and Centre) of magnetic field from VFM measurement.

## 3.3.5 **B**<sub>NEC</sub> vs Chaos7 model residuals

The magnetic field measurement is compared to magnetic field estimated from the Chaos 7 global geomagnetic field model (only Core and Crustal contributions). Currently in the monitoring routines the external contribution based on Dst index is not taken into account.

Left side of Figure 3-10, Figure 3-11 and Figure 3-12 show field residuals dB=BNEC - BChaos (all versus co-latitude in degrees), from top to bottom: 1) Br, 2) B $\theta$  and 3) B $\phi$ .

As a general feature one can see the field residuals to be steady and usually below 50 nT at low and middle latitudes, up to |55| - |60| degrees; then the residual increases at high latitudes because the Chaos model does not take into account the contribution from the external field ([RD. 04]).







Doc. no: SW-RP-SER-GS-010, Rev: 1

Page 16 of 23

Right side of Figure 3-10, Figure 3-11 and Figure 3-12 show, from top to bottom, the time series on first day of the week of: (1-2-3) residuals of BNEC-BChaos by components, related to S/C A, B and C respectively.

The component most affected by residual spikes and variations is  $B\theta NEC$ , i.e. the component that shows the variations of the field wrt to co-latitude. At high latitudes, the order of magnitude of the variability is about +/- 200 nT.

#### 3.3.5.1 **Swarm A**




Figure 3-10: S/C A day 25/03: time series of BNEC - Bchaos residuals (right) and BNEC - Bchaos vs colatitude (left).

### 3.3.5.2 **Swarm B**




Figure 3-11: S/C B day 25/03: time series of BNEC – BChaos residuals (right) and BNEC - BChaos vs colatitude (left).







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 17 of 23

#### 3.3.5.3 **Swarm C**




Figure 3-12: S/C C day 25/03: time series of BNEC - Bchaos residuals (right) and BNEC - Bchaos vs colatitude (left)

#### 3.4 Plasma Products

The monitored plasma products are the electron density (Ne) and electron temperature (Te) measured by the EFI-LP instruments. The monitoring of the data is done on different temporal basis (daily, weekly, monthly, yearly) in order to have a comprehensive view on the data quality. Here we report only two examples of the performed data monitoring, which are the most representative of the data quality.

Figures from Figure 3-13 to Figure 3-15 show the weekly profiles of the electron density and temperature as a function of time for the last week of operations. Data have been down sampled from 0.5s to 2min in order to have a clearer representation (grey lines). Also, the 20 minutes moving window average is shown in the figures (black points). From these figures, it is possible to see if there are measurements with large discrepancies from the average behaviour, and their time location. Information on the local magnetic time is reported in the captions.

Figures from Figure 3-16 to Figure 3-18 show the variations of the electron density and temperature as a function of the latitudes in quasi-dipole (QD) coordinate system, during the last week of operation. These analyses are useful to study the dependence of the variables on the QD magnetic coordinate system.

These analyses are shown for the ascending (upper panels) and descending (bottom panels) phase orbits, separately.

It is visible that sometimes the electron temperature reaches very high values, exceeding ten thousand Kelvin, particularly at high latitudes. The nature of this feature is currently under investigation.








Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 18 of 23


## 3.4.1 Plasma time series visual inspection

#### 3.4.1.1 **Swarm A**



**Figure 3-13:** The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 11 a.m. for ascending phase and 11 p.m. for descending phase.

#### 3.4.1.2 **Swarm B**



**Figure 3-14:** The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 2:30 p.m. for descending phase and 2:30 a.m. for ascending phase.








Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 19 of 23


#### 3.4.1.3 **Swarm C**



**Figure 3-15:** The panels show the electron density (left) and temperature (right) weekly time series (grey lines) together with the 20 min moving windows average (black lines). The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits. The average magnetic local time during the week is 11 a.m. for ascending phase and 11 p.m. for descending phase.

## 3.4.2 Plasma products latitudinal variations

#### 3.4.2.1 **Swarm A**



**Figure 3-16:** The panels shown the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.









Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 20 of 23

#### 3.4.2.2 **Swarm B**



**Figure 3-17:** The panels shown the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.

#### 3.4.2.3 **Swarm C**



**Figure 3-18:** The panels shown the electron density (left) and temperature (right) profile as a function of QD Latitudes for the last week of operation. The analysis is made separately for ascending (upper panels) and descending (bottom panels) orbits.







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 21 of 23

## 4. Special Investigations

Nothing to report.







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 22 of 23

**End of Document** 







Doc. no: SW-RP-SER-GS-010, Rev: 1 Page 23 of 23